

Characterizations for the parabolic nonsingular integral operator on parabolic generalized Orlicz-Morrey spaces

Mehriban N. Omarova *

Received: 12.04.2019 / Revised: 29.09.2019 / Accepted: 02.11.2019

Abstract. We show continuity in parabolic generalized Orlicz-Morrey spaces $M^{\Phi, \varphi}(\mathbb{D}_+^{n+1})$ of parabolic nonsingular integral operators. We shall give necessary and sufficient conditions for the boundedness of the parabolic nonsingular integral operator on parabolic generalized Orlicz-Morrey spaces $M^{\Phi, \varphi}(\mathbb{D}_+^{n+1})$.

Keywords. Parabolic generalized Orlicz-Morrey spaces; parabolic nonsingular integral.

Mathematics Subject Classification (2010): 42B20, 42B35.

1 Introduction and main results

The classical Morrey spaces were introduced by Morrey [23] to study the local behavior of solutions to second-order elliptic partial differential equations. Although such spaces allow to describe local properties of functions better than Lebesgue spaces, they have some unpleasant issues. It is well known that Morrey spaces are non separable and that the usual classes of nice functions are not dense in such spaces. Moreover, various Morrey spaces are defined in the process of study. Guliyev, Mizuhara and Nakai [10, 22, 24] introduced generalized Morrey spaces $M^{p, \varphi}(\mathbb{R}^n)$ (see, also [11, 12, 32]). Later, Guliyev [12] defined the generalized Morrey spaces $M^{p, \varphi}(\mathbb{R}^n)$ with normalized norm

$$\|f\|_{M^{p, \varphi}} \equiv \sup_{x \in \mathbb{R}^n, r > 0} \varphi(x, r)^{-1} |B(x, r)|^{-1/p} \|f\|_{L^p(B(x, r))},$$

where the function φ is a positive measurable function on $\mathbb{R}^n \times (0, \infty)$. Here and everywhere in the sequel $B(x, r)$ is the ball in \mathbb{R}^n of radius r centered at x and $|B(x, r)| = v_n r^n$ is its Lebesgue measure, where v_n is the volume of the unit ball in \mathbb{R}^n .

The Orlicz space were first introduced by Orlicz in [29, 30] as generalizations of Lebesgue spaces $L^p(\mathbb{R}^n)$. Since then, the theory of Orlicz spaces themselves has been well developed and the spaces have been widely used in probability, statistics, potential theory, partial differential equations, as well as harmonic analysis and some other fields of analysis.

In [5], the generalized Orlicz-Morrey space $M^{\Phi, \varphi}(\mathbb{R}^n)$ was introduced to unify Orlicz and generalized Morrey spaces. Other definitions of generalized Orlicz-Morrey spaces can

* Corresponding author

be found in [25] and [31]. In words of [18], our generalized Orlicz-Morrey space is the third kind and the ones in [25] and [31] are the first kind and the second kind, respectively. According to the examples in [9], one can say that the generalized Orlicz-Morrey spaces of the first kind and the second kind are different and that second kind and third kind are different. However, we do not know the relation between the first and the second kind.

Note that, Orlicz-Morrey spaces unify Orlicz and generalized Morrey spaces. We extend some results on generalized Morrey space in the papers [7, 12, 14, 15, 19, 20] to the case of Orlicz-Morrey space in [5, 8, 16–18].

Consider the half-space $\mathbb{R}_+^{n+1} = \mathbb{R}^n \times (0, \infty)$. For $x = (x', t) \in \mathbb{R}_+^{n+1}$, $x = (x'', x_n, t) \in \mathbb{D}_+^{n+1} = \mathbb{R}^{n-1} \times \mathbb{R}_+ \times \mathbb{R}_+$, $\mathbb{D}_-^{n+1} = \mathbb{R}^{n-1} \times \mathbb{R}_- \times \mathbb{R}_+$. In the following, besides the standard parabolic metric $\varrho(x) = \max(|x'|, |t|^{1/2})$ we use the equivalent one $\rho(x) = \left(\frac{|x'|^2 + \sqrt{|x'|^4 + 4t^2}}{2} \right)^{1/2}$ introduced by Fabes and Rivi re in [4]. The induced by it topology consists of ellipsoids (parabolic balls)

$$\mathcal{E}_r(x) = \left\{ y \in \mathbb{R}^{n+1} : \frac{|x' - y'|^2}{r^2} + \frac{|t - \tau|^2}{r^4} < 1 \right\}, \quad |\mathcal{E}_r| = Cr^{n+2}.$$

It is easy to see that $\mathcal{E}_1(x)$ and \mathbb{S}^n are the unit ball and the unit sphere, respectively, with respect to the both metrics and $\rho(x)$. On the other hand, the equivalence between the both parabolic metrics $\varrho(x)$ and $\rho(x)$ follows by the inclusion: for each \mathcal{E}_r there exist parabolic cylinders $\underline{\mathcal{C}}$ and $\overline{\mathcal{C}}$ with measure comparable with r^{n+2} such that $\underline{\mathcal{C}} \subset \mathcal{E}_r \subset \overline{\mathcal{C}}$. In what follows all estimate obtained over ellipsoids hold true also over parabolic cylinders and we shall use this property without explicit references.

Let $\tilde{x} = (x'', -x_n, t)$ be the "reflected point". The parabolic nonsingular integral operator \mathcal{R} is defined by (see [2])

$$\mathcal{R}f(x) = \int_{\mathbb{D}_+^{n+1}} \frac{|f(y)|}{\rho(\tilde{x} - y)^{n+2}} dy. \quad (1.1)$$

The operator \mathcal{R} and its commutator appear in [2] in connection with boundary estimates for solutions to parabolic equations.

In [28] the author was study the boundedness of the parabolic nonsingular integral operator \mathcal{R} on Orlicz spaces $L^\Phi(\mathbb{D}_+^{n+1})$. Therefore, the purpose of this paper is mainly to study the boundedness of the operator \mathcal{R} on parabolic generalized Orlicz-Morrey spaces of the third kind $M^{\Phi, \varphi}(\mathbb{D}_+^{n+1})$.

A function $\varphi : (0, \infty) \rightarrow (0, \infty)$ is said to be almost increasing (resp. almost decreasing) if there exists a constant $C > 0$ such that

$$\varphi(r) \leq C\varphi(s) \quad (\text{resp. } \varphi(r) \geq C\varphi(s)) \quad \text{for } r \leq s.$$

For a Young function Φ , we denote by \mathcal{G}_Φ the set of all decreasing functions $\varphi : (0, \infty) \rightarrow (0, \infty)$ such that $t \in (0, \infty) \mapsto \Phi^{-1}(t^{-n-2})\varphi(t)^{-1}$ is almost decreasing.

The following results are the fundamental theorems in this paper:

Theorem 1.1 *Let $\Phi \in \Delta_2$ and $\varphi_1, \varphi_2 \in \Omega_\Phi$.*

1. *The condition*

$$\int_r^\infty \left(\text{ess inf}_{t < s < \infty} \frac{\varphi_1(x, s)}{\Phi^{-1}(s^{-n-2})} \right) \Phi^{-1}(t^{-n-2}) \frac{dt}{t} \leq C \varphi_2(x, r), \quad (1.2)$$

where C does not depend on x and r , is sufficient for the boundedness of \mathcal{R} from $M^{\Phi, \varphi_1}(\mathbb{D}_+^{n+1})$ to $WM^{\Phi, \varphi_2}(\mathbb{D}_+^{n+1})$. If, in addition, $\Phi \in \nabla_2$, then the condition (1.2) is sufficient for the boundedness of \mathcal{R} from $M^{\Phi, \varphi_1}(\mathbb{D}_+^{n+1})$ to $M^{\Phi, \varphi_2}(\mathbb{D}_+^{n+1})$.

2. If $\varphi_1 \in \mathcal{G}_\Phi$, then the condition

$$\varphi_1(x, r) \leq C\varphi_2(x, r), \quad (1.3)$$

where C does not depend on x and r , is necessary for the boundedness of \mathcal{R} from $M^{\Phi, \varphi_1}(\mathbb{D}_+^{n+1})$ to $WM^{\Phi, \varphi_2}(\mathbb{D}_+^{n+1})$ and from $M^{\Phi, \varphi_1}(\mathbb{D}_+^{n+1})$ to $M^{\Phi, \varphi_2}(\mathbb{D}_+^{n+1})$.

3. If $\varphi_1 \in \mathcal{G}_\Phi$ satisfies the regularity type condition

$$\int_t^\infty \varphi_1(r) \frac{dr}{r} \leq C\varphi_1(t), \quad (1.4)$$

for all $t > 0$, where $C > 0$ does not depend on t , then the condition (1.3) is necessary and sufficient for the boundedness of \mathcal{R} from $M^{\Phi, \varphi_1}(\mathbb{D}_+^{n+1})$ to $WM^{\Phi, \varphi_2}(\mathbb{D}_+^{n+1})$. If, in addition, $\Phi \in \nabla_2$, then the condition (1.3) is necessary and sufficient for the boundedness of \mathcal{R} from $M^{\Phi, \varphi_1}(\mathbb{D}_+^{n+1})$ to $M^{\Phi, \varphi_2}(\mathbb{D}_+^{n+1})$.

If we take $\Phi(t) = t^p$, $p \in [1, \infty)$ at Theorem 1.1 we get the following new result for generalized Morrey spaces.

Corollary 1.1 Let $p \in [1, \infty)$ and $\varphi_1, \varphi_2 \in \Omega_p \equiv \Omega_{t^p}$.

1. The condition

$$\int_r^\infty \frac{\operatorname{ess\,inf}_{t < s < \infty} \varphi_1(s) s^{\frac{n+2}{p}}}{t^{\frac{n+2}{p}+1}} dt \leq C\varphi_2(r), \quad (1.5)$$

for all $r > 0$, where $C > 0$ does not depend on r , is sufficient for the boundedness of \mathcal{R} from $M^{p, \varphi_1}(\mathbb{D}_+^{n+1})$ to $WM^{p, \varphi_2}(\mathbb{D}_+^{n+1})$. If $1 < p < \infty$, then the condition (1.5) is sufficient for the boundedness of \mathcal{R} from $M^{p, \varphi_1}(\mathbb{D}_+^{n+1})$ to $M^{p, \varphi_2}(\mathbb{D}_+^{n+1})$.

2. If $\varphi_1 \in \mathcal{G}_p$, then the condition (1.3) is necessary for the boundedness of \mathcal{R} from $M^{p, \varphi_1}(\mathbb{D}_+^{n+1})$ to $WM^{p, \varphi_2}(\mathbb{D}_+^{n+1})$ and from $M^{p, \varphi_1}(\mathbb{D}_+^{n+1})$ to $M^{p, \varphi_2}(\mathbb{D}_+^{n+1})$.

3. If $\varphi_1 \in \mathcal{G}_p$ satisfies the regularity condition (1.4), then the condition (1.3) is necessary and sufficient for the boundedness of \mathcal{R} from $M^{p, \varphi_1}(\mathbb{D}_+^{n+1})$ to $WM^{p, \varphi_2}(\mathbb{D}_+^{n+1})$. If, in addition, $1 < p < \infty$, then the condition (1.3) is necessary and sufficient for the boundedness of \mathcal{R} from $M^{p, \varphi_1}(\mathbb{D}_+^{n+1})$ to $M^{p, \varphi_2}(\mathbb{D}_+^{n+1})$.

Remark 1.1 Note that, in [8] was study the boundedness of the nonsingular integral operator on generalized Orlicz-Morrey spaces of the third kind $M^{\Phi, \varphi}(\mathbb{R}_+^n)$.

By $A \lesssim B$ we mean that $A \leq CB$ with some positive constant C independent of appropriate quantities. If $A \lesssim B$ and $B \lesssim A$, we write $A \approx B$ and say that A and B are equivalent.

2 Definitions and Preliminary Results

2.1 On Young Functions and Orlicz Spaces

We recall the definition of Young functions.

Definition 2.1 A function $\Phi : [0, \infty) \rightarrow [0, \infty]$ is called a Young function if Φ is convex, left-continuous, $\lim_{r \rightarrow +0} \Phi(r) = \Phi(0) = 0$ and $\lim_{r \rightarrow \infty} \Phi(r) = \infty$.

From the convexity and $\Phi(0) = 0$ it follows that any Young function is increasing. If there exists $s \in (0, \infty)$ such that $\Phi(s) = \infty$, then $\Phi(r) = \infty$ for $r \geq s$. The set of Young functions such that

$$0 < \Phi(r) < \infty \quad \text{for} \quad 0 < r < \infty$$

will be denoted by \mathcal{Y} . If $\Phi \in \mathcal{Y}$, then Φ is absolutely continuous on every closed interval in $[0, \infty)$ and bijective from $[0, \infty)$ to itself.

For a Young function Φ and $0 \leq s \leq \infty$, let

$$\Phi^{-1}(s) = \inf\{r \geq 0 : \Phi(r) > s\}.$$

If $\Phi \in \mathcal{Y}$, then Φ^{-1} is the usual inverse function of Φ . We note that

$$\Phi(\Phi^{-1}(r)) \leq r \leq \Phi^{-1}(\Phi(r)) \quad \text{for } 0 \leq r < \infty.$$

It is well known that

$$r \leq \Phi^{-1}(r)\tilde{\Phi}^{-1}(r) \leq 2r \quad \text{for } r \geq 0, \quad (2.1)$$

where $\tilde{\Phi}(r)$ is defined by

$$\tilde{\Phi}(r) = \begin{cases} \sup\{rs - \Phi(s) : s \in [0, \infty)\}, & r \in [0, \infty) \\ \infty, & r = \infty. \end{cases}$$

A Young function Φ is said to satisfy the Δ_2 -condition, denoted also as $\Phi \in \Delta_2$, if

$$\Phi(2r) \leq k\Phi(r) \quad \text{for } r > 0$$

for some $k > 1$. If $\Phi \in \Delta_2$, then $\Phi \in \mathcal{Y}$. A Young function Φ is said to satisfy the ∇_2 -condition, denoted also by $\Phi \in \nabla_2$, if

$$\Phi(r) \leq \frac{1}{2k}\Phi(kr), \quad r \geq 0,$$

for some $k > 1$.

Definition 2.2 (*Orlicz Space*). For a Young function Φ , the set

$$L^\Phi(\mathbb{D}_+^{n+1}) = \left\{ f \in L_{\text{loc}}^1(\mathbb{D}_+^{n+1}) : \int_{\mathbb{D}_+^{n+1}} \Phi(k|f(x)|)dx < \infty \text{ for some } k > 0 \right\}$$

is called *Orlicz space*. If $\Phi(r) = r^p$, $1 \leq p < \infty$, then $L^\Phi(\mathbb{D}_+^{n+1}) = L^p(\mathbb{D}_+^{n+1})$. If $\Phi(r) = 0$, ($0 \leq r \leq 1$) and $\Phi(r) = \infty$, ($r > 1$), then $L^\Phi(\mathbb{D}_+^{n+1}) = L^\infty(\mathbb{D}_+^{n+1})$. The space $L_{\text{loc}}^\Phi(\mathbb{D}_+^{n+1})$ is defined as the set of all functions f such that $f\chi_B \in L^\Phi(\mathbb{D}_+^{n+1})$ for all balls $B \subset \mathbb{D}_+^{n+1}$.

$L^\Phi(\mathbb{D}_+^{n+1})$ is a Banach space with respect to the norm

$$\|f\|_{L^\Phi(\mathbb{D}_+^{n+1})} = \inf \left\{ \lambda > 0 : \int_{\mathbb{D}_+^{n+1}} \Phi\left(\frac{|f(x)|}{\lambda}\right)dx \leq 1 \right\}.$$

We note that

$$\int_{\mathbb{D}_+^{n+1}} \Phi\left(\frac{|f(x)|}{\|f\|_{L^\Phi(\mathbb{D}_+^{n+1})}}\right)dx \leq 1. \quad (2.2)$$

The weak Orlicz space

$$WL^\Phi(\mathbb{D}_+^{n+1}) = \{f \in L_{\text{loc}}^1(\mathbb{D}_+^{n+1}) : \|f\|_{WL^\Phi(\mathbb{D}_+^{n+1})} < +\infty\}$$

is defined by the norm

$$\|f\|_{WL^\Phi(\mathbb{D}_+^{n+1})} = \inf \left\{ \lambda > 0 : \sup_{t>0} \Phi(t) m\left(\frac{f}{\lambda}, t\right) \leq 1 \right\}.$$

The following lemmas are valid.

Lemma 2.1 [1, 21] *Let Φ be a Young function and \mathcal{E} a set in \mathbb{D}_+^{n+1} with finite Lebesgue measure. Then*

$$\|\chi_{\mathcal{E}}\|_{WL^\Phi(\mathbb{D}_+^{n+1})} = \|\chi_{\mathcal{E}}\|_{L^\Phi(\mathbb{D}_+^{n+1})} = \frac{1}{\Phi^{-1}(|\mathcal{E}|^{-1})}.$$

Lemma 2.2 *For a Young function Φ and for all parabolic balls \mathcal{E} in \mathbb{D}_+^{n+1} , the following inequality is valid*

$$\|f\|_{L^1(\mathcal{E})} \leq 2|\mathcal{E}|\Phi^{-1}(|\mathcal{E}|^{-1})\|f\|_{L^\Phi(\mathcal{E})}.$$

The following theorem were proved in [28].

Theorem 2.1 [28] *Let Φ be a Young function and \mathcal{R} be a parabolic nonsingular integral operator, defined by (1.1). If $\Phi \in \Delta_2 \cap \nabla_2$, then the operator \mathcal{R} is bounded on $L^\Phi(\mathbb{D}_+^{n+1})$ and if $\Phi \in \Delta_2$, then the operator \mathcal{R} is bounded from $L^\Phi(\mathbb{D}_+^{n+1})$ to $WL^\Phi(\mathbb{D}_+^{n+1})$.*

In the case $\Phi(t) = t^p$, we get the following result proved in [2].

Corollary 2.1 [2] *Let $1 \leq p < \infty$ and $f \in L^p(\mathbb{D}_+^{n+1})$. Then there exists a constant C_p independent of f , such that*

$$\|\mathcal{R}f\|_{L^p(\mathbb{D}_+^{n+1})} \leq C_p \|f\|_{L^p(\mathbb{D}_+^{n+1})}, \quad 1 < p < \infty$$

and

$$\|\mathcal{R}f\|_{WL^1(\mathbb{D}_+^{n+1})} \leq C_1 \|f\|_{L^1(\mathbb{D}_+^{n+1})}.$$

2.2 Parabolic generalized Orlicz-Morrey Space

Various versions of generalized Orlicz-Morrey spaces were introduced in [25], [31] and [5]. We used the definition of [5] which runs as follows.

We now define parabolic generalized Orlicz-Morrey spaces of the third kind. The *parabolic generalized Orlicz-Morrey space $M^{\Phi, \phi}(\mathbb{D}_+^{n+1})$ of the third kind* is defined as the set of all measurable functions f for which the norm

$$\|f\|_{M^{\Phi, \phi}(\mathbb{D}_+^{n+1})} \equiv \sup_{x \in \mathbb{D}_+^{n+1}, r > 0} \frac{1}{\phi(x, r)} \Phi^{-1} \left(\frac{1}{|\mathcal{E}^+(x, r)|} \right) \|f\|_{L^\Phi(\mathcal{E}^+(x, r))}$$

is finite, where $\mathcal{E}^+(x, r) = \mathcal{E}(x, r) \cap \mathbb{D}_+^{n+1}$. Also by $WM^{\Phi, \varphi}(\mathbb{D}_+^{n+1})$ we denote the *weak parabolic generalized Orlicz-Morrey space of the third kind* of all functions $f \in WL_{\text{loc}}^\Phi(\mathbb{D}_+^{n+1})$ for which

$$\|f\|_{WM^{\Phi, \varphi}(\mathbb{D}_+^{n+1})} = \sup_{x \in \mathbb{D}_+^{n+1}, r > 0} \varphi(x, r)^{-1} \Phi^{-1}(|\mathcal{E}^+(x, r)|^{-1}) \|f\|_{WL^\Phi(\mathcal{E}^+(x, r))} < \infty,$$

where $WL^\Phi(\mathcal{E}^+(x, r))$ denotes the weak L^Φ -space of measurable functions f for which

$$\|f\|_{WL^\Phi(\mathcal{E}^+(x, r))} \equiv \|f\chi_{\mathcal{E}^+(x, r)}\|_{WL^\Phi(\mathbb{D}_+^{n+1})}.$$

Note that $M^{\Phi, \phi}(\mathbb{D}_+^{n+1})$ covers many classical function spaces.

Example 2.1 Let $1 \leq q \leq p < \infty$ and $\Phi \in \Delta_2 \cap \nabla_2$. From the following special cases, we see that our results will cover the Lebesgue space $L^p(\mathbb{D}_+^{n+1})$, the *classical Morrey space* $M_q^p(\mathbb{D}_+^{n+1})$, the generalized Morrey space $M^{\phi,p}(\mathbb{D}_+^{n+1})$ and the Orlicz space $L^\Phi(\mathbb{D}_+^{n+1})$ with norm coincidence:

- 1 If $\Phi(t) = t^p$ and $\phi(t) = t^{-\frac{n+2}{p}}$, then $M^{\Phi,\phi}(\mathbb{D}_+^{n+1}) = L^p(\mathbb{D}_+^{n+1})$ with norm equivalence.
- 2 If $\Phi(t) = t^q$ and $\phi(t) = t^{-\frac{n+2}{p}}$, then $M^{\Phi,\phi}(\mathbb{D}_+^{n+1})$, which is denoted by $M_q^p(\mathbb{D}_+^{n+1})$, is the parabolic Morrey space.
- 3 If $\Phi(t) = t^p$, then $M^{\Phi,\phi}(\mathbb{D}_+^{n+1}) = M^{p,\phi}(\mathbb{D}_+^{n+1})$ is the parabolic generalized Morrey space which were discussed in [10], see also [12, 22, 24].
- 4 If $\phi(t) = \Phi^{-1}(t^{-n-2})$, then $M^{\Phi,\phi}(\mathbb{D}_+^{n+1}) = L^\Phi(\mathbb{D}_+^{n+1})$.

Other definitions of generalized Orlicz-Morrey spaces can be found in [9, 25–27]. Therefore, our definition of generalized Orlicz-Morrey spaces here is named “third kind”.

In the case $\varphi(x, r) = \frac{\Phi^{-1}(|\mathcal{E}(x, r)|^{-1})}{\Phi^{-1}(|\mathcal{E}(x, r)|^{-\lambda/n})}$, we get the parabolic Orlicz-Morrey space $M^{\Phi,\lambda}(\mathbb{R}^n)$ from parabolic generalized Orlicz-Morrey space $M^{\Phi,\varphi}(\mathbb{R}^n)$. We refer to [6, Lemmas 2.8 and 2.9] for more information about Orlicz-Morrey spaces.

Lemma 2.3 [6, Lemma 2.12] *Let Φ be a Young function and φ be a positive measurable function on $\mathbb{R}^n \times (0, \infty)$.*

(i) *If*

$$\sup_{t < r < \infty} \frac{\Phi^{-1}(|\mathcal{E}(x, r)|^{-1})}{\varphi(x, r)} = \infty \quad \text{for some } t > 0 \text{ and for all } x \in \mathbb{R}^n, \quad (2.3)$$

then $M^{\Phi,\varphi}(\mathbb{R}^n) = \emptyset$.

(ii) *If*

$$\sup_{0 < r < \tau} \varphi(x, r)^{-1} = \infty \quad \text{for some } \tau > 0 \text{ and for all } x \in \mathbb{R}^n, \quad (2.4)$$

then $M^{\Phi,\varphi}(\mathbb{R}^n) = \emptyset$.

Remark 2.1 Let Φ be a Young function. We denote by Ω_Φ the sets of all positive measurable functions φ on $\mathbb{R}^n \times (0, \infty)$ such that for all $t > 0$,

$$\sup_{x \in \mathbb{R}^n} \left\| \frac{\Phi^{-1}(|\mathcal{E}(x, r)|^{-1})}{\varphi(x, r)} \right\|_{L^\infty(t, \infty)} < \infty,$$

and

$$\sup_{x \in \mathbb{R}^n} \left\| \varphi(x, r)^{-1} \right\|_{L^\infty(0, t)} < \infty,$$

respectively. In what follows, keeping in mind Lemma 2.3, we always assume that $\varphi \in \Omega_\Phi$.

The following lemma plays a key role in our main results.

Lemma 2.4 *Let $\mathcal{E}_0^+ := \mathcal{E}^+(x_0, r_0)$ a parabolic ball in \mathbb{D}_+^{n+1} . If $\varphi \in \mathcal{G}_\Phi$, then there exist $C > 0$ such that*

$$\frac{1}{\varphi(r_0)} \leq \|\chi_{\mathcal{E}_0^+}\|_{M^{\Phi,\varphi}(\mathbb{D}_+^{n+1})} \leq \frac{C}{\varphi(r_0)}.$$

Proof. Let $\mathcal{E}^+ = \mathcal{E}^+(x, r)$ denote an arbitrary parabolic ball in \mathbb{D}_+^{n+1} . By the definition and Lemma 2.1, it is easy to see that

$$\begin{aligned} \|\chi_{\mathcal{E}_0^+}\|_{M^{\Phi, \varphi}} &= \sup_{x \in \mathbb{D}_+^{n+1}, r > 0} \varphi(r)^{-1} \Phi^{-1}(|\mathcal{E}^+|^{-1}) \frac{1}{\Phi^{-1}(|\mathcal{E}^+ \cap \mathcal{E}_0^+|^{-1})} \\ &\geq \varphi(r_0)^{-1} \Phi^{-1}(|\mathcal{E}_0^+|^{-1}) \frac{1}{\Phi^{-1}(|\mathcal{E}_0^+ \cap \mathcal{E}_0^+|^{-1})} = \frac{1}{\varphi(r_0)}. \end{aligned}$$

Now if $r \leq r_0$, then $\varphi(r_0) \leq C\varphi(r)$ and

$$\varphi(r)^{-1} \Phi^{-1}(|\mathcal{E}^+|^{-1}) \|\chi_{\mathcal{E}_0^+}\|_{L^{\Phi}(\mathcal{E}^+)} \leq \frac{1}{\varphi(r)} \leq \frac{C}{\varphi(r_0)}.$$

On the other hand if $r \geq r_0$, then $\frac{\varphi(r_0)}{\Phi^{-1}(|\mathcal{E}_0^+|^{-1})} \leq C \frac{\varphi(r)}{\Phi^{-1}(|\mathcal{E}^+|^{-1})}$ and

$$\varphi(r)^{-1} \Phi^{-1}(|\mathcal{E}^+|^{-1}) \|\chi_{\mathcal{E}_0^+}\|_{L^{\Phi}(\mathcal{E}^+)} \leq \frac{C}{\varphi(r_0)}.$$

This completes the proof.

3 Parabolic nonsingular integral operators in the space $M^{\Phi, \varphi}(\mathbb{D}_+^{n+1})$

We will use the following statement on the boundedness of the weighted Hardy operator

$$H_w^* g(t) := \int_t^\infty g(s) w(s) ds, \quad 0 < t < \infty,$$

where w is a weight.

The following theorem were proved in [13, 15] and in the case $w = 1$ in [3].

Theorem 3.1 *Let v_1, v_2 and w be weights on $(0, \infty)$ and $v_1(t)$ be bounded outside a neighborhood of the origin. The inequality*

$$\sup_{t > 0} v_2(t) H_w^* g(t) \leq C \sup_{t > 0} v_1(t) g(t) \quad (3.1)$$

holds for some $C > 0$ for all non-negative and non-decreasing g on $(0, \infty)$ if and only if

$$B := \sup_{t > 0} v_2(t) \int_t^\infty \frac{w(s) ds}{\sup_{s < \tau < \infty} v_1(\tau)} < \infty. \quad (3.2)$$

Moreover, the value $C = B$ is the best constant for (3.1).

Remark 3.1 In (3.1) and (3.2) it is assumed that $\frac{1}{\infty} = 0$ and $0 \cdot \infty = 0$.

For any $x \in \mathbb{D}_+^{n+1}$ define $\tilde{x} = (x', -x_n)$ and recall that $x^0 = (x', 0)$. Also define $\mathcal{E}_r^+ \equiv \mathcal{E}^+(x^0, r) = \mathcal{E}(x^0, r) \cap \mathbb{D}_+^{n+1}$, $2\mathcal{E}_r^+ = \mathcal{E}^+(x^0, 2r)$.

Lemma 3.1 *Let Φ any Young function, $f \in L_{\text{loc}}^{\Phi}(\mathbb{D}_+^{n+1})$, be such that*

$$\int_1^{\infty} \|f\|_{L^{\Phi}(\mathcal{E}^+(x^0, t))} \Phi^{-1}(t^{-n-2}) \frac{dt}{t} < \infty \quad (3.3)$$

i) *If $\Phi \in \Delta_2 \cap \nabla_2$, then*

$$\|\mathcal{R}f\|_{L^{\Phi}(\mathcal{E}^+(x^0, r))} \leq \frac{C}{\Phi^{-1}(r^{-n-2})} \int_{2r}^{\infty} \|f\|_{L^{\Phi}(\mathcal{E}^+(x^0, t))} \Phi^{-1}(t^{-n-2}) \frac{dt}{t}. \quad (3.4)$$

ii) *If $\Phi \in \Delta_2$, then*

$$\|\mathcal{R}f\|_{WL^{\Phi}(\mathcal{E}^+(x^0, r))} \leq \frac{C}{\Phi^{-1}(r^{-n-2})} \int_{2r}^{\infty} \|f\|_{L^{\Phi}(\mathcal{E}^+(x^0, t))} \Phi^{-1}(t^{-n-2}) \frac{dt}{t}, \quad (3.5)$$

where the constants are independent of x^0 , r and f .

Proof. i) Denote by $\mathcal{E}_r^+ = \mathcal{E}^+(x^0, r)$, $\mathcal{E}_t^+ = \mathcal{E}^+(x^0, t)$ and for any $f \in L_{\text{loc}}^{\Phi}(\mathbb{D}_+^{n+1})$ write $f = f_1 + f_2$ with $f_1 = f\chi_{2\mathcal{E}_r^+}$ and $f_2 = f\chi_{(2\mathcal{E}_r^+)^c}$. Because of the Φ -boundedness of the operator \mathcal{R} (see Theorem 2.1) and $f_1 \in L^{\Phi}(\mathbb{D}_+^{n+1})$ we have

$$\|\mathcal{R}f_1\|_{L^{\Phi}(\mathcal{E}_r^+)} \leq \|\mathcal{R}f_1\|_{L^{\Phi}(\mathbb{D}_+^{n+1})} \leq C\|f_1\|_{L^{\Phi}(\mathbb{D}_+^{n+1})} = C\|f\|_{L^{\Phi}(2\mathcal{E}_r^+)}.$$

It is easy to see that for arbitrary points $x \in \mathcal{E}_r^+$ and $y \in (2\mathcal{E}_r^+)^c$ it holds

$$\frac{1}{2}\rho(x^0 - y) \leq \rho(\tilde{x} - y) \leq \frac{3}{2}\rho(x^0 - y). \quad (3.6)$$

Applying the Fubini theorem to $\mathcal{R}f_2$ we get

$$\begin{aligned} |\mathcal{R}f_2(x)| &\leq C \int_{\mathbb{D}_+^{n+1}} \frac{|f_2(y)|}{\rho(\tilde{x} - y)^{n+2}} dy \\ &\leq C \int_{(2\mathcal{E}_r^+)^c} \frac{|f(y)|}{\rho(x^0 - y)^{n+2}} dy \leq C \int_{(2\mathcal{E}_r^+)^c} |f(y)| dy \int_{\rho(x^0 - y)}^{\infty} \frac{dt}{t^{n+3}} \\ &\leq C \int_{2r}^{\infty} \left(\int_{2r \leq \rho(x^0 - y) < t} |f(y)| dy \right) \frac{dt}{t^{n+3}} \leq C \int_{2r}^{\infty} \left(\int_{\mathcal{E}_t^+} |f(y)| dy \right) \frac{dt}{t^{n+3}}. \end{aligned}$$

Applying the Hölder's inequality (see, Lemma 2.2), we get

$$\begin{aligned} \int_{(2\mathcal{E}_r^+)^c} \frac{|f(y)|}{\rho(x^0 - y)^{n+2}} dy &\lesssim \int_{2r}^{\infty} \|f\|_{L^{\Phi}(\mathcal{E}_t^+)} \|1\|_{L^{\tilde{\Phi}}(\mathcal{E}_t^+)} \frac{dt}{t^{n+3}} \\ &= \int_{2r}^{\infty} \|f\|_{L^{\Phi}(\mathcal{E}_t^+)} \frac{1}{\tilde{\Phi}^{-1}(|\mathcal{E}_t^+|^{-1})} \frac{dt}{t^{n+3}} \approx \int_{2r}^{\infty} \|f\|_{L^{\Phi}(\mathcal{E}_t^+)} \Phi^{-1}(t^{-n-2}) \frac{dt}{t}. \end{aligned} \quad (3.7)$$

Direct calculations give

$$\|\mathcal{R}f_2\|_{L^{\Phi}(\mathcal{E}_r^+)} \leq \frac{C}{\Phi^{-1}(r^{-n-2})} \int_{2r}^{\infty} \|f\|_{L^{\Phi}(\mathcal{E}_t^+)} \Phi^{-1}(t^{-n-2}) \frac{dt}{t} \quad (3.8)$$

and the last estimate holds for all $f \in L^{\Phi}(\mathbb{D}_+^{n+1})$ satisfying (3.3). Thus

$$\|\mathcal{R}f\|_{L^{\Phi}(\mathcal{E}_r^+)} \lesssim \|f\|_{L^{\Phi}(2\mathcal{E}_r^+)} + \frac{1}{\Phi^{-1}(r^{-n-2})} \int_{2r}^{\infty} \|f\|_{L^{\Phi}(\mathcal{E}_t^+)} \Phi^{-1}(t^{-n-2}) \frac{dt}{t}. \quad (3.9)$$

On the other hand,

$$\begin{aligned} \|f\|_{L^\Phi(2\mathcal{E}_r)} &= \frac{C}{\Phi^{-1}(r^{-n-2})} \|f\|_{L^\Phi(2\mathcal{E}_r)} \int_{2r}^{\infty} \Phi^{-1}(t^{-n-2}) \frac{dt}{t} \\ &\leq \frac{C}{\Phi^{-1}(r^{-n-2})} \int_{2r}^{\infty} \|f\|_{L^\Phi(\mathcal{E}_t^+)} \Phi^{-1}(t^{-n-2}) \frac{dt}{t} \end{aligned} \quad (3.10)$$

which unified with (3.9) gives (3.4).

ii) Let now $f \in L^\Phi(\mathbb{D}_+^{n+1})$, the weak Φ -boundedness of \mathcal{R} (see Theorem 2.1) implies

$$\|\mathcal{R}f_1\|_{WL^\Phi(\mathcal{E}_r^+)} \leq \|\mathcal{R}f_1\|_{WL^\Phi(\mathbb{D}_+^{n+1})} \lesssim \|f_1\|_{L^\Phi(\mathbb{D}_+^{n+1})} = \|f\|_{L^\Phi(2\mathcal{E}_r^+)}.$$

The estimate (3.5) follows by (3.8).

For proving our main results, we need the following estimate.

Lemma 3.2 *If $\mathcal{E}_0^+ := \mathcal{E}^+(x_0, r_0)$, then $C \leq \mathcal{R}\chi_{\mathcal{E}_0^+}(x)$ for every $x \in \mathcal{E}_0^+$.*

Proof. If $x, y \in \mathcal{E}_0^+$, then $\rho(\tilde{x} - y) \leq \rho(\tilde{x} - x_0) + \rho(y - x_0) < 2r_0$. We get $Cr_0^{-n-2} \leq \rho(\tilde{x} - y)^{-n-2}$. Therefore

$$\mathcal{R}\chi_{\mathcal{E}_0^+}(x) = \int_{\mathbb{R}^n} \chi_{\mathcal{E}_0^+}(y) \rho(\tilde{x} - y)^{-n-2} dy = \int_{\mathcal{E}_0^+} \rho(\tilde{x} - y)^{-n-2} dy \geq Cr_0^{-n-2} |\mathcal{E}_0^+| = C.$$

Theorem 3.2 *Let Φ any Young function, $\varphi_1, \varphi_2 : \mathbb{D}_+^{n+1} \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be measurable functions satisfying (1.2).*

i) *If $\Phi \in \Delta_2 \cap \nabla_2$, then it is bounded from $M^{\Phi, \varphi_1}(\mathbb{D}_+^{n+1})$ in $M^{\Phi, \varphi_2}(\mathbb{D}_+^{n+1})$ and*

$$\|\mathcal{R}f\|_{M^{\Phi, \varphi_2}(\mathbb{D}_+^{n+1})} \leq C \|f\|_{M^{\Phi, \varphi_1}(\mathbb{D}_+^{n+1})}. \quad (3.11)$$

ii) *If $\Phi \in \Delta_2$, then it is bounded from $M^{\Phi, \varphi_1}(\mathbb{D}_+^{n+1})$ to $WM^{\Phi, \varphi_2}(\mathbb{D}_+^{n+1})$ and*

$$\|\mathcal{R}f\|_{M^{\Phi, \varphi_2}(\mathbb{D}_+^{n+1})} \leq C \|f\|_{WM^{\Phi, \varphi_1}(\mathbb{D}_+^{n+1})}$$

with constants independent of f .

Proof. Let \mathcal{R} bounded in $L^\Phi(\mathbb{D}_+^{n+1})$ than by Lemma 3.1 we have

$$\|\mathcal{R}f\|_{M^{\Phi, \varphi_2}(\mathbb{D}_+^{n+1})} \leq C \sup_{x^0, r > 0} \varphi_2(x^0, r)^{-1} \int_r^{\infty} \|f\|_{L^\Phi(\mathcal{E}^+(x^0, t))} \Phi^{-1}(t^{-n-2}) \frac{dt}{t}.$$

Applying the Theorem 3.1 to the above integral with

$$w(r) = \Phi^{-1}(r^{-n-2}), \quad v_2(x^0, r) = \varphi_2(x^0, r)^{-1}, \quad v_1(x^0, r) = \varphi_1(x^0, r)^{-1} \Phi^{-1}(r^{-n-2}),$$

$$g(x^0, r) = \|f\|_{L^\Phi(\mathcal{E}^+(x^0, r))}, \quad H_w^* g(x^0, r) = \int_r^{\infty} \|f\|_{L^\Phi(\mathcal{E}^+(x^0, t))} w(t) dt$$

where the condition (3.2) is equivalent to (1.2) we get

$$\|\mathcal{R}f\|_{M^{\Phi, \varphi_2}(\mathbb{D}_+^{n+1})} \leq C \sup_{x \in \mathbb{D}_+^{n+1}, r > 0} \varphi_1(x^0, r)^{-1} \Phi^{-1}(r^{-n-2}) \|f\|_{L^\Phi(\mathcal{E}^+(x^0, r))} = C \|f\|_{M^{\Phi, \varphi_1}(\mathbb{D}_+^{n+1})}.$$

The case $p = 1$ is treated in the same manner making use of (3.5) and (3.2)

$$\begin{aligned} \|\mathcal{R}f\|_{WM^{\Phi, \varphi_2}(\mathbb{D}_+^{n+1})} &\leq C \sup_{x^0 \in \mathbb{D}_+^{n+1}, r > 0} \varphi_2(x^0, r)^{-1} \int_r^{\infty} \|f\|_{L^\Phi(\mathcal{E}^+(x^0, t))} \Phi^{-1}(t^{-n-2}) \frac{dt}{t} \\ &= C \sup_{x^0, r > 0} \varphi_1(x^0, r)^{-1} \Phi^{-1}(r^{-n-2}) \|f\|_{L^\Phi(\mathcal{E}^+(x^0, r))} = C \|f\|_{M^{\Phi, \varphi_1}(\mathbb{D}_+^{n+1})}. \end{aligned}$$

Proof of Theorem 1.1. The first part of the theorem follows from Lemma 3.1 and Theorem 3.2. We shall now prove the second part. Let $\mathcal{E}_0^+ = \mathcal{E}^+(x_0, r_0)$ and $x \in \mathcal{E}_0^+$. It is easy to see that $\mathcal{R}\chi_{\mathcal{E}_0^+}(x) = 1$ for every $x \in \mathcal{E}_0^+$. Therefore, by Lemmas 2.1 and 3.2

$$\begin{aligned} 1 &= \Phi^{-1}(w(\mathcal{E}_0^+)^{-1}) \|\mathcal{R}\chi_{\mathcal{E}_0^+}\|_{L^\Phi(\mathcal{E}_0^+)} \leq \varphi_2(\mathcal{E}_0^+) \|\mathcal{R}\chi_{\mathcal{E}_0^+}\|_{M^{\Phi, \varphi_2}} \\ &\leq C\varphi_2(\mathcal{E}_0^+) \|\chi_{\mathcal{E}_0^+}\|_{M^{\Phi, \varphi_1}} \leq C \frac{\varphi_2(\mathcal{E}_0^+)}{\varphi_1(\mathcal{E}_0^+)}. \end{aligned}$$

Since this is true for every \mathcal{E}_0^+ , we are done.

The third statement of the theorem follows from the other statements of the theorem. \square

Acknowledgements. We thank the referee(s) for careful reading the paper and useful comments. The research of M. Omarova was partially supported by the Grant of 1st Azerbaijan-Russia Joint Grant Competition (Agreement Number No. EIF-BGM-4-RFTF-1/2017-21/01/1).

References

1. Bennett, C., Sharpley, R.: *Interpolation of operators*, Academic Press, Boston, (1988).
2. Bramanti, M., Cerutti, M.C.: $W_p^{1,2}$ solvability for the Cauchy-Dirichlet problem for parabolic equations with VMO coefficients, Comm. in Partial Diff. Eq. **18**, 1735-1763 (1993).
3. Burenkov, V., Gogatishvili, A., Guliyev, V.S., Mustafayev, R.: *Boundedness of the fractional maximal operator in local Morrey-type spaces*, Complex Var. Elliptic Equ. **55** (8-10), 739-758 (2010).
4. Fabes, E.B., Riviere, N.: Singular integrals with mixed homogeneity, Studia Math. **27**, 19-38 (1966).
5. Deringoz, F., Guliyev, V.S., Samko, S.: *Boundedness of maximal and singular operators on generalized Orlicz-Morrey spaces*. Operator Theory, Operator Algebras and Applications, Series: Operator Theory: Advances and Applications **242**, 139-158 (2014).
6. Deringoz, F., Guliyev, V.S., Hasanov, S.G.: *A characterization for Adams type boundedness of the fractional maximal operator on generalized Orlicz-Morrey spaces*, arXiv:1610.09823
7. Eroglu, A.: *Boundedness of fractional oscillatory integral operators and their commutators on generalized Morrey spaces*, Bound. Value Probl. 2013:70, 12 pp. (2013).
8. Eroglu, A., Guliyev, V.S., Omarova, M.N.: *Characterizations for the nonsingular integral operator and its commutators on generalized Orlicz-Morrey spaces*, Azerb. J. Math. **7** (2), 66-88 (2017).
9. Gala, S., Sawano, Y., Tanaka, H.: *A remark on two generalized Orlicz-Morrey spaces*, J. Approx. Theory **98**, 1-9 (2015).
10. Guliyev, V.S.: *Integral operators on function spaces on the homogeneous groups and on domains in \mathbb{R}^n* . (Russian) Doctor of Sciences, Mat. Inst. Steklova, Moscow, 329 pp. (1994).
11. Guliyev, V.S.: *Function spaces, integral operators and two weighted inequalities on homogeneous groups. Some applications*, Baku, 1-332 (1999). (Russian)
12. Guliyev, V.S.: *Boundedness of the maximal, potential and singular operators in the generalized Morrey spaces*, J. Inequal. Appl. Art. ID 503948, 20 pp (2009).
13. Guliyev, V.S.: *Generalized local Morrey spaces and fractional integral operators with rough kernel*, J. Math. Sci. (N. Y.) **193** (2), 211-227 (2013).

14. Guliyev, V.S., Aliyev, S.S., Karaman, T., Shukurov, P.S.: *Boundedness of sublinear operators and commutators on generalized Morrey space*, Int. Eq. Op. Theory. **71** (3), 327-355 (2011).
15. Guliyev, V.S.: *Local generalized Morrey spaces and singular integrals with rough kernel*, Azerb. J. Math. **3** (2), 79-94 (2013).
16. Guliyev, V.S., Deringoz, F., Hasanov, J.J.: *Φ -admissible singular operators and their commutators on vanishing generalized Orlicz-Morrey spaces*, J. Inequal. Appl. 2014:143, 18 pp. (2014).
17. Guliyev, V.S., Deringoz, F.: *On the Riesz potential and its commutators on generalized Orlicz-Morrey spaces*, J. Funct. Spaces. Article ID 617414, 11 pages (2014).
18. Guliyev, V.S., Hasanov, S.G., Sawano, Y., Noi, T.: *Non-smooth atomic decompositions for generalized Orlicz-Morrey spaces of the third kind*, Acta Appl. Math. **145** (1), 133-174 (2016).
19. Guliyev, V.S., Omarova, M.N.: *Multilinear singular and fractional integral operators on generalized weighted Morrey spaces*, Azerb. J. Math. **5** (1), 104-132 (2015).
20. Hakim, D.I., Sawano, Y., Shimomura, T.: *Boundedness of generalized fractional integral operators from the Morrey space $L_{1,\phi}(X; \mu)$ to the Campanato space $\mathcal{L}_{1,\phi}(X; \mu)$ over non-doubling measure spaces*, Azerb. J. Math. **6** (2), 117-122 (2016).
21. PeiDe Liu, YouLiang Hou, and MaoFa Wang, *Weak Orlicz space and its applications to the martingale theory*, Science China Mathematics **53** (4), 905-916 (2010).
22. Mizuhara, T.: *Boundedness of some classical operators on generalized Morrey spaces*, Harmonic Analysis, ICM 90 Statellite Proceedings, Springer, Tokyo, 183-189 (1991).
23. Morrey, C.B.: *On the solutions of quasi-linear elliptic partial differential equations*, Trans. Amer. Math. Soc. **43**, 126-166 (1938).
24. Nakai, E.: *Hardy-Littlewood maximal operator, singular integral operators, and the Riesz potential on generalized Morrey spaces*, Math. Nachr. **166**, 95-103 (1994).
25. Nakai, E.: *Generalized fractional integrals on Orlicz-Morrey spaces*, In: Banach and Function Spaces. (Kitakyushu, 2003), Yokohama Publishers, Yokohama, 323-333 (2004).
26. Nakai, E.: *Orlicz-Morrey spaces and Hardy-Littlewood maximal function*, Studia Math. **188**, 193-221 (2008).
27. Nakai, E.: *Orlicz-Morrey spaces and their preduals*, in: Banach and Function spaces (Kitakyushu, 2009) Yokohama Publ. Yokohama 187-205 (2011).
28. Omarova, M.N.: *Parabolic nonsingular integral in Orlicz spaces*, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci. **39** (1), Mathematics, 162-169 (2019).
29. Orlicz, W.: *Über eine gewisse Klasse von Räumen vom Typus B*, Bull. Acad. Polon. A, 207-220 (1932). ; reprinted in: Collected Papers, PWN, Warszawa, 217-230 (1988).
30. Orlicz, W.: *Über Räume (L^M)*, Bull. Acad. Polon. A (1936), 93-107. ; reprinted in: Collected Papers, PWN, Warszawa, 345-359 (1988).
31. Sawano, Y., Sugano, S., Tanaka, H.: *Orlicz-Morrey spaces and fractional operators*, Potential Anal. **36** (4), 517-556 (2012).
32. Sawano, Y.: *A thought on generalized Morrey spaces*, ArXiv Math. 78 pp. (2018).