

Multi-sublinear operators generated by multilinear Calderón-Zygmund operators on product generalized Morrey spaces

Afaq F. Ismayilova* · Fatayi A. Isayev

Received: 08.08.2019 / Revised: 08.03.2020 / Accepted: 05.06.2020

Abstract. *In this paper, we established the boundedness for a large class of multi-sublinear operators T_m generated by multilinear Calderón-Zygmund operators on product generalized Morrey spaces $\mathcal{M}_{p_1, \varphi_1}(\mathbb{R}^n) \times \dots \times \mathcal{M}_{p_m, \varphi_m}(\mathbb{R}^n)$. We find the sufficient conditions on $(\varphi_1, \dots, \varphi_m, \varphi)$ which ensures the boundedness of the operators T_m from $\mathcal{M}_{p_1, \varphi_1}(\mathbb{R}^n) \times \dots \times \mathcal{M}_{p_m, \varphi_m}(\mathbb{R}^n)$ to $\mathcal{M}_{p, \varphi}(\mathbb{R}^n)$ for $1/p = 1/p_1 + \dots + 1/p_m$. The multi-sublinear operators under consideration contain integral operators of harmonic analysis such as multi-sublinear maximal operator \mathcal{M}_m , multilinear Calderón-Zygmund operators \mathcal{T}_m , etc.*

Keywords. Multi-sublinear maximal operator; multilinear Calderón-Zygmund operator; product generalized Morrey space.

Mathematics Subject Classification (2010): 42B20, 42B25, 42B35.

1 Introduction

Multilinear Calderón-Zygmund theory is a natural generalization of the linear case. The initial work on the class of multilinear Calderón-Zygmund operators was done by Coifman and Meyer in [2] and was later systematically studied by Grafakos and Torres in [9, 10].

The classical Morrey spaces, introduced by Morrey [23] in 1938, have been studied intensively by various authors and together with Lebesgue spaces play an important role in the theory of partial differential equations. Although such spaces allow to describe local properties of functions better than Lebesgue spaces, they have some unpleasant issues. It is well known that Morrey spaces are non separable and that the usual classes of nice functions are not dense in such spaces. Moreover, various Morrey spaces are defined in the process of study. Guliyev, Mizuhara and Nakai [11, 21, 24] introduced generalized Morrey spaces $\mathcal{M}_{p, \varphi}(\mathbb{R}^n)$ (see, also [12, 13, 17, 26]). In [13] is defined the generalized Morrey spaces $\mathcal{M}_{p, \varphi}$ with normalized norm

$$\|f\|_{\mathcal{M}_{p, \varphi}} \equiv \sup_{x \in \mathbb{R}^n, r > 0} \varphi(x, r)^{-1} |B(x, r)|^{-1/p} \|f\|_{L_p(B(x, r))},$$

* Corresponding author

A.F. Ismayilova
Azerbaijan University of Cooperation, Baku, Azerbaijan
E-mail: afaismayilova28@gmail.com

F.A. Isayev
Institute of Mathematics and Mechanics of NAS of Azerbaijan, Baku, Azerbaijan
E-mail: isayevfatai@yahoo.com

where the function φ is a positive measurable function on $\mathbb{R}^n \times (0, \infty)$. Here and everywhere in the sequel $B(x, r)$ is the ball in \mathbb{R}^n of radius r centered at x and $|B(x, r)| = v_n r^n$ is its Lebesgue measure, where v_n is the volume of the unit ball in \mathbb{R}^n . In [11] Guliyev also studied the boundedness of the classical operators in these spaces $\mathcal{M}_{p,\varphi}$, see also [3–8, 18, 19, 25].

For $x \in \mathbb{R}^n$ and $r > 0$, we denote by $B(x, r)$ the open ball centered at x of radius r , and by ${}^c B(x, r)$ denote its complement. Let $|B(x, r)|$ be the Lebesgue measure of the ball $B(x, r)$. We denote by \vec{f} the m -tuple (f_1, f_2, \dots, f_m) , $\vec{y} = (y_1, \dots, y_m)$ and $d\vec{y} = dy_1 \cdots dy_m$.

Let $\vec{f} \in L_{p_1}^{loc}(\mathbb{R}^n) \times \dots \times L_{p_m}^{loc}(\mathbb{R}^n)$. The multi-sublinear maximal operator M_m is defined by

$$M_m(\vec{f})(x) = \sup_{r>0} \prod_{i=1}^m \frac{1}{|B(x, r)|} \int_{B(x, r)} |f_i(y_i)| dy_i.$$

In [10] Grafakos and Torres studied the multilinear Calderón-Zygmund operator which can be written for $x \notin \cap_{j=1}^m \text{supp} f_j$ as

$$K_m(\vec{f})(x) = \int_{(\mathbb{R}^n)^m} K(x, y_1, \dots, y_m) f_1(y_1) \cdots f_m(y_m) dy_1 dy_2 \cdots dy_m,$$

where $K(x, y_1, \dots, y_m)$ is the kernel function defined of the diagonal $x = y_1 = \dots = y_m =$ in $(\mathbb{R}^n)^{m+1}$ satisfying

$$|K(y_0, y_1, \dots, y_m)| \leq c_1 \left(\sum_{k,l=0}^m |y_k - y_l| \right)^{-mn},$$

and whenever $2|y_j - y'_j| \leq \frac{1}{2} \max_{0 \leq k \leq m} |y_j - y_k|$,

$$|K(y_0, \dots, y_j, \dots, y_m) - K(y_0, \dots, y'_j, \dots, y_m)| \leq \frac{c_1 |y_j - y'_j|^\epsilon}{\left(\sum_{k,l=0}^m |y_k - y_l| \right)^{mn+\epsilon}},$$

for some $\epsilon > 0$ and all $0 \leq j \leq m$. Grafakos and Torres [10] proved that the operator $K_m(\vec{f})$ is bounded from $L_{p_1}(\mathbb{R}^n) \times \dots \times L_{p_m}(\mathbb{R}^n)$ to $L_p(\mathbb{R}^n)$ for $p_i > 1 (i = 1, \dots, m)$ and $1/p = 1/p_1 + \dots + 1/p_m$, and bounded from $L_1(\mathbb{R}^n) \times \dots \times L_1(\mathbb{R}^n)$ to $L_{\frac{1}{m}, \infty}(\mathbb{R}^n)$.

It is well known that multi-sublinear maximal operator and multilinear Calderón-Zygmund operators play an important role in harmonic analysis (see [1, 10, 22]).

Suppose that T_m represents a multilinear or a multi-sublinear operator, which satisfies that for any $\vec{f} \in L_1(\mathbb{R}^n) \times \dots \times L_1(\mathbb{R}^n)$ with compact support and $x \notin \cap_{j=1}^m \text{supp} f_j$

$$|T_m(\vec{f})(x)| \leq c_0 \int_{(\mathbb{R}^n)^m} \frac{|f_1(y_1) \cdots f_m(y_m)|}{|(x - y_1, \dots, x - y_m)|^{mn}} dy_1 dy_2 \cdots dy_m, \quad (1.1)$$

where c_0 is independent of \vec{f} and x .

The condition (1.1) is satisfied by many interesting operators in harmonic analysis, such as the multilinear Calderón-Zygmund operators, multi-sublinear maximal operator, and so on (see [10, 20] for details).

In this work, we prove the boundedness of the multi-sublinear operator T_m satisfies the condition (1.1) generated by multilinear Calderón-Zygmund operator from product generalized Morrey space $\mathcal{M}_{p_1, \varphi_1} \times \dots \times \mathcal{M}_{p_m, \varphi_m}$ to $\mathcal{M}_{p, \varphi}$, if $1 < p_1, \dots, p_m < \infty$ and $1/p = 1/p_1 + \dots + 1/p_m$, and from the space $\mathcal{M}_{p_1, \varphi_1} \times \dots \times \mathcal{M}_{p_m, \varphi_m}$ to the weak space $WM_{1, \varphi}$, if $1 \leq p_1, \dots, p_m < \infty$, $1/p = 1/p_1 + \dots + 1/p_m$ and at least one p_i equals one (Theorem 2.3). Finally, as applications we apply this result to several particular operators such as the multi-sublinear maximal operator and multilinear Calderón-Zygmund operator.

By $A \lesssim B$ we mean that $A \leq CB$ with some positive constant C independent of appropriate quantities. If $A \lesssim B$ and $B \lesssim A$, we write $A \approx B$ and say that A and B are equivalent.

2 Main Results

In this section, we will discuss the boundedness properties of multi-sublinear operators T_m generated by multilinear Calderón-Zygmund operators on product generalized Morrey spaces $\mathcal{M}_{p_1, \varphi_1}(\mathbb{R}^n) \times \dots \times \mathcal{M}_{p_m, \varphi_m}(\mathbb{R}^n)$.

We find it convenient to define the generalized Morrey spaces in the form as follows.

Definition 2.1 Let $\varphi(x, r)$ be a positive measurable function on $\mathbb{R}^n \times (0, \infty)$ and $1 \leq p < \infty$. We denote by $M_{p, \varphi} \equiv M_{p, \varphi}(\mathbb{R}^n)$ the generalized Morrey space, the space of all functions $f \in L_p^{\text{loc}}(\mathbb{R}^n)$ with finite quasinorm

$$\|f\|_{M_{p, \varphi}} = \sup_{x \in \mathbb{R}^n, r > 0} \varphi(x, r)^{-1} |B(x, r)|^{-\frac{1}{p}} \|f\|_{L_p(B(x, r))}.$$

Also by $WM_{p, \varphi} \equiv WM_{p, \varphi}(\mathbb{R}^n)$ we denote the weak generalized Morrey space of all functions $f \in WL_p^{\text{loc}}(\mathbb{R}^n)$ for which

$$\|f\|_{WM_{p, \varphi}} = \sup_{x \in \mathbb{R}^n, r > 0} \varphi(x, r)^{-1} |B(x, r)|^{-\frac{1}{p}} \|f\|_{WL_p(B(x, r))} < \infty.$$

Lemma 2.1 [4] Let $\varphi(x, r)$ be a positive measurable function on $\mathbb{R}^n \times (0, \infty)$.

(i) If

$$\sup_{t < r < \infty} \frac{r^{-\frac{n}{p}}}{\varphi(x, r)} = \infty \quad \text{for some } t > 0 \text{ and for all } x \in \mathbb{R}^n, \quad (2.1)$$

then $M_{p, \varphi}(\mathbb{R}^n) = \emptyset$.

(ii) If

$$\sup_{0 < r < \tau} \varphi(x, r)^{-1} = \infty \quad \text{for some } \tau > 0 \text{ and for all } x \in \mathbb{R}^n, \quad (2.2)$$

then $M_{p, \varphi}(\mathbb{R}^n) = \emptyset$.

Remark 2.1 We denote by Ω_p the sets of all positive measurable functions φ on $\mathbb{R}^n \times (0, \infty)$ such that for all $t > 0$,

$$\sup_{x \in \mathbb{R}^n} \left\| \frac{r^{-\frac{n}{p}}}{\varphi(x, r)} \right\|_{L_\infty(t, \infty)} < \infty, \quad \text{and} \quad \sup_{x \in \mathbb{R}^n} \left\| \varphi(x, r)^{-1} \right\|_{L_\infty(0, t)} < \infty,$$

respectively. In what follows, keeping in mind Lemma 2.1, we always assume that $\varphi \in \Omega_p$.

We will use the following statements on the boundedness of the weighted Hardy operator

$$H_w g(r) := \int_r^\infty g(t)w(t)dt, \quad 0 < t < \infty,$$

where w is a fixed function non-negative and measurable on $(0, \infty)$. The following theorem was proved in [14] (see also [16]).

Theorem 2.1 [14] *Let v_1, v_2 and w be positive almost everywhere and measurable functions on $(0, \infty)$. The inequality*

$$\operatorname{ess\,sup}_{r>0} v_2(r)H_w g(r) \leq C \operatorname{ess\,sup}_{r>0} v_1(r)g(r) \quad (2.3)$$

holds for some $C > 0$ for all non-negative and non-decreasing g on $(0, \infty)$ if and only if

$$B := \sup_{r>0} v_2(r) \int_r^\infty \frac{w(t)dt}{\sup_{t<s<\infty} v_1(s)} < \infty. \quad (2.4)$$

Moreover, the value $C = B$ is the best constant for (2.3).

Remark 2.2 In (2.3) – (2.4) it is assumed that $0 \cdot \infty = 0$.

In the following lemma we get Guliyev local estimate (see, for example, [11–13] in the case $m = 1$ and [15] in the case $m > 1$) for the operator T_m .

Theorem 2.2 *Let $1 \leq p_1, \dots, p_m < \infty$ and $1/p = 1/p_1 + \dots + 1/p_m$. Let T_m be a multi-sublinear operator which satisfies the condition (1.1) bounded from $L_{p_1}(\mathbb{R}^n) \times \dots \times L_{p_m}(\mathbb{R}^n)$ to $L_p(\mathbb{R}^n)$ for $p_i > 1, i = 1, \dots, m$, and bounded from $L_{p_1}(\mathbb{R}^n) \times \dots \times L_{p_m}(\mathbb{R}^n)$ to $WL_p(\mathbb{R}^n)$ for $p_i \geq 1, i = 1, \dots, m$.*

Then, for $1 < p_1, \dots, p_m < \infty$ the inequality

$$\|T_m(\vec{f})\|_{L_p(B(x_0, r))} \lesssim r^{\frac{n}{p}} \prod_{i=1}^m \int_{2r}^\infty t^{-\frac{n}{p_i}-1} \|f_i\|_{L_{p_i}(B(x_0, t))} dt \quad (2.5)$$

holds for any ball $B(x_0, r)$ and for all $\vec{f} \in L_{p_1}^{loc}(\mathbb{R}^n) \times \dots \times L_{p_m}^{loc}(\mathbb{R}^n)$.

Moreover, if at least one p_i equals one, the inequality

$$\|T_m(\vec{f})\|_{WL_p(B(x_0, r))} \lesssim r^{\frac{n}{p}} \prod_{i=1}^m \int_{2r}^\infty t^{-\frac{n}{p_i}-1} \|f_i\|_{L_{p_i}(B(x_0, t))} dt \quad (2.6)$$

holds for any ball $B(x_0, r)$ and for all $\vec{f} \in L_{p_1}^{loc}(\mathbb{R}^n) \times \dots \times L_{p_m}^{loc}(\mathbb{R}^n)$.

Proof. Let $1 < p_1, \dots, p_m < \infty$ and $1/p = 1/p_1 + \dots + 1/p_m$. For arbitrary $x_0 \in \mathbb{R}^n$, set $B = B(x_0, r)$ for the ball centered at x_0 and of radius r , $2B = B(x_0, 2r)$. We represent $\vec{f} = (f_1, \dots, f_m)$ as

$$f_j = f_j^0 + f_j^\infty, \quad f_j^0 = f_j \chi_{2B}, \quad f_j^\infty = f_j \chi_{\mathbb{R}^n \setminus 2B}, \quad j = 1, \dots, m. \quad (2.7)$$

Then we split $T_m(\vec{f})$ as follows

$$\left| T_m(\vec{f})(x) \right| \leq c_0 \left| T_m(f_1^0, \dots, f_m^0)(x) \right| + \left| \sum_{\beta_1, \dots, \beta_m} T_m(f_1^{\beta_1}, \dots, f_m^{\beta_m})(x) \right|,$$

where $\beta_1, \dots, \beta_m \in \{0, \infty\}$ and each term of \sum' contains at least $\beta_i \neq 0$. Then,

$$\begin{aligned} \|T_m(\vec{f})\|_{L_p(B(x,r))} &\leq \|T_m(f_1^0, \dots, f_m^0)\|_{L_p(B(x,r))} \\ &+ \left\| \sum'_{\beta_1, \dots, \beta_m} T_m(f_1^{\beta_1}, \dots, f_m^{\beta_m}) \right\|_{L_p(B(x,r))} \leq I + II. \end{aligned}$$

For I , by the boundedness of T_m from product $L_{p_1}(\mathbb{R}^n) \times \dots \times L_{p_m}(\mathbb{R}^n)$ to $L_p(\mathbb{R}^n)$ with $1/p = 1/p_1 + \dots + 1/p_m$ for each $p_i > 1$ ($i = 1, \dots, m$), we have,

$$\begin{aligned} \|T_m(\vec{f}^0)\|_{L_p(B(x,r))} &\leq \|T_m(\vec{f}^0)\|_{L_p(\mathbb{R}^n)} \\ &\lesssim \prod_{i=1}^m \|f_i^0\|_{L_{p_i}(\mathbb{R}^n)} \lesssim \prod_{i=1}^m \|f_i\|_{L_{p_i}(B(x,2r))}. \end{aligned}$$

Taking into account that

$$\|f_i\|_{L_{p_i}(B(x,2r))} \lesssim r^{\frac{n}{p_i}} \int_{2r}^{\infty} t^{-\frac{n}{p_i}-1} \|f_i\|_{L_{p_i}(B(x,t))} dt, \quad i = 1, \dots, m$$

we get

$$\|T_m(\vec{f}^0)\|_{L_p(B(x,r))} \lesssim r^{\frac{n}{p}} \prod_{i=1}^m \int_{2r}^{\infty} t^{-\frac{n}{p_i}-1} \|f_i\|_{L_{p_i}(B(x,t))} dt. \quad (2.8)$$

For II , first we consider the case $\beta_1 = \dots = \beta_m = \infty$.

When $|x - y_i| \leq r$, $|z - y_i| \geq 2r$, we have $\frac{1}{2}|z - y_i| \leq |x - y_i| \leq \frac{3}{2}|z - y_i|$, and so by the condition (1.1) we have

$$\begin{aligned} |T_m(\vec{f}^{\infty})(z)| &\lesssim \int_{(\mathfrak{c}_{B(x,2r)})^m} \frac{|f_1(y_1) \cdots f_m(y_m)|}{|(x - y_1, \dots, x - y_m)|^{mn}} d\vec{y} \\ &\lesssim \prod_{i=1}^m \int_{\mathfrak{c}_{B(x,2r)}} \frac{|f_i(y_i)|}{|x - y_i|^n} dy_i \end{aligned}$$

and

$$\begin{aligned} \|T_m(\vec{f}^{\infty})\|_{L_p(B(x,r))} &\leq \prod_{i=1}^m \int_{\mathfrak{c}_{B(x,2r)}} \frac{|f_i(y_i)|}{|x - y_i|^n} dy_i \|\chi_{B(x,r)}\|_{L_p(\mathbb{R}^n)} \\ &\lesssim r^{\frac{n}{p}} \prod_{i=1}^m \int_{\mathfrak{c}_{B(x,2r)}} \frac{|f_i(y_i)|}{|x - y_i|^n} dy_i. \end{aligned}$$

By Fubini's theorem we have

$$\begin{aligned} \int_{\mathfrak{c}_{B(x,2r)}} \frac{|f_i(y_i)|}{|x - y_i|^n} dy_i &\approx \int_{\mathfrak{c}_{B(x,2r)}} |f_i(y_i)| \int_{|x_0 - y_i|}^{\infty} \frac{dt}{t^{n+1}} dy_i \\ &\approx \int_{2r}^{\infty} \int_{2r \leq |x_0 - y_i| < t} |f_i(y_i)| dy_i \frac{dt}{t^{n+1}} \\ &\lesssim \int_{2r}^{\infty} \int_{B(x_0,t)} |f_i(y_i)| dy_i \frac{dt}{t^{n+1}}. \end{aligned}$$

Applying Hölder's inequality, we get

$$\int_{\mathfrak{C}_{B(x,2r)}} \frac{|f_i(y_i)|}{|x-y_i|^n} dy_i \lesssim \int_{2r}^{\infty} t^{-\frac{n}{p_i}-1} \|f_i\|_{L_{p_i}(B(x,t))} dt. \quad (2.9)$$

Moreover, for all $p_i \in [1, \infty)$, $i = 1, \dots, m$ the inequality

$$\|T_m(\vec{f}^{\infty})\|_{L_p(B(x,r))} \lesssim r^{\frac{n}{p}} \prod_{i=1}^m \int_{2r}^{\infty} t^{-\frac{n}{p_i}-1} \|f_i\|_{L_{p_i}(B(x,t))} dt \quad (2.10)$$

is valid.

Next we consider the case that some $\alpha_i = 0$ and other $\alpha_j = \infty$. To this end we may assume that $\alpha_1 = \alpha_2 = \infty$ and $\alpha_3 = \dots = \alpha_m = 0$. Recall the condition (1.1) and the fact that $|x-y_i| \approx |z-y_i|$ for $z \in B(x,r)$ and $y_i \in \mathfrak{C}_{B(x,2r)}$, we have that

$$\begin{aligned} & T(f_1^{\infty}, f_2^{\infty}, f_3^0, \dots, f_m^0)(z) \\ & \lesssim \int_{\mathfrak{C}_{B(x,2r)} \times \mathfrak{C}_{B(x,2r)}} \frac{|f_1(y_1)| |f_2(y_2)|}{(|x-y_1| + |x-y_2|)^{mn}} dy_1 dy_2 \prod_{i=3}^m \int_{B(x,2r)} |f_i(y_i)| dy_i \\ & \lesssim \int_{\mathfrak{C}_{B(x,2r)}} \frac{|f_1(y_1)|}{|x-y_1|^n} dy_1 \int_{\mathfrak{C}_{B(x,2r)}} \frac{|f_2(y_2)|}{|x-y_2|^n} dy_2 \prod_{i=3}^m \int_{B(x,2r)} |f_i(y_i)| dy_i. \end{aligned}$$

By the inequality (2.9) and use the Hölder's inequality for integrals, we get

$$\begin{aligned} & \|T(f_1^{\infty}, f_2^{\infty}, f_3^0, \dots, f_m^0)\|_{L_p(B(x,r))} \\ & \lesssim r^{\frac{n}{p}} \int_{\mathfrak{C}_{B(x,2r)}} \frac{|f_1(y_1)|}{|x-y_1|^n} dy_1 \int_{\mathfrak{C}_{B(x,2r)}} \frac{|f_2(y_2)|}{|x-y_2|^n} dy_2 \prod_{i=3}^m \int_{B(x,2r)} |f_i(y_i)| dy_i \\ & \leq r^{\frac{n}{p}} \prod_{i=3}^m \int_r^{\infty} t^{-\frac{n}{p_i}-1} \|f_i\|_{L_{p_i}(B(x,t))} dt. \end{aligned}$$

For the proof of the inequality (2.6), by a similar argument as in the proof of (2.5) and pay attention to the fact that $\vec{f} \rightarrow T_m(\vec{f})$ is bounded from $L_{p_1}(\mathbb{R}^n) \times \dots \times L_{p_m}(\mathbb{R}^n)$ to $WL_p(\mathbb{R}^n)$, we can similarly prove (2.6) and we omit the details here.

Now we give the boundedness of multi-sublinear operators generated by multilinear Calderón-Zygmund operators on product generalized Morrey space.

Theorem 2.3 *Let $1 \leq p_1, \dots, p_m < \infty$ with $1/p = 1/p_1 + \dots + 1/p_m$ and $(\varphi_1, \dots, \varphi_m, \varphi) \in \Omega_{p_1} \times \dots \times \Omega_{p_1} \times \Omega_p$ satisfies the condition*

$$\prod_{i=1}^m \int_r^{\infty} \frac{\operatorname{ess\,inf}_{t < s < \infty} \varphi_i(x, s) s^{\frac{n}{p_i}}}{t^{\frac{n}{p_i} + 1}} dt \lesssim \varphi(x, r). \quad (2.11)$$

Let also T_m be a multi-sublinear operator which satisfies the condition (1.1) and bounded from $L_{p_1}(\mathbb{R}^n) \times \dots \times L_{p_m}(\mathbb{R}^n)$ to $L_p(\mathbb{R}^n)$ for $p_i > 1$, $i = 1, \dots, m$, and bounded from $L_{p_1}(\mathbb{R}^n) \times \dots \times L_{p_m}(\mathbb{R}^n)$ to $WL_p(\mathbb{R}^n)$ for $p_i \geq 1$, $i = 1, \dots, m$. Then the operator T_m is bounded from product space $\mathcal{M}_{p_1, \varphi_1}(\mathbb{R}^n) \times \dots \times \mathcal{M}_{p_m, \varphi_m}(\mathbb{R}^n)$ to $\mathcal{M}_{p, \varphi}(\mathbb{R}^n)$ for $p_i > 1$, $i = 1, \dots, m$, and from product space $\mathcal{M}_{p_1, \varphi_1}(\mathbb{R}^n) \times \dots \times \mathcal{M}_{p_m, \varphi_m}(\mathbb{R}^n)$ to $W\mathcal{M}_{p, \varphi}(\mathbb{R}^n)$ for at least one p_i equals one.

Proof. Let $1 < p_1, \dots, p_m < \infty$ and $\vec{f} \in \mathcal{M}_{p_1, \varphi_1}(\mathbb{R}^n) \times \dots \times \mathcal{M}_{p_m, \varphi_m}(\mathbb{R}^n)$. By Theorems 2.1 and 2.2 we have

$$\begin{aligned} \|T_{\alpha, m}(\vec{f})\|_{\mathcal{M}_{p, \varphi}} &\lesssim \sup_{x \in \mathbb{R}^n, r > 0} \varphi(x, r)^{-1} \prod_{i=1}^m \int_r^\infty t^{-\frac{n}{p_i}-1} \|f_i\|_{L_{p_i}(B(x, t))} dt \\ &\lesssim \prod_{i=1}^m \sup_{x \in \mathbb{R}^n, r > 0} \varphi_i(x, r)^{-1} r \|f_i\|_{L_{p_i}(B(x, r))} = \prod_{i=1}^m \|f_i\|_{\mathcal{M}_{p_i, \varphi_i}}. \end{aligned}$$

When $p_i = 1, i = 1, \dots, m$, the proof is similar and we omit the details here.

Corollary 2.1 [15] *Let $1 \leq p_1, \dots, p_m < \infty$ with $1/p = 1/p_1 + \dots + 1/p_m$. Let also $(\varphi_1, \dots, \varphi_m, \varphi) \in \Omega_{p_1} \times \dots \times \Omega_{p_1} \times \Omega_p$ satisfies the condition (2.11). Then the operators M_m and K_m are bounded from product space $\mathcal{M}_{p_1, \varphi_1}(\mathbb{R}^n) \times \dots \times \mathcal{M}_{p_m, \varphi_m}(\mathbb{R}^n)$ to $\mathcal{M}_{p, \varphi}(\mathbb{R}^n)$ for $p_i > 1, i = 1, \dots, m$ and from product space $\mathcal{M}_{p_1, \varphi_1}(\mathbb{R}^n) \times \dots \times \mathcal{M}_{p_m, \varphi_m}(\mathbb{R}^n)$ to $W\mathcal{M}_{p, \varphi}(\mathbb{R}^n)$ for at least one p_i equals one.*

Acknowledgements The authors thank the referee(s) for careful reading the paper and useful comments. The research of F. Isayev was partially supported by the Grant of 1st Azerbaijan-Russia Joint Grant Competition (Agreement Number No. EIF-BGM-4-RFTF1/2017-21/01/1).

References

1. Coifman R., Grafakos L.: *Hardy spaces estimates for multilinear operators I*, Rev. Math. Iberoamericana, **8**, 45-68 (1992).
2. Coifman R.R., Meyer Y.: *On commutators of singular integrals and bilinear singular integrals*, Trans. Amer. Math. Soc. **212**, 315-331 (1975).
3. Eroglu A., Hasanov A.A., Omarova M.N.: *(p, q)-admissible multilinear fractional integral operators and their commutators in product generalized local Morrey spaces*, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci. **36** (4), Mathematics, 77-98 (2016).
4. Eroglu, A., Guliyev, V.S., Azizov, C.V.: *Characterizations for the fractional integral operators in generalized Morrey spaces on Carnot groups*, Math. Notes. **102** (5-6), 722-734 (2017).
5. Eroglu A, Azizov C.V.: *A note on the fractional integral operators in generalized Morrey spaces on the Heisenberg group*, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci. **37** (1), Mathematics, 86-91 (2017).
6. Eroglu A, Isayev F.A., Namazov F.M.: *Fractional integral associated to Schrödinger operator on the Heisenberg groups in vanishing generalized Morrey spaces*, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci. **38** (4), Mathematics, 54-62 (2018).
7. Eroglu A, Azizov C.V., Guliyev, V.S.: *Fractional maximal operator and its commutators in generalized Morrey spaces on Heisenberg group*, Proc. Inst. Math. Mech. Natl. Acad. Sci. Azerb. **44** (2), 304-317 (2018).
8. Gadjev T., Aliev S., Galandarova Sh.: *A priori estimates for solutions to Dirichlet boundary value problems for polyharmonic equations in generalized Morrey spaces*, TWMS J. Pure Appl. Math. **9** (2), 231-242 (2018).
9. Grafakos L., Torres R.H.: *Multilinear Calderón-Zygmund theory*, Adv. Math. **165** (1), 124-164 (2002).
10. Grafakos L., Torres R.H.: *On multilinear singular integrals of Calderón-Zygmund type*, Publ. Mat. Vol. Extra, 57-91 (2002).

11. Guliyev V.S.: *Integral operators on function spaces on the homogeneous groups and on domains in \mathbb{R}^n* , (Russian) Doctor's degree dissertation, Mat. Inst. Steklov, Moscow, 329 pp. (1994).
12. Guliyev V.S.: *Function spaces, Integral Operators and Two Weighted Inequalities on Homogeneous Groups. Some Applications*, (Russian) Casioglu, Baku, 332 pp. (1999).
13. Guliyev V. S.: *Boundedness of the maximal, potential and singular operators in the generalized Morrey spaces*, J. Inequal. Appl. Art. ID 503948, 20 pp. (2009).
14. Guliyev V.S.: *Generalized weighted Morrey spaces and higher order commutators of sublinear operators*, Eurasian Math. J. **3** (3), 33-61 (2012).
15. Guliyev V.S., Ismayilova A.F.: *Multi-sublinear maximal operator and multilinear singular integral operators on generalized Morrey spaces*, Proc. Inst. Math. Mech. Natl. Acad. Sci. Azerb. **40** (2), 65-77 (2014).
16. Guliyev V.S., Omarova M.N., Ragusa M.A., Scapellato A.: *Commutators and generalized local Morrey spaces*, J. Math. Anal. Appl. **457** (2), 1388-1402 (2018).
17. Guliyev V.S., Guliyev R.V., Omarova, M.N.: *Riesz transforms associated with Schrödinger operator on vanishing generalized Morrey spaces*, Appl. Comput. Math. **17** (1), 56-71 (2018).
18. Guliyev V.S., Akbulut A., Celik S., Omarova M.N.: *Higher order Riesz transforms related to Schrödinger type operator on local generalized Morrey spaces*, TWMS J. Pure Appl. Math. **10** (1), 58-75 (2019).
19. Ibrahimov E.J., Dadashova G.A., Ekincioglu S.E.: *On the boundedness of the G -maximal operator and G -Riesz potential in the generalized G -Morrey spaces*, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci. **40** (1), Mathematics, 111-125 (2020).
20. Lin Yan, Lu Shanzhen: *Multilinear Calderón-Zygmund operator on Morrey type spaces*, Anal. Theory Appl. **22**, 387-400 (2006).
21. Mizuhara T.: *Boundedness of some classical operators on generalized Morrey spaces*, Harmonic Analysis (S. Igari, Editor), ICM 90 Satellite Proceedings, Springer - Verlag, Tokyo, 183-189 (1991).
22. Moen K.: *Weighted inequalities for multilinear fractional integral operators*, Collect. Math, **60**, 213-238 (2009).
23. Morrey C.B.: *On the solutions of quasi-linear elliptic partial differential equations*, Trans. Amer. Math. Soc, **43**, 126-166 (1938).
24. Nakai E.: *Hardy-Littlewood maximal operator, singular integral operators and Riesz potentials on generalized Morrey spaces*, Math. Nachr, **166**, 95-103 (1994).
25. Rahimova K.R.: *Characterizations of parabolic fractional integral operators on generalized parabolic Morrey spaces*, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci. **36** (4), Mathematics, 156-166 (2016).
26. Sawano Y.: *A thought on generalized Morrey spaces*, J. Indonesian Math. Soc. **25** (3), 210-281 (2019).