

## Calderón-Zygmund operators with kernels of Dini's type and their multilinear commutators on generalized Morrey spaces

Afag F. Ismayilova

Received: 28.12.2020 / Revised: 12.03.2021 / Accepted: 25.04.2021

**Abstract.** *In this paper, we obtain the endpoint boundedness for the Calderón-Zygmund operators with kernels of Dini's type on generalized Morrey spaces. We also get similar results for the multilinear commutators of Calderón-Zygmund operators with kernels of Dini's type with BMO functions.*

**Keywords.** Generalized Morrey spaces; Calderón-Zygmund operator; commutator; BMO.

**Mathematics Subject Classification (2010):** 42B20, 42B35.

### 1 Introduction

The theory of Calderón-Zygmund operators has played very important roles in modern harmonic analysis with lots of extensive applications in the others fields of mathematics, which has been extensively studied (see [2, 3, 4, 20, 21, 28]). In particular, Yabuta introduced certain  $\omega$ -type Calderón-Zygmund operators to facilitate his study of certain classes of pseudodifferential operators (see [31]). Let  $\omega$  be a non-negative and non-decreasing function on  $\mathbb{R}_+ = (0, \infty)$ . We say that  $\omega$  satisfies the *Dini* condition and write  $\omega \in \text{Dini}$ , if

$$\int_0^\infty \frac{\omega(t)}{t} dt < \infty. \quad (1.1)$$

A measurable function  $K(\cdot, \cdot)$  on  $\mathbb{R}^n \times \mathbb{R}^n$  is said to be a  $\omega$ -type Calderón-Zygmund kernel if it satisfies

$$|K(x, y)| \leq C |x - y|^{-n} \quad (1.2)$$

and for all distinct  $x, y \in \mathbb{R}^n$ , and all  $z$  with  $2|x - z| < |x - y|$ , there exist positive constants  $C$  and  $\gamma$  such that

$$|K(x, y) - K(z, y)| + |K(y, x) - K(y, z)| \leq C\omega\left(\frac{|x - z|}{|x - y|}\right) |x - y|^{-n}. \quad (1.3)$$

**Definition 1.1** *Let  $T$  be a linear operator from  $\mathcal{S}(\mathbb{R}^n)$  into its dual  $\mathcal{S}'(\mathbb{R}^n)$ , where  $\mathcal{S}(\mathbb{R}^n)$  denotes the Schwartz class. One can say that  $T$  is a  $\omega$ -type Calderón-Zygmund operator if it satisfies the following conditions:*

i)  $T$  can be extended to be a bounded linear operator on  $L_2(\mathbb{R}^n)$ ;

ii) there is a  $\omega$ -type Calderón-Zygmund kernel  $K(x, y)$  such that

$$Tf(x) = \int_{\mathbb{R}^n} K(x, y)f(y)dy, \text{ as } f \in C_c^\infty \text{ and } x \notin \text{supp } f. \quad (1.4)$$

It is easy to see that the classical Calderón-Zygmund operator with standard kernel is a special case of  $\omega$ -type operator  $T$  as  $\omega(t) = t^\varepsilon$  with  $0 < \varepsilon \leq 1$ . Given a locally integrable function  $b$ , the commutator generated by  $T$  and  $b$  is defined by

$$[b, T]f(x) = b(x)Tf(x) - T(bf)(x) = \int_{\mathbb{R}^n} [b(x) - b(y)]K(x, y)f(y)dy. \quad (1.5)$$

Let  $\mathbf{b} = (b_1, \dots, b_m)$  and  $b_j, 1 \leq j \leq m$  be locally integrable functions when we consider multilinear commutators as defined by

$$T_{\mathbf{b}}f(x) = \int_{\mathbb{R}^n} \prod_{j=1}^m (b_j(x) - b_j(y))K(x, y)f(y)dy. \quad (1.6)$$

Furthermore, if we take  $b_i = b, i = 1, \dots, m$ , then we define the following integral equation

$$T_{\mathbf{b}}f(x) = \int_{\mathbb{R}^n} (b(x) - b(y))^m K(x, y)f(y)dy = [b, T]^m f(x).$$

It is well known that Calderón-Zygmund operators play an important role in harmonic analysis (see [28]).

The classical Morrey spaces were introduced by Morrey [23] to study the local behavior of solutions to second-order elliptic partial differential equations. Moreover, various Morrey spaces are defined in the process of study. Guliyev, Mizuhara and Nakai [8, 24, 25] introduced generalized Morrey spaces  $M_{p,\varphi}(\mathbb{R}^n)$  (see, also [1, 5, 6, 9, 10, 13, 14, 15, 16, 17, 18, 27]).

The main purpose of this paper is to establish a number of results concerning generalized Morrey boundedness of Calderón-Zygmund operators with kernels of mild regularity. Let  $T$  be a linear Calderón-Zygmund operator of type  $\omega(t)$  with  $\omega$  being nondecreasing and  $\omega \in Dini$ , but without assuming to be concave. We show that the  $\omega$ -type Calderón-Zygmund operators  $T$  and their multilinear commutators  $T_{\mathbf{b}}$  are bounded from one generalized Morrey space  $M_{p,\varphi_1}$  to another  $M_{p,\varphi_2}$ ,  $1 < p < \infty$ . We find the sufficient conditions on the pair  $(\varphi_1, \varphi_2)$  with  $\mathbf{b} \in BMO^m(\mathbb{R}^n)$  which ensures the boundedness of the operators  $T$  and  $T_{\mathbf{b}}$  from  $M_{p,\varphi_1}$  to  $M_{p,\varphi_2}$  for  $1 < p < \infty$ .

By  $A \lesssim B$  we mean that  $A \leq CB$  with some positive constant  $C$  independent of appropriate quantities. If  $A \lesssim B$  and  $B \lesssim A$ , we write  $A \approx B$  and say that  $A$  and  $B$  are equivalent.

## 2 Generalized Morrey spaces

We define the generalized Morrey spaces as follows.

**Definition 2.1** Let  $1 \leq p < \infty$ ,  $\varphi$  be a positive measurable function on  $\mathbb{R}^n \times (0, \infty)$ . We denote by  $M_{p,\varphi}$  the generalized Morrey space, the space of all functions  $f \in L_p^{loc}(\mathbb{R}^n)$  with finite norm

$$\|f\|_{M_{p,\varphi}} = \sup_{x \in \mathbb{R}^n, r > 0} \varphi(x, r)^{-1} |B(x, r)|^{-\frac{1}{p}} \|f\|_{L_p(B(x, r))},$$

where  $L_p(B(x, r))$  denotes the weighted  $L_p$ -space of measurable functions  $f$  for which

$$\|f\|_{L_p(B(x, r))} \equiv \|f\chi_{B(x, r)}\|_{L_p(\mathbb{R}^n)} = \left( \int_{B(x, r)} |f(y)|^p w(y) dy \right)^{\frac{1}{p}}.$$

Furthermore, by  $WM_{p, \varphi}$  we denote the weak generalized weighted Morrey space of all functions  $f \in WL_p^{\text{loc}}(\mathbb{R}^n)$  for which

$$\|f\|_{WM_{p, \varphi}} = \sup_{x \in \mathbb{R}^n, r > 0} \varphi(x, r)^{-1} |B(x, r)|^{-\frac{1}{p}} \|f\|_{WL_p(B(x, r))} < \infty,$$

where  $WL_p(B(x, r))$  denotes the weak  $L_p$ -space of measurable functions  $f$  for which

$$\|f\|_{WL_p(B(x, r))} \equiv \|f\chi_{B(x, r)}\|_{WL_p(\mathbb{R}^n)} = \sup_{t > 0} t \left( \int_{\{y \in B(x, r) : |f(y)| > t\}} w(y) dy \right)^{\frac{1}{p}}.$$

**Remark 2.1** If  $\varphi(x, r) = r^{\frac{\lambda-n}{p}}$  with  $0 < \lambda < n$ , then  $M_{p, \varphi} = L_{p, \lambda}(\mathbb{R}^n)$  is the classical Morrey space and  $WM_{p, \varphi} = WL_{p, \lambda}(\mathbb{R}^n)$  is the weak Morrey space; If  $\varphi(x, r) \equiv |B(x, r)|^{-\frac{1}{p}}$ , then  $M_{p, \varphi} = L_p(\mathbb{R}^n)$  is the Lebesgue space.

We will use the following statement on the boundedness of the weighted Hardy operator

$$H_w g(t) := \int_t^\infty g(s) w(s) ds, \quad H_w^* g(t) := \int_t^\infty \left(1 + \frac{s}{t}\right) g(s) w(s) ds, \quad 0 < t < \infty,$$

where  $w$  is a weight. The following theorem was proved in [12].

**Theorem 2.1** [12] Let  $v_1, v_2$  and  $w$  be weights on  $(0, \infty)$  and  $v_1(t)$  be bounded outside a neighborhood of the origin. The inequality

$$\sup_{t > 0} v_2(t) H_w g(t) \leq C \sup_{t > 0} v_1(t) g(t)$$

holds for some  $C > 0$  for all non-negative and non-decreasing  $g$  on  $(0, \infty)$  if and only if

$$B := \sup_{t > 0} v_2(t) \int_t^\infty \frac{w(s) ds}{\sup_{s < \tau < \infty} v_1(\tau)} < \infty.$$

**Theorem 2.2** [11] Let  $v_1, v_2$  and  $w$  be weights on  $(0, \infty)$  and  $v_1(t)$  be bounded outside a neighborhood of the origin. The inequality

$$\sup_{t > 0} v_2(t) H_w^* g(t) \leq C \sup_{t > 0} v_1(t) g(t)$$

holds for some  $C > 0$  for all non-negative and non-decreasing  $g$  on  $(0, \infty)$  if and only if

$$B := \sup_{t > 0} v_2(t) \int_t^\infty \left(1 + \frac{s}{t}\right) \frac{w(s) ds}{\sup_{s < \tau < \infty} v_1(\tau)} < \infty.$$

### 3 $\omega$ -type Calderón-Zygmund operators in the spaces $M_{p,\varphi}(\mathbb{R}^n)$

The following theorem was proved in [26].

**Theorem 3.1** [26] *Let  $1 \leq p < \infty$  and  $T$  be  $\omega$ -type Calderón-Zygmund operator defined by (1.4) with  $\omega$  satisfies (1.1). Then, the operator  $T$  is bounded on  $L_p(\mathbb{R}^n)$  for  $p > 1$  and bounded from  $L_1(\mathbb{R}^n)$  into  $WL_1(\mathbb{R}^n)$  for  $p = 1$ .*

The following Guliyev local estimates are valid (see [10]).

**Theorem 3.2** *Let  $1 \leq p < \infty$  and  $T$  be  $\omega$ -type Calderón-Zygmund operator defined by (1.4) with  $\omega$  satisfies (1.1). Then, for  $p > 1$  the inequality*

$$\|Tf\|_{L_p(B)} \lesssim |B|^{\frac{1}{p}} \int_{2r}^{\infty} \|f\|_{L_p(B(x_0,t))} |B(x_0,t)|^{-\frac{1}{p}} \frac{dt}{t}$$

holds for any ball  $B = B(x_0, r)$  and for all  $f \in L_p^{\text{loc}}(\mathbb{R}^n)$ .

Moreover, for  $p = 1$  the inequality

$$\|Tf\|_{WL_1(B)} \lesssim |B| \int_{2r}^{\infty} \|f\|_{L_1(B(x_0,t))} |B(x_0,t)|^{-1} \frac{dt}{t} \quad (3.1)$$

holds for any ball  $B = B(x_0, r)$  and for all  $f \in L_1^{\text{loc}}(\mathbb{R}^n)$ .

**Proof.** Let  $p \in (1, \infty)$ . For arbitrary  $x_0 \in \mathbb{R}^n$ , set  $B = B(x_0, r)$  for the ball centered at  $x_0$  and of radius  $r$ ,  $2B = B(x_0, 2r)$ . We represent  $f$  as

$$f = f_1 + f_2, \quad f_1(y) = f(y)\chi_{2B}(y), \quad f_2(y) = f(y)\chi_{\mathbb{R}^n \setminus 2B}(y), \quad r > 0. \quad (3.2)$$

Then we have

$$\|Tf\|_{L_p(B)} \leq \|Tf_1\|_{L_p(B)} + \|Tf_2\|_{L_p(B)}.$$

Since  $f_1 \in L_p$ ,  $Tf_1 \in L_p$  and from the boundedness of  $T$  in  $L_p$  (see Theorem (3.1)) it follows that

$$\|Tf_1\|_{L_p(B)} \leq \|Tf_1\|_{L_p} \leq C\|f_1\|_{L_p} = C\|f\|_{L_p(2B)},$$

where constant  $C > 0$  is independent of  $f$ .

It is clear that  $x \in B$ ,  $y \in \mathbb{R}^n \setminus 2B$  implies  $\frac{1}{2}|x_0 - y| \leq |x - y| \leq \frac{3}{2}|x_0 - y|$ . We get

$$|Tf_2(x)| \leq 2^n c_0 \int_{\mathbb{R}^n \setminus 2B} \frac{|f(y)|}{|x_0 - y|^n} dy \lesssim \int_{2r}^{\infty} \int_{B(x_0,t)} |f(y)| dy \frac{dt}{t^{n+1}}.$$

Applying Hölder's inequality, we get

$$\int_{\mathbb{R}^n \setminus 2B} \frac{|f(y)|}{|x_0 - y|^n} dy \lesssim \int_{2r}^{\infty} \|f\|_{L_p(B(x_0,t))} |B(x_0,t)|^{-\frac{1}{p}} \frac{dt}{t}. \quad (3.3)$$

Moreover, for all  $p \in [1, \infty)$  the inequality

$$\|Tf_2\|_{L_p(B)} \lesssim |B|^{\frac{1}{p}} \int_{2r}^{\infty} \|f\|_{L_p(B(x_0,t))} |B(x_0,t)|^{-\frac{1}{p}} \frac{dt}{t} \quad (3.4)$$

is valid. Thus

$$\begin{aligned} \|Tf\|_{L_p(B)} &\lesssim \|f\|_{L_p(2B)} + |B|^{\frac{1}{p}} \int_{2r}^{\infty} \|f\|_{L_p(B(x_0,t))} |B(x_0,t)|^{-\frac{1}{p}} \frac{dt}{t} \\ &\lesssim |B|^{\frac{1}{p}} \int_{2r}^{\infty} \|f\|_{L_p(B(x_0,t))} |B(x_0,t)|^{-\frac{1}{p}} \frac{dt}{t}. \end{aligned}$$

Let  $p = 1$ . From the weak  $(1, 1)$  boundedness of  $T$  it follows that:

$$\begin{aligned} \|Tf_1\|_{WL_1(B)} &\leq \|Tf_1\|_{WL_1} \lesssim \|f_1\|_{L_1} = \|f\|_{L_1(2B)} \\ &\lesssim |B| \int_{2r}^{\infty} \|f\|_{L_1(B(x_0,t))} |B(x_0,t)|^{-1} \frac{dt}{t}. \end{aligned} \quad (3.5)$$

Then by (3.4) and (3.5) we get the inequality (3.1).

**Theorem 3.3** *Let  $1 \leq p < \infty$ ,  $T$  be  $\omega$ -type Calderón-Zygmund operator defined by (1.4) with  $\omega$  satisfies (1.1), and  $(\varphi_1, \varphi_2)$  satisfy the condition*

$$\int_r^{\infty} \frac{\operatorname{ess\,inf}_{t < s < \infty} \varphi_1(x, s) |B(x, s)|^{1/p}}{|B(x, t)|^{1/p}} \frac{dt}{t} \leq C \varphi_2(x, r) \quad (3.6)$$

where  $C$  does not depend on  $x$  and  $r$ . Then the operator  $T$  is bounded from  $M_{p,\varphi_1}$  to  $M_{p,\varphi_2}$  for  $p > 1$  and from  $M_{1,\varphi_1}$  to  $WM_{1,\varphi_2}$  for  $p = 1$ .

**Proof.** For  $p > 1$  from Theorem 2.1 and Theorem 3.2 we get

$$\begin{aligned} \|Tf\|_{M_{p,\varphi_2}} &\lesssim \sup_{x \in \mathbb{R}^n, r > 0} \varphi_2(x, r)^{-1} \int_r^{\infty} \|f\|_{L_p(B(x_0,t))} |B(x, t)|^{-\frac{1}{p}} \frac{dt}{t} \\ &\lesssim \sup_{x \in \mathbb{R}^n, r > 0} \varphi_1(x, r)^{-1} |B|^{-\frac{1}{p}} \|f\|_{L_p(B)} = \|f\|_{M_{p,\varphi_1}} \end{aligned}$$

and for  $p = 1$

$$\begin{aligned} \|Tf\|_{WM_{1,\varphi_2}} &\lesssim \sup_{x \in \mathbb{R}^n, r > 0} \varphi_2(x, r)^{-1} \int_r^{\infty} \|f\|_{L_1(B(x_0,t))} |B(x_0, t)|^{-1} \frac{dt}{t} \\ &\lesssim \sup_{x \in \mathbb{R}^n, r > 0} \varphi_1(x, r)^{-1} |B|^{-1} \|f\|_{L_1(B)} = \|f\|_{M_{1,\varphi_1}}. \end{aligned}$$

**Remark 3.1** Let  $0 \leq \kappa < 1$ . Assume that  $\psi$  is a positive increasing function defined in  $(0, \infty)$  and satisfies the following  $\mathcal{D}_\kappa$  condition :

$$\frac{\psi(t_2)}{t_2^\kappa} \leq C \frac{\psi(t_1)}{t_1^\kappa}, \text{ for any } 0 < t_1 < t_2 < \infty,$$

where  $C > 0$  is a constant independent of  $t_1$  and  $t_2$ . If  $\varphi_1(x, r) = \varphi_2(x, r) = \psi(|B(x, r)|)$  and  $\psi$  satisfy the  $\mathcal{D}_\kappa$  condition, Theorems 3.2 and 3.3 were proved in [29]. Also, in the case  $\omega(t) = t^\varepsilon$  with  $0 < \varepsilon \leq 1$ , Theorems 3.2 and 3.3 were proved in [10].

#### 4 Commutators of $\omega$ -type Calderón-Zygmund operators in the spaces $M_{p,\varphi}(\mathbb{R}^n)$

We recall the definition of the space of  $BMO(\mathbb{R}^n)$ .

**Definition 4.1** *Suppose that  $b \in L_1^{\text{loc}}(\mathbb{R}^n)$ , and let*

$$\|b\|_* = \sup_{x \in \mathbb{R}^n, r > 0} \frac{1}{|B(x, r)|} \int_{B(x, r)} |b(y) - b_{B(x, r)}| dy < \infty,$$

where

$$b_{B(x, r)} = \frac{1}{|B(x, r)|} \int_{B(x, r)} b(y) dy.$$

Define

$$BMO(\mathbb{R}^n) = \{b \in L_1^{\text{loc}}(\mathbb{R}^n) : \|b\|_* < \infty\}.$$

Modulo constants, the space  $BMO(\mathbb{R}^n)$  is a Banach space with respect to the norm  $\|\cdot\|_*$ . The following lemma is valid.

**Lemma 4.1** [19, 28] (1) *Let  $b \in BMO(\mathbb{R}^n)$ . Then*

$$\|b\|_* \approx \sup_{x \in \mathbb{R}^n, r > 0} \left( \frac{1}{|B(x, r)|} \int_{B(x, r)} |b(y) - b_{B(x, r)}|^p dy \right)^{\frac{1}{p}} \quad (4.1)$$

for  $1 < p < \infty$ .

(2) *Let  $b \in BMO(\mathbb{R}^n)$ . Then there is a constant  $C > 0$  such that*

$$|b_{B(x, r)} - b_{B(x, \tau)}| \leq C \|b\|_* \log \frac{\tau}{r} \text{ for } 0 < 2r < \tau, \quad (4.2)$$

where  $C$  is independent of  $f$ ,  $x$ ,  $r$  and  $\tau$ .

Since linear commutator has a greater degree of singularity than the corresponding  $\omega$ -type Calderón-Zygmund operator, we need a slightly stronger version of condition

$$\int_0^1 \frac{\omega(t)}{t} \left(1 + \log \frac{1}{t}\right)^m dt < \infty. \quad (4.3)$$

The following weighted endpoint estimate for commutator  $T_{\mathbf{b}}$  of the  $\omega$ -type Calderón-Zygmund operator was established in [30] under a stronger version of condition (4.3) assumed on  $\omega$ , if  $\mathbf{b} \in BMO^m(\mathbb{R}^n)$  (for the unweighted case, see [22]).

The following theorem was proved in [30].

**Theorem 4.1** [30] *Let  $T$  be linear  $\omega$ -CZO and  $\mathbf{b} \in BMO^m(\mathbb{R}^n)$ . If  $\omega$  satisfies condition (4.3) and  $1 < p < \infty$ , then there exists a constant  $C > 0$  such that*

$$\|T_{\mathbf{b}}f\|_{L_p} \leq C \|\mathbf{b}\|_* \|f\|_{L_p}.$$

The following Guliyev local estimates are valid (see [10]).

**Theorem 4.2** *Let  $T$  be linear  $\omega$ -CZO and  $\mathbf{b} \in BMO^m(\mathbb{R}^n)$ . Let also  $\omega$  satisfies condition (4.3) and  $1 < p < \infty$ . Then*

$$\|T_{\mathbf{b}}f\|_{L_p(B)} \leq C \|\mathbf{b}\|_* |B|^{\frac{1}{p}} \int_{2r}^{\infty} \ln^m \left(e + \frac{t}{r}\right) \|f\|_{L_p(B(x_0, t))} |B(x_0, t)|^{-1/p} \frac{dt}{t}$$

holds for any ball  $B = B(x_0, r)$  and for all  $f \in L_p^{\text{loc}}(\mathbb{R}^n)$ , where  $C$  does not depend on  $f$ ,  $x_0 \in \mathbb{R}^n$  and  $r > 0$ .

**Proof.** Let  $p \in (1, \infty)$ . For arbitrary  $x_0 \in \mathbb{R}^n$  and  $r > 0$ , set  $B = B(x_0, r)$ . Write  $f = f_1 + f_2$  with  $f_1 = f\chi_{2B}$  and  $f_2 = f\chi_{\mathbb{R}^n \setminus 2B}$ . For all  $f \in L_p^{\text{loc}}(\mathbb{R}^n)$  we define

$$T_{\mathbf{b}}f(x) := T_{\mathbf{b}}f_1(x) + \int_{\mathbb{R}^n} \prod_{j=1}^m (b_j(x) - b_j(y)) K(x, y) f_2(y) dy, \quad (4.4)$$

here  $T_{\mathbf{b}}$  denotes the commutator as a bounded linear operator on  $L_p$  with  $1 \leq p < \infty$  and  $w \in A_p(\mathbb{R}^n)$  (see [30]). It is easy to check that the definition of  $T_{\mathbf{b}}f(x)$  does not depend on the choice of the ball  $B$ .

Hence

$$\|T_{\mathbf{b}}f\|_{L_p(B)} \leq \|T_{\mathbf{b}}f_1\|_{L_p(B)} + \|T_{\mathbf{b}}f_2\|_{L_p(B)}.$$

From the boundedness of  $T_{\mathbf{b}}$  in  $L_p(\mathbb{R}^n)$  ( see Theorem 4.1) it follows that:

$$\|T_{\mathbf{b}}f_1\|_{L_p(B)} \leq \|T_{\mathbf{b}}f_1\|_{L_p} \lesssim \|\mathbf{b}\|_* \|f_1\|_{L_p} = \|\mathbf{b}\|_* \|f\|_{L_p(2B)}.$$

For the term  $\|T_{\mathbf{b}}f_2\|_{L_p(B)}$ , without loss of generality, we can assume  $m = 2$ . Thus, the operator  $T_{\mathbf{b}}f_2$  can be divided into four parts

$$\begin{aligned} T_{\mathbf{b}}f_2(x) &= (b_1(x) - (b_1)_B)(b_2(x) - (b_2)_B) \int_{\mathbb{R}^n} K(x, y)f_2(y)dy \\ &+ \int_{\mathbb{R}^n} K(x, y)(b_1(y) - (b_1)_B)(b_2(y) - (b_2)_B)f_2(y)dy \\ &- (b_1(x) - (b_1)_B) \int_{\mathbb{R}^n} K(x, y)(b_2(y) - (b_2)_B)f_2(y)dy \\ &- (b_2(x) - (b_2)_B) \int_{\mathbb{R}^n} K(x, y)(b_1(y) - (b_1)_B)f_2(y)dy \\ &= I_1(x) + I_2(x) + I_3(x) + I_4(x). \end{aligned}$$

For  $x \in B$  we have

$$\begin{aligned} |T_{\mathbf{b}}f_2(x)| &\leq |I_1(x) + I_2(x)| + |I_3(x)| + |I_4(x)| \\ &\lesssim |b_1(x) - (b_1)_B| |b_2(x) - (b_2)_B| \int_{\mathfrak{c}_{(2B)}} \frac{|f(y)|}{|x_0 - y|^n} dy \\ &+ \int_{\mathfrak{c}_{(2B)}} |b_1(y) - (b_1)_B| |b_2(y) - (b_2)_B| \frac{|f(y)|}{|x_0 - y|^n} dy \\ &+ |b_1(x) - (b_1)_B| \int_{\mathfrak{c}_{(2B)}} |b_2(y) - (b_2)_B| \frac{|f(y)|}{|x_0 - y|^n} dy \\ &+ |b_2(x) - (b_2)_B| \int_{\mathfrak{c}_{(2B)}} |b_1(y) - (b_1)_B| \frac{|f(y)|}{|x_0 - y|^n} dy. \end{aligned}$$

Then

$$\begin{aligned} \|T_{\mathbf{b}}f_2\|_{L_p(B)} &\lesssim \left( \int_B \left( \int_{\mathfrak{c}_{(2B)}} \frac{\prod_{j=1}^2 |b_j(y) - (b_j)_B|}{|x_0 - y|^n} |f(y)| dy \right)^p dx \right)^{\frac{1}{p}} \\ &+ \left( \int_B |b_1(x) - (b_1)_B| \left( \int_{\mathfrak{c}_{(2B)}} \frac{|b_2(y) - (b_2)_B|}{|x_0 - y|^n} |f(y)| dy \right)^p dx \right)^{\frac{1}{p}} \\ &+ \left( \int_B |b_2(x) - (b_2)_B| \left( \int_{\mathfrak{c}_{(2B)}} \frac{|b_1(y) - (b_1)_B|}{|x_0 - y|^n} |f(y)| dy \right)^p dx \right)^{\frac{1}{p}} \\ &+ \left( \int_B \left( \int_{\mathfrak{c}_{(2B)}} \frac{\prod_{j=1}^2 |b_j(x) - (b_j)_B|}{|x_0 - y|^n} |f(y)| dy \right)^p dx \right)^{\frac{1}{p}} \\ &= I_1 + I_2 + I_3 + I_4. \end{aligned}$$

Let us estimate  $I_1$ .

$$\begin{aligned}
I_1 &= |B|^{\frac{1}{p}} \int_{\mathfrak{C}_{(2B)}} \frac{\prod_{j=1}^2 |b_j(y) - (b_j)_B|}{|x_0 - y|^n} |f(y)| dy \\
&\approx |B|^{\frac{1}{p}} \int_{\mathfrak{C}_{(2B)}} \prod_{j=1}^2 |b_j(y) - (b_j)_B| |f(y)| \int_{|x_0-y|}^{\infty} \frac{dt}{t^{n+1}} dy \\
&\approx |B|^{\frac{1}{p}} \int_{2r}^{\infty} \int_{2r \leq |x_0-y| \leq t} \prod_{j=1}^2 |b_j(y) - (b_j)_B| |f(y)| dy \frac{dt}{t^{n+1}} \\
&\lesssim |B|^{\frac{1}{p}} \int_{2r}^{\infty} \int_{B(x_0,t)} \prod_{j=1}^2 |b_j(y) - (b_j)_B| |f(y)| dy \frac{dt}{t^{n+1}}.
\end{aligned}$$

Applying Hölder's inequality and by Lemma 4.1, we get

$$\begin{aligned}
I_1 &\lesssim |B|^{\frac{1}{p}} \int_{2r}^{\infty} \prod_{j=1}^2 \left( \int_{B(x_0,t)} |b_j(y) - (b_j)_B|^{2p'} w(y)^{1-2p'} dy \right)^{\frac{1}{2p'}} \|f\|_{L_p(B(x_0,t))} \frac{dt}{t^{n+1}} \\
&\lesssim \prod_{j=1}^2 \|b_j\|_* |B|^{\frac{1}{p}} \int_{2r}^{\infty} \left(1 + \ln \frac{t}{r}\right)^2 \|1\|_{L_{p'}(B(x_0,t))} \|f\|_{L_p(B(x_0,t))} \frac{dt}{t^{n+1}} \\
&\lesssim \|\mathbf{b}\|_* |B|^{\frac{1}{p}} \int_{2r}^{\infty} \ln^2 \left(e + \frac{t}{r}\right) \|f\|_{L_p(B(x_0,t))} |B(x_0,t)|^{-1/p} \frac{dt}{t}.
\end{aligned}$$

Let us estimate  $I_2$ .

$$\begin{aligned}
I_2 &= \left( \int_B |b_1(x) - (b_1)_B|^p dx \right)^{\frac{1}{p}} \int_{\mathfrak{C}_{(2B)}} \frac{|b_2(y) - (b_2)_B|}{|x_0 - y|^n} |f(y)| dy \\
&\lesssim \|b_1\|_* |B|^{\frac{1}{p}} \int_{\mathfrak{C}_{(2B)}} |b_2(y) - (b_2)_B| |f(y)| \int_{|x_0-y|}^{\infty} \frac{dt}{t^{n+1}} dy \\
&\approx \|b_1\|_* |B|^{\frac{1}{p}} \int_{2r}^{\infty} \int_{2r \leq |x_0-y| \leq t} |b_2(y) - (b_2)_B| |f(y)| dy \frac{dt}{t^{n+1}} \\
&\lesssim \|b_1\|_* |B|^{\frac{1}{p}} \int_{2r}^{\infty} \int_{B(x_0,t)} |b_2(y) - (b_2)_B| |f(y)| dy \frac{dt}{t^{n+1}}.
\end{aligned}$$

Applying Hölder's inequality and by Lemma 4.1, we get

$$\begin{aligned}
I_2 &\lesssim \|b_1\|_* |B|^{\frac{1}{p}} \int_{2r}^{\infty} \left( \int_{B(x_0,t)} |b_2(y) - (b_2)_B|^{p'} dy \right)^{\frac{1}{p'}} \|f\|_{L_p(B(x_0,t))} \frac{dt}{t^{n+1}} \\
&\lesssim \prod_{j=1}^2 \|b_j\|_* |B|^{\frac{1}{p}} \int_{2r}^{\infty} \left(1 + \ln \frac{t}{r}\right) \|1\|_{L_{p'}(B(x_0,t))} \|f\|_{L_p(B(x_0,t))} \frac{dt}{t^{n+1}} \\
&\lesssim \|\mathbf{b}\|_* |B|^{\frac{1}{p}} \int_{2r}^{\infty} \ln^2 \left(e + \frac{t}{r}\right) \|f\|_{L_p(B(x_0,t))} |B(x_0,t)|^{-1/p} \frac{dt}{t}.
\end{aligned}$$



In the same way, we shall get the result of  $I_3$

$$I_3 \lesssim \|\mathbf{b}\|_* |B|^{\frac{1}{p}} \int_{2r}^{\infty} \ln^2 \left( e + \frac{t}{r} \right) \|f\|_{L_p(B(x_0,t))} |B(x_0,t)|^{-1/p} \frac{dt}{t}.$$

In order to estimate  $I_4$  note that

$$\begin{aligned} I_4 &= \left( \int_B \prod_{j=1}^2 |b_j(x) - (b_j)_B|^p dx \right)^{\frac{1}{p}} \int_{\mathfrak{C}_{(2B)}} \frac{|f(y)|}{|x_0 - y|^n} dy \\ &\leq \prod_{j=1}^2 \left( \int_B |b_j(x) - (b_j)_B|^{2p} dx \right)^{\frac{1}{2p}} \int_{\mathfrak{C}_{(2B)}} \frac{|f(y)|}{|x_0 - y|^n} dy. \end{aligned}$$

By Lemma 4.1, we get

$$I_4 \lesssim \|\mathbf{b}\|_* |B|^{\frac{1}{p}} \int_{\mathfrak{C}_{(2B)}} \frac{|f(y)|}{|x_0 - y|^n} dy.$$

Applying Hölder's inequality, we get

$$\begin{aligned} \int_{\mathfrak{C}_{(2B)}} \frac{|f(y)|}{|x_0 - y|^n} dy &\lesssim \int_{2r}^{\infty} \|f\|_{L_p(B(x_0,t))} \|1\|_{L_{p'}(B(x_0,t))} \frac{dt}{t^{n+1}} \\ &\leq \int_{2r}^{\infty} \|f\|_{L_p(B(x_0,t))} |B(x_0,t)|^{-1/p} \frac{dt}{t}. \end{aligned} \quad (4.5)$$

Thus, by (4.5)

$$I_4 \lesssim \|\mathbf{b}\|_* |B|^{\frac{1}{p}} \int_{2r}^{\infty} \|f\|_{L_p(B(x_0,t))} |B(x_0,t)|^{-1/p} \frac{dt}{t}.$$

Summing up  $I_1$  and  $I_4$ , for all  $p \in [1, \infty)$  we get

$$\|T_{\mathbf{b}} f_2\|_{L_p(B)} \lesssim \|\mathbf{b}\|_* |B|^{\frac{1}{p}} \int_{2r}^{\infty} \ln^2 \left( e + \frac{t}{r} \right) \|f\|_{L_p(B(x_0,t))} |B(x_0,t)|^{-1/p} \frac{dt}{t}. \quad (4.6)$$

On the other hand,

$$\begin{aligned} \|f\|_{L_p(2B)} &\lesssim |B| \int_{2r}^{\infty} \|f\|_{L_p(B(x_0,t))} \frac{dt}{t^{n+1}} \\ &\leq |B|^{\frac{1}{p}} \|1\|_{L_{p'}(B)} \int_{2r}^{\infty} \|f\|_{L_p(B(x_0,t))} \frac{dt}{t^{n+1}} \\ &\leq |B|^{\frac{1}{p}} \int_{2r}^{\infty} \|f\|_{L_p(B(x_0,t))} \|1\|_{L_{p'}(B(x_0,t))} \frac{dt}{t^{n+1}} \\ &\leq |B|^{\frac{1}{p}} \int_{2r}^{\infty} \|f\|_{L_p(B(x_0,t))} |B(x_0,t)|^{-1/p} \frac{dt}{t}. \end{aligned} \quad (4.7)$$

Finally,

$$\begin{aligned} \|T_{\mathbf{b}} f\|_{L_p(B)} &\lesssim \|\mathbf{b}\|_* \|f\|_{L_p(2B)} \\ &\quad + \|\mathbf{b}\|_* |B|^{\frac{1}{p}} \int_{2r}^{\infty} \ln^m \left( e + \frac{t}{r} \right) \|f\|_{L_p(B(x_0,t))} |B(x_0,t)|^{-1/p} \frac{dt}{t}, \end{aligned}$$

and the statement of Theorem 4.2 follows by (4.7).

**Theorem 4.3** Let  $T$  be linear  $\omega$ -CZO and  $\mathbf{b} \in BMO^m(\mathbb{R}^n)$ . Let also  $\omega$  satisfies condition (4.3),  $1 < p < \infty$  and  $(\varphi_1, \varphi_2)$  satisfy the condition

$$\int_r^\infty \ln^m \left( e + \frac{t}{r} \right) \frac{\operatorname{ess\,inf}_{t < s < \infty} \varphi_1(x, s) |B(x, s)|^{1/p}}{|B(x, t)|^{1/p}} \frac{dt}{t} \leq C \varphi_2(x, r) \quad (4.8)$$

where  $C$  does not depend on  $x$  and  $r$ . Then the operator  $T_{\mathbf{b}}$  is bounded from  $M_{p, \varphi_1}$  to  $M_{p, \varphi_2}$ . Moreover,

$$\|T_{\mathbf{b}}f\|_{M_{p, \varphi_2}} \lesssim \|\mathbf{b}\|_* \|f\|_{M_{p, \varphi_1}}.$$

**Proof.** Using the Theorem 2.2 and the Theorem 4.2 we have

$$\begin{aligned} \|T_{\mathbf{b}}f\|_{M_{p, \varphi_2}} &= \sup_{x \in \mathbb{R}^n, r > 0} \varphi_2(x, r)^{-1} |B(x, t)|^{-\frac{1}{p}} \|T_{\mathbf{b}}f\|_{L_p(B(x, r))} \\ &\lesssim \|\mathbf{b}\|_* \sup_{x \in \mathbb{R}^n, r > 0} \varphi_2(x, r)^{-1} \int_r^\infty \ln^m \left( e + \frac{t}{r} \right) \|f\|_{L_p(B(x, t))} |B(x, t)|^{-1/p} \frac{dt}{t} \\ &\lesssim \|\mathbf{b}\|_* \sup_{x \in \mathbb{R}^n, r > 0} \varphi_1(x, r)^{-1} |B(x, t)|^{-\frac{1}{p}} \|f\|_{L_p(B(x, r))} = \|\mathbf{b}\|_* \|f\|_{M_{p, \varphi_1}}. \end{aligned}$$

**Remark 4.1** Note that, if  $\varphi_1(x, r) = \varphi_2(x, r) = \psi(w(x, r))$  and  $\psi$  satisfy the  $\mathcal{D}_\kappa$  condition, Theorems 4.2 and 4.3 were proved in [29]. Also, in the case  $m = 1$  and  $\omega(t) = t^\varepsilon$  with  $0 < \varepsilon \leq 1$ , Theorems 4.2 and 4.3 were proved in [11].

**Acknowledgements.** The author would like to express their gratitude to the referees for his very valuable comments and suggestions. The author would like to express his deep gratitude to corresponding member of the National Academy of Sciences of Azerbaijan, Professor Vagif S. Guliyev for his attention to this work.

## References

1. Akbulut, A., Burenkov, V.I., Guliyev, V.S.: *Anisotropic fractional maximal commutators with BMO functions on anisotropic morrey-type spaces*, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci. **40**(4), Mathematics, 13-32 (2020).
2. Calderon, A.P.: *Commutators of singular integral operators*, Proc. Natl. Acad. Sci. USA, **53**, 1092-1099 (1965).
3. Calderon, A.P.: *Cauchy integrals on Lipschitz curves and related operators*, Proc. Natl. Acad. Sci. USA, **74**(4), 1324-1327 (1977).
4. Coifman, R.R., Meyer, Y.: *Au delà des Opérateurs Pseudo-Différentiels*, *Astérisque 57. Société Maématique de France, Paris, France*, (1978).
5. Eroglu, A., Azizov, J.V.: *A note on the fractional integral operators in generalized Morrey spaces on the Heisenberg group*, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci. **37**(1), Mathematics, 86-91 (2017).
6. Eroglu, A., Isayev, F.A., Namazov, F.M.: *Fractional integral associated to Schrödinger operator on the Heisenberg groups in vanishing generalized Morrey spaces*, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci., **38**(4), Mathematics, 54-62 (2018).
7. Giaquinta, M.: *Multiple integrals in the calculus of variations and nonlinear elliptic systems*, *Princeton Univ. Press, Princeton, NJ*, 1983.
8. Guliyev, V.S.: *Integral operators on function spaces on the homogeneous groups and on domains in  $\mathbb{R}^n$* , *Doctoral dissertation, Moscow, Mat. Inst. Steklov*, 329 pp. (1994) (in Russian).

9. Guliyev, V.S.: Function spaces, integral operators and two weighted inequalities on homogeneous groups. Some applications, *Baku, Elm.* 332 pp. (1999) (Russian).
10. Guliyev, V.S.: *Boundedness of the maximal, potential and singular operators in the generalized Morrey spaces*, J. Inequal. Appl. Art. ID 503948, 20 pp. (2009).
11. Guliyev, V.S.: *Generalized weighted Morrey spaces and higher order commutators of sublinear operators*, Eurasian Math. J. **3**(3), 33-61 (2012).
12. Guliyev, V.S.: *Generalized local Morrey spaces and fractional integral operators with rough kernel*, J. Math. Sci., (N. Y.) **193**(2), 211-227 (2013).
13. Guliyev, V.S., Ismayilova, A.F.: *Multi-sublinear maximal operator and multilinear singular integral operators on generalized Morrey spaces*, Proc. Inst. Math. Mech. Natl. Acad. Sci. Azerb. **40**(2), 65-77 (2014).
14. Guliyev, V.S., Guliyev, R.V., Omarova, M.N.: *Riesz transforms associated with Schrödinger operator on vanishing generalized Morrey spaces*, Appl. Comput. Math. **17**(1), 56-71 (2018).
15. Guliyev, V.S., Akbulut, A., Celik, S., Omarova, M.N.: *Higher order Riesz transforms related to Schrödinger type operator on local generalized Morrey spaces*, TWMS Jour. Pure Appl. Math. **10**(1), 58-75 (2019).
16. Hasanov, A.A.: *(p, p)-admissible multi-sublinear singular integral operators in product generalized local Morrey spaces*, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci., **40**(4), Mathematics, 85-96 (2020).
17. Ibrahimov, E.J., Dadashova, G.A., Ekincioglu, S.E.: *On the boundedness of the G-maximal operator and G-Riesz potential in the generalized G-Morrey spaces*, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci., **40**(1), Mathematics, 111-125 (2020).
18. Ismayilova, A.F., Isayev, F.A.: *Multi-sublinear operators generated by multilinear Calderón-Zygmund operators on product generalized Morrey spaces*, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci., **40**(4), Mathematics, 110-117 (2020).
19. Kufner, A., John, O., Fučík, S.: *Function Spaces*, Noordhoff International Publishing: Leyden, Publishing House Czechoslovak Academy of Sciences: Prague, (1977).
20. Lin, Y.: *Strongly singular Calderón-Zygmund operator and commutator on Morrey type spaces*, Acta Math. Sin. (Engl. Ser.) **23**(11), 2097-2110 (2007).
21. Lin, Y., Lu, Sh.: *Strongly singular Calderón-Zygmund operators and their commutators*, Jordan Journal of Mathematics and Statistics, **1**(1), 31-49 (2008).
22. Liu, Z.G., Lu, S.Z.: *Endpoint estimates for commutators of Calderón-Zygmund type operators*, Kodai Math. J., **25**(1), 79-88 (2002).
23. Morrey, C.B.: *On the solutions of quasi-linear elliptic partial differential equations*, Trans. Amer. Math. Soc., **43**, 126-166 (1938).
24. Mizuhara, T.: *Boundedness of some classical operators on generalized Morrey spaces*, Harmonic Analysis (Sendai, 1990), S. Igari, Ed. ICM 90 Satellite Conference Proceedings, Springer, Tokyo, Japan, pp. 183-189 (1991).
25. Nakai, E.: *Hardy-Littlewood maximal operator, singular integral operators and Riesz potentials on generalized Morrey spaces*, Math. Nachr. **166**, 95-103 (1994).
26. Quek, T., Yang, D.: *Calderón-Zygmund-type operators on weighted weak Hardy spaces over  $\mathbb{R}^n$* , Acta Math. Sin. (Engl. Ser.), **16**(1), 141-160 (2000).
27. Sawano, Y.: *A thought on generalized Morrey spaces*, J. Indonesian Math. Soc., **25**(3), 210-281 (2019).
28. Stein, E.M.: *Harmonic Analysis: Real Variable Methods, Orthogonality and Oscillatory Integrals*, vol. 43 of Princeton Mathematical Series, Princeton University Press, Princeton, NJ, USA, (1993).
29. Wang, H.: *Boundedness of  $\theta$ -type Calderón-Zygmund operators and commutators in the generalized weighted Morrey spaces*, J. Funct. Spaces, Art. ID 1309348, 18 pp. (2016).

30. Zhang, P., Sun J.: *Commutators of multilinear Calderón-Zygmund operators with kernels of Dini's type and applications*, J. Math. Inequal. **13**(4), 1071-1093 (2019).
31. Yabuta, K.: *Generalizations of Calderón-Zygmund operators*, Studia Math. **82**, 17-31 (1985).