

Multilinear Calderón-Zygmund operators with kernels of Dini's type and their commutators on generalized local Morrey spaces

Vagif S. Guliyev*, Fatai A. Isayev, Ayhan Serbetci

Received: 22.05.2022 / Revised: 21.09.2022 / Accepted: 02.11.2022

Abstract. Let T be a multilinear Calderón-Zygmund operator of type ω with $\omega(t)$ being nondecreasing and satisfying a kind of Dini's type condition and $T_{\Pi\mathbf{b}}$ be the iterated commutators of the operator T with BMO^m functions. In this paper, we study the boundedness of the operators T and $T_{\Pi\mathbf{b}}$ on generalized local Morrey spaces $M_{p,\varphi}^{\{x_0\}}(\mathbb{R}^n)$ and generalized Morrey spaces $M_{p,\varphi}(\mathbb{R}^n)$. We find the sufficient conditions on the pair (φ_1, φ_2) with $\mathbf{b} \in BMO^m$ which ensures the boundedness of the operators T and $T_{\Pi\mathbf{b}}$ from $M_{p_1,\varphi_{11}}^{\{x_0\}}(\mathbb{R}^n) \times \dots \times M_{p_m,\varphi_{1m}}^{\{x_0\}}(\mathbb{R}^n)$ to $M_{p,\varphi_2}^{\{x_0\}}(\mathbb{R}^n)$.

Keywords. Multilinear Calderón-Zygmund operator, generalized local Morrey spaces, commutator, BMO

Mathematics Subject Classification (2010): 42B20, 42B25, 47G30, 35S05, 46E30

1 Introduction and Main Results

Let T be a multilinear Calderon-Zygmund operator such that

$$T(f_1, \dots, f_m)(x) = \int_{(\mathbb{R}^n)^m} K(x, y_1, \dots, y_m) f_1(y_1) \dots f_m(y_m) dy_1 \dots dy_m,$$

$x \notin \bigcap_{j=1}^m \text{supp} f_j$, where $K(x, y_1, \dots, y_m)$ is an m -linear Calderon-Zygmund kernel of type $\omega(t)$ and each $f_j \in C_c^\infty(\mathbb{R}^n)$, $j = 1, \dots, m$. Let $\mathbf{b} = (b_1, \dots, b_m) \in BMO^m$ such that

* Corresponding author

V.S. Guliyev
Institute of Applied Mathematics, Baku State University, Baku, Azerbaijan
Institute of Mathematics and Mechanics of NAS of Azerbaijan, Baku, Azerbaijan
E-mail: vagif@guliyev.com

F.A. Isayev
Institute of Mathematics and Mechanics of NAS of Azerbaijan, Baku, Azerbaijan
E-mail: isayevfatai@yahoo.com

A. Serbetci
Department of Mathematics, Ankara University, Ankara, Turkey
E-mail: serbetci@ankara.edu.tr

$b_j \in BMO(\mathbb{R}^n)$, $j = 1, \dots, m$. The m -linear commutator of T with \mathbf{b} is defined by

$$T_{\Sigma\mathbf{b}}(f_1, \dots, f_m) = \sum_{j=1}^m T_{b_j}^j(\mathbf{f}),$$

where

$$T_{b_j}^j(\mathbf{f}) = b_j T(f_1, \dots, f_j, \dots, f_m) - T(f_1, \dots, f_j, \dots, f_m). \quad (1.1)$$

The iterated commutator of T with \mathbf{b} is defined by

$$T_{\Pi\mathbf{b}}(\mathbf{f})(x) = [b_1, [b_2, \dots [b_{m-1}, [b_m, T]_m]_{m-1} \dots]_2,]_1(\mathbf{f})(x). \quad (1.2)$$

For an m -linear Calderón-Zygmund operator with associated kernel $K(x, \mathbf{y})$, the iterated commutator $T_{\Pi\mathbf{b}}$ is given formally by

$$T_{\Pi\mathbf{b}}(\mathbf{f})(x) = \int_{(\mathbb{R}^n)^m} \left(\prod_{j=1}^m (b_j(x) - b_j(y_j)) \right) K(x, \mathbf{y}) f_1(y_1) \dots f_m(y_m) d\mathbf{y}.$$

Here and in what follows, $\mathbf{y} = (y_1, \dots, y_m)$, $(x, \mathbf{y}) = (x, y_1, \dots, y_m)$ and $d\mathbf{y} = dy_1 \dots dy_m$.

The study of multilinear Calderón-Zygmund theory goes back to the pioneering works of Coifman and Meyer in 1970s, see e.g. [2, 3]. This topic was then further investigated by many authors in the last few decades, see for example [6, 15, 18, 23]. The classical Morrey spaces $M_{p,\lambda}$ were introduced by Morrey [21] to study the local behavior of solutions to second-order elliptic partial differential equations. Moreover, various Morrey spaces are defined in the process of study. The first author, Mizuhara and Nakai [7, 20, 22] introduced generalized Morrey spaces $M_{p,\varphi}(\mathbb{R}^n)$ (see, also [8, 12, 24]). In [4], the boundedness of the multilinear commutator of Calderón-Zygmund operator $T_{\mathbf{b}}$ from one generalized variable exponent Morrey space $M_{p(\cdot),\varphi_1}$ to another $M_{p(\cdot),\varphi_2}$ for $\mathbf{b} = (b_1, \dots, b_m) \in BMO^m$ was proved, see also [1, 5, 13, 14].

The purpose of this paper is to prove the boundedness of multilinear Calderón-Zygmund operators T of type ω and their iterated commutators $T_{\Pi\mathbf{b}}$ with BMO^m functions from $M_{p_1,\varphi_{11}}^{\{x_0\}}(\mathbb{R}^n) \times \dots \times M_{p_m,\varphi_{1m}}^{\{x_0\}}(\mathbb{R}^n)$ to $M_{p,\varphi_2}^{\{x_0\}}(\mathbb{R}^n)$.

Our main results formulated as follows:

Theorem 1.1 *Let $m \geq 2$, $x_0 \in \mathbb{R}^n$, T be an m -linear ω - CZO and $\omega \in \text{Dini}(1)$. Suppose that $p_1, \dots, p_m \in [1, \infty)$, $p \in (0, \infty)$ with $1/p = \sum_{i=1}^m 1/p_i$, $\varphi_1 = (\varphi_{11}, \dots, \varphi_{1m})$ and (φ_1, φ_2) satisfies the condition*

$$\int_r^\infty t^{-\frac{n}{p}} \operatorname{ess\,inf}_{t < s < \infty} s^{\frac{n}{p}} \prod_{i=1}^m \varphi_{1i}(x_0, s) \frac{dt}{t} \leq C \varphi_2(x_0, r), \quad (1.3)$$

where C does not depend on r .

(1) *If $1 < p_j < \infty$ for all $j = 1, \dots, m$, then there exists a constant $C > 0$ such that*

$$\|T(\mathbf{f})\|_{M_{p,\varphi_2}^{\{x_0\}}} \leq C \prod_{j=1}^m \|f_j\|_{M_{p_j,\varphi_{1j}}^{\{x_0\}}}.$$

(2) *If $1 \leq p_j < \infty$ for all $j = 1, \dots, m$, and at least one of the $p_j = 1$, then there exists a constant $C > 0$ such that*

$$\|T(\mathbf{f})\|_{WM_{p,\varphi_2}^{\{x_0\}}} \leq C \prod_{j=1}^m \|f_j\|_{M_{p_j,\varphi_{1j}}^{\{x_0\}}}.$$

Corollary 1.1 Let T be an m -linear ω -CZO, $m \geq 2$, and $\omega \in \text{Dini}(1)$. Suppose that $p_1, \dots, p_m \in [1, \infty)$, $p \in (0, \infty)$ with $1/p = \sum_{i=1}^m 1/p_i$, $\varphi_1 = (\varphi_{11}, \dots, \varphi_{1m})$ and (φ_1, φ_2) satisfies the condition

$$\int_r^\infty t^{-\frac{n}{p}} \operatorname{ess\,inf}_{t < s < \infty} s^{\frac{n}{p}} \prod_{i=1}^m \varphi_{1i}(x, s) \frac{dt}{t} \leq C \varphi_2(x, r), \quad (1.4)$$

where C does not depend on x and r .

(1) If $1 < p_j < \infty$ for all $j = 1, \dots, m$, then there exists a constant $C > 0$ such that

$$\|T(\mathbf{f})\|_{M_{p, \varphi_2}} \leq C \prod_{j=1}^m \|f_j\|_{M_{p_i, \varphi_{1i}}}.$$

(2) If $1 \leq p_j < \infty$ for all $j = 1, \dots, m$, and at least one of the $p_j = 1$, then there exists a constant $C > 0$ such that

$$\|T(\mathbf{f})\|_{WM_{p, \varphi_2}} \leq C \prod_{j=1}^m \|f_j\|_{M_{p_i, \varphi_{1i}}}.$$

Theorem 1.2 Let $m \geq 2$, $x_0 \in \mathbb{R}^n$, T be an m -linear ω -CZO and ω satisfies

$$\int_0^1 \frac{\omega(t)}{t} \left(1 + \log \frac{1}{t}\right)^m dt < \infty. \quad (1.5)$$

Suppose that $p_1, \dots, p_m \in (1, \infty)$, $p \in (0, \infty)$ with $1/p = \sum_{i=1}^m 1/p_i$, and (φ_1, φ_2) satisfies the condition

$$\int_r^\infty \left(1 + \ln \frac{t}{r}\right)^m t^{-\frac{n}{p}} \operatorname{ess\,inf}_{t < s < \infty} s^{\frac{n}{p}} \prod_{i=1}^m \varphi_{1i}(x_0, s) \frac{dt}{t} \leq C \varphi_2(x_0, r), \quad (1.6)$$

where C does not depend on r .

If $b_i \in BMO$, $\mathbf{b} = (b_1, \dots, b_m)$, then there exist constants $C > 0$ independent of $\mathbf{f} = (f_1, \dots, f_m)$ such that

$$\|T_{\Pi \mathbf{b}}(\mathbf{f})\|_{M_{p, \varphi_2}^{\{x_0\}}} \leq C \|\mathbf{b}\|_* \prod_{i=1}^m \|f_i\|_{M_{p_i, \varphi_{1i}}^{\{x_0\}}}, \quad (1.7)$$

where $\|\mathbf{b}\|_* = \prod_{i=1}^m \|b_i\|_*$.

Corollary 1.2 Let $m \geq 2$, T be an m -linear ω -CZO and ω satisfies (1.5). Suppose that $p_1, \dots, p_m \in (1, \infty)$, $p \in (0, \infty)$ with $1/p = \sum_{i=1}^m 1/p_i$, and (φ_1, φ_2) satisfies the condition

$$\int_r^\infty \left(1 + \ln \frac{t}{r}\right)^m t^{-\frac{n}{p}} \operatorname{ess\,inf}_{t < s < \infty} s^{\frac{n}{p}} \prod_{i=1}^m \varphi_{1i}(x, s) \frac{dt}{t} \leq C \varphi_2(x, r), \quad (1.8)$$

where C does not depend on x and r .

If $b_i \in BMO$, $\mathbf{b} = (b_1, \dots, b_m)$, then there exist constants $C > 0$ independent of $\mathbf{f} = (f_1, \dots, f_m)$ such that

$$\|T_{\Pi\mathbf{b}}(\mathbf{f})\|_{M_{p,\varphi_2}} \leq C \|\mathbf{b}\|_* \prod_{i=1}^m \|f_i\|_{M_{p_i,\varphi_{1i}}}. \quad (1.9)$$

Theorem 1.3 Let $m \geq 2$, $x_0 \in \mathbb{R}^n$, T be an m -linear ω -CZO and ω satisfies (1.5). Suppose that (φ_1, φ_2) satisfies the condition

$$\int_r^\infty \left(1 + \ln \frac{t}{r}\right)^m t^{-n} \operatorname{ess\,inf}_{t < s < \infty} s^n \prod_{i=1}^m \varphi_{1i}(x_0, s) \frac{dt}{t} \leq C \varphi_2(x_0, r), \quad (1.10)$$

where C does not depend on r .

If $b_i \in BMO$, $\mathbf{b} = (b_1, \dots, b_m)$, then there exist constants $C > 0$ independent of $\mathbf{f} = (f_1, \dots, f_m)$ such that

$$\|T_{\Pi\mathbf{b}}(\mathbf{f})\|_{WM_{1,\varphi_2}^{\{x_0\}}} \leq C \|\mathbf{b}\|_* \prod_{i=1}^m \|f_i\|_{M_{\Phi^{(m)},\varphi_{1i}}^{\{x_0\}}}, \quad (1.11)$$

where $\Phi(t) = t(1 + \log^+ t)$ and $\Phi^{(m)} = \underbrace{\Phi \circ \dots \circ \Phi}_{m \text{ times}}$.

Corollary 1.3 Let $m \geq 2$, T be an m -linear ω -CZO and ω satisfies (1.5). Suppose that (φ_1, φ_2) satisfies the condition

$$\int_r^\infty \left(1 + \ln \frac{t}{r}\right)^m t^{-n} \operatorname{ess\,inf}_{t < s < \infty} s^n \prod_{i=1}^m \varphi_{1i}(x, s) \frac{dt}{t} \leq C \varphi_2(x, r),$$

where C does not depend on x and r .

If $b_i \in BMO$, $\mathbf{b} = (b_1, \dots, b_m)$, then there exist constants $C > 0$ independent of $\mathbf{f} = (f_1, \dots, f_m)$ such that

$$\|T_{\Pi\mathbf{b}}(\mathbf{f})\|_{WM_{1,\varphi_2}} \leq C \|\mathbf{b}\|_* \prod_{i=1}^m \|f_i\|_{M_{\Phi^{(m)},\varphi_{1i}}},$$

This paper is arranged as follows. In Section 2, we recall some basic definitions and known results. Section 3 is devoted to proving our main theorems.

Throughout this article, we will use C to denote a positive constant, which is independent of the main parameters and not necessarily the same at each occurrence. Besides, we will denote the conjugate exponent of $p > 1$ by $p' = p/(p-1)$. By $A \lesssim B$, we mean that $A \leq CB$ for some constant $C > 0$, and $A \approx B$ means that $A \lesssim B$ and $B \lesssim A$.

2 Notations and definitions

We now recall the definition of multilinear Calderón-Zygmund operators of type ω .

Definition 2.1 Let $w(t) : [0, \infty) \rightarrow [0, \infty)$ be a nondecreasing function. A locally integrable function $K(x, y_1, \dots, y_m)$, defined away from the diagonal $x = y_1 = \dots = y_m$ in $(\mathbb{R}^n)^{m+1}$, is called an m -linear Calderón-Zygmund kernel of type ω if, for some constants $0 < \tau < 1$, there exists a constant $A > 0$ such that

$$|K(x, y_1, \dots, y_m)| \leq \frac{A}{(|x - y_1| + \dots + |x - y_m|)^{mn}}$$

for all $(x, y_1, \dots, y_m) \in (\mathbb{R}^n)^{m+1}$ with $x \neq y_j$ for some $j \in \{1, 2, \dots, m\}$, and

$$\begin{aligned} & |K(x, y_1, \dots, y_m) - K(x', y_1, \dots, y_m)| \\ & \leq \frac{A}{(|x - y_1| + \dots + |x - y_m|)^{mn}} \omega\left(\frac{|x - x'|}{|x - y_1| + \dots + |x - y_m|}\right), \end{aligned}$$

whenever $|x - x'| \leq \tau \max_{1 \leq j \leq m} |x - y_j|$, and

$$\begin{aligned} & |K(x, y_1, \dots, y_j, \dots, y_m) - K(x', y_1, \dots, y'_j, \dots, y_m)| \\ & \leq \frac{A}{(|x - y_1| + \dots + |x - y_m|)^{mn}} \omega\left(\frac{|y_j - y'_j|}{|x - y_1| + \dots + |x - y_m|}\right), \end{aligned}$$

whenever $|y_j - y'_j| \leq \tau \max_{1 \leq i \leq m} |x - y_i|$.

We say $T : \mathcal{S}(\mathbb{R}^n) \times \dots \times \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}'(\mathbb{R}^n)$ is an m -linear Calderón-Zygmund kernel of type ω , $K(x, y_1, \dots, y_m)$, if

$$T(f_1, \dots, f_m)(x) = \int_{(\mathbb{R}^n)^m} K(x, y_1, \dots, y_m) f_1(y_1) \dots f_m(y_m) dy_1 \dots dy_m,$$

whenever $x \notin \bigcap_{j=1}^m \text{supp} f_j$ and each $f_j \in C_c^\infty(\mathbb{R}^n)$, $j = 1, \dots, m$.

If T can be extended to a bounded multilinear operator from $L_{q_1}(\mathbb{R}^n) \times \dots \times L_{q_m}(\mathbb{R}^n)$ to $L_{q, \infty}(\mathbb{R}^n)$ for some $1 \leq q_1, \dots, q_m < \infty$, and $\frac{1}{q} = \frac{1}{q_1} + \dots + \frac{1}{q_m}$, then T is called an m -linear Calderón-Zygmund operator of type w , abbreviated to m -linear w -CZO.

When $\omega(t) = t^\varepsilon$ for some $\varepsilon > 0$, the m -linear ω -CZO is exactly the multilinear Calderón-Zygmund operator studied in [6] and [16]. The linear ω -CZO was studied by Yabuta [25].

Definition 2.2 Let $\omega(t) : [0, \infty) \rightarrow [0, \infty)$ be a nondecreasing function. For $a > 0$, we say that ω satisfies the Dini(a) condition and write $\omega \in \text{Dini}(a)$, if

$$|\omega|_{\text{Dini}(a)} := \int_0^1 \frac{\omega^a(t)}{t} dt < \infty.$$

We would like to note that Maldonado and Naibo [19] studied the bilinear ω -CZOs when ω is a nondecreasing, concave function and belongs to $\text{Dini}(1/2)$. Recently, Lu and Zhang [18] improve and extend their results by removing the hypothesis that ω is concave and reducing the condition $w \in \text{Dini}(1/2)$ to a weaker condition $w \in \text{Dini}(1)$.

Theorem 2.1 [18] Let $w \in \text{Dini}(1)$ and T be an m -linear operator with an m -linear Calderón-Zygmund kernel of type ω . Suppose that for some $1 \leq q_1, \dots, q_m \leq \infty$, and some $0 < q < \infty$ with $\frac{1}{q} = \frac{1}{q_1} + \dots + \frac{1}{q_m}$, T maps $L_{q_1}(\mathbb{R}^n) \times \dots \times L_{q_m}(\mathbb{R}^n)$ into $L_{q, \infty}(\mathbb{R}^n)$. Then T can be extended to a bounded operator from $L_1(\mathbb{R}^n) \times \dots \times L_1(\mathbb{R}^n)$ to $L_{\frac{1}{m}, \infty}(\mathbb{R}^n)$.

Remark 2.1 Perez et al. [23] proved the same results in Theorems 1.3 and 1.4 in case $\omega(t) = t^\varepsilon$ for some $\varepsilon > 0$. We also note that similar results for T_{Σ^b} were proved in [18] for commutators of the linear Calderón-Zygmund operator of type ω , see [17].

The following result holds.

Theorem 2.2 [18] *Let T be an m -linear w -CZO and $w \in \text{Dini}(1)$. Let also $p_1, \dots, p_m \in [1, \infty)$, $p \in (0, \infty)$ with $\frac{1}{p} = \frac{1}{p_1} + \dots + \frac{1}{p_m}$.*

(1) *If $1 < p_j < \infty$ for all $j = 1, \dots, m$, then there exists a constant $C > 0$ such that*

$$\|T(\mathbf{f})\|_{L_p} \leq C \prod_{j=1}^m \|f_j\|_{L_{p_j}}.$$

(2) *If $1 \leq p_j < \infty$ for all $j = 1, \dots, m$, and at least one of the $p_j = 1$, then there exists a constant $C > 0$ such that*

$$\|T(\mathbf{f})\|_{WL_p} \leq C \prod_{j=1}^m \|f_j\|_{L_{p_j}}.$$

In [8] and [10] the first author defined the generalized Morrey spaces $M_{p,\varphi}(\mathbb{R}^n)$, the generalized local Morrey spaces $M_{p,\varphi}^{\{x_0\}}(\mathbb{R}^n)$ and their weak versions as follows.

Definition 2.3 *Let $1 \leq p < \infty$ and φ be a positive measurable function on $\mathbb{R}^n \times (0, \infty)$. We denote by $M_{p,\varphi}^{\{x_0\}}(\mathbb{R}^n)$ and $M_{p,\varphi}(\mathbb{R}^n)$ the generalized local Morrey space, the generalized Morrey space respectively, the spaces of all functions $f \in L_p^{\text{loc}}(\mathbb{R}^n)$ with finite norms*

$$\|f\|_{M_{p,\varphi}^{\{x_0\}}} = \sup_{r>0} \varphi(x_0, r)^{-1} |B(x_0, r)|^{-\frac{1}{p}} \|f\|_{L_p(B(x_0, r))},$$

$$\|f\|_{M_{p,\varphi}} = \sup_{x \in \mathbb{R}^n, r>0} \varphi(x, r)^{-1} |B(x, r)|^{-\frac{1}{p}} \|f\|_{L_p(B(x, r))},$$

respectively.

Furthermore, by $WM_{p,\varphi}(\mathbb{R}^n)^{\{x_0\}}$ and $WM_{p,\varphi}(\mathbb{R}^n)$ we denote the weak generalized Morrey space and weak generalized Morrey space respectively of all functions $f \in WL_p^{\text{loc}}(\mathbb{R}^n)$ for which,

$$\|f\|_{WM_{p,\varphi}^{\{x_0\}}} = \sup_{r>0} \varphi(x_0, r)^{-1} |B(x_0, r)|^{-\frac{1}{p}} \|f\|_{WL_p(B(x_0, r))} < \infty,$$

$$\|f\|_{WM_{p,\varphi}} = \sup_{x \in \mathbb{R}^n, r>0} \varphi(x, r)^{-1} |B(x, r)|^{-\frac{1}{p}} \|f\|_{WL_p(B(x, r))} < \infty,$$

respectively.

Remark 2.2 (1) If $\varphi(x, r) = r^{\frac{\lambda-n}{p}}$ with $0 < \lambda < n$, then $M_{p,\varphi}^{\{x_0\}}(\mathbb{R}^n) = L_{p,\lambda}^{\{x_0\}}(\mathbb{R}^n)$ is the local Morrey space, $M_{p,\varphi}(\mathbb{R}^n) = L_{p,\lambda}(\mathbb{R}^n)$ is the classical Morrey space and $WM_{p,\varphi}^{\{x_0\}}(\mathbb{R}^n) = WL_{p,\lambda}^{\{x_0\}}(\mathbb{R}^n)$ is the weak local Morrey space, $WM_{p,\varphi}(\mathbb{R}^n) = WL_{p,\lambda}(\mathbb{R}^n)$ is the weak Morrey space.

(2) If $\varphi(x, r) \equiv |B(x, r)|^{-\frac{1}{p}}$, then $M_{p,\varphi}^{\{x_0\}}(\mathbb{R}^n) = M_{p,\varphi}(\mathbb{R}^n) = L_p(\mathbb{R}^n)$ is the Lebesgue space and $WM_{p,\varphi}^{\{x_0\}}(\mathbb{R}^n) = WM_{p,\varphi}(\mathbb{R}^n) = WL_p(\mathbb{R}^n)$ is the weak Lebesgue space.

Let us recall the definition and some properties of BMO . A locally integrable function b belongs to BMO if

$$\|b\|_* := \sup_{B \subset \mathbb{R}^n} \frac{1}{|B|} \int_B |b(x) - b_B| dx < \infty, \quad (2.1)$$

where $b_B = |B|^{-1} \int_B b(y) dy$.

Lemma 2.1 [9] Let $b \in BMO$, $1 \leq p < \infty$, and $r_1, r_2 > 0$. Then,

$$\left(\frac{1}{|B(x_0, r_1)|} \int_{B(x_0, r_1)} |b(y) - b_{B(x_0, r_2)}|^p dy \right)^{1/p} \leq C \|b\|_* \left(1 + \left| \ln \frac{r_1}{r_2} \right| \right), \quad (2.2)$$

where $C > 0$ is independent of f , x_0 , r_1 , and r_2 .

We will use the following statement on the boundedness of the weighted Hardy operator

$$H_w g(t) := \int_t^\infty g(s) w(s) ds, \quad H_w^* g(t) := \int_t^\infty \left(1 + \ln \frac{s}{t} \right)^m g(s) w(s) ds, \quad t > 0,$$

where w is a weight. The following theorem was proved in [9].

Theorem 2.3 [9] Let v_1, v_2 and w be weights on $(0, \infty)$ and $v_1(t)$ be bounded outside a neighborhood of the origin. The inequality

$$\sup_{t>0} v_2(t) H_w g(t) \leq C \sup_{t>0} v_1(t) g(t)$$

holds for some $C > 0$ for all non-negative and non-decreasing g on $(0, \infty)$ if and only if

$$B := \sup_{t>0} v_2(t) \int_t^\infty \frac{w(s) ds}{\sup_{s<\tau<\infty} v_1(\tau)} < \infty.$$

Theorem 2.4 [9] Let v_1, v_2 and w be weights on $(0, \infty)$ and $v_1(t)$ be bounded outside a neighborhood of the origin. The inequality

$$\sup_{t>0} v_2(t) H_w^* g(t) \leq C \sup_{t>0} v_1(t) g(t)$$

holds for some $C > 0$, for all non-negative and non-decreasing g on $(0, \infty)$ if and only if

$$B := \sup_{t>0} v_2(t) \int_t^\infty \left(1 + \ln \frac{s}{t} \right)^m \frac{w(s) ds}{\sup_{s<\tau<\infty} v_1(\tau)} < \infty.$$

3 Proof of Main Results

We first prove a theorem below, such that we prove Theorem 1.1 with the help of this theorem.

Theorem 3.1 Let $m \geq 2$, $x_0 \in \mathbb{R}^n$, T_m be an m -linear w -CZO, and $w \in Dini(1)$. If $p_1, \dots, p_m \in (1, \infty)$ and $p \in (0, \infty)$ with $1/p = \sum_{i=1}^m 1/p_i$, then the inequality

$$\|T_m(\vec{f})\|_{L_p(B(x_0, r))} \lesssim r^{\frac{n}{p}} \int_{2r}^\infty t^{-\frac{n}{p}} \prod_{i=1}^m \|f_i\|_{L_{p_i}(B(x_0, t))} \frac{dt}{t} \quad (3.1)$$

holds for any balls $B(x_0, r)$, and for all $\vec{f} \in L_{p_1}^{loc}(\mathbb{R}^n) \times \dots \times L_{p_m}^{loc}(\mathbb{R}^n)$.

If $p_1, \dots, p_m \in [1, \infty)$, $\min\{p_1, \dots, p_m\} = 1$, and $p \in (0, \infty)$ with $1/p = \sum_{i=1}^m 1/p_i$, then the inequality

$$\|T(\mathbf{f})\|_{W_{L_p, B(x_0, r)}} \lesssim r^{\frac{n}{p}} \int_{2r}^\infty t^{-\frac{n}{p}} \prod_{i=1}^m \|f_i\|_{L_{p_i}(B(x_0, t))} \frac{dt}{t} \quad (3.2)$$

holds for any balls $B(x_0, r)$ and for all $\vec{f} \in L_{p_1}^{loc}(\mathbb{R}^n) \times \dots \times L_{p_m}^{loc}(\mathbb{R}^n)$.

Proof. Let $(p_1, \dots, p_m) \in (1, \infty)^m$ and $p \in (0, \infty)$ with $1/p = \sum_{i=1}^m 1/p_i$. For the moment, we denote the multilinear singular integral operator on $L_{p_1}(\mathbb{R}^n) \times \dots \times L_{p_m}(\mathbb{R}^n)$ by T_0 to avoid confusion.

For arbitrary $x \in \mathbb{R}^n$, set $B = B(x_0, r)$ for the ball centered at x with a radius r , $2B = B(x_0, 2r)$. We represent $\vec{f} = (f_1, \dots, f_m)$ as

$$f_j = f_j^0 + f_j^\infty, \quad f_j^0 = f_j \chi_{2B}, \quad f_j^\infty = f_j \chi_{\mathbb{C}(2B)}, \quad j = 1, \dots, m. \quad (3.3)$$

Then we write

$$\begin{aligned} \prod_{i=1}^m f_i(y_i) &= \prod_{i=1}^m (f_i^0(y_i) + f_i^\infty(y_i)) = \sum_{\beta_1, \dots, \beta_m \in \{0, \infty\}} f_1^{\beta_1}(y_1) \dots f_m^{\beta_m}(y_m) \\ &= \prod_{i=1}^m f_i^0(y_i) + \sum'_{\beta_1, \dots, \beta_m} f_1^{\beta_1}(y_1) \dots f_m^{\beta_m}(y_m), \end{aligned}$$

where each term in \sum' contains at least one $\beta_i \neq 0$.

For all $\vec{f} \in L_{p_1}^{loc}(\mathbb{R}^n) \times \dots \times L_{p_m}^{loc}(\mathbb{R}^n)$, we define

$$T(\vec{f})(x) := T_0(f_1^0, \dots, f_m^0)(x) + \sum'_{\beta_1, \dots, \beta_m} T(f_1^{\beta_1}, \dots, f_m^{\beta_m})(x), \quad (3.4)$$

where $\beta_1, \dots, \beta_m \in \{0, \infty\}$ and each term in \sum' contains at least one $\beta_i \neq 0$.

First we show that $T(\vec{f})(x)$ is well-defined *a.e.* x and independent of the choice B containing x . As T_0 is bounded on $L_{p_1}(\mathbb{R}^n) \times \dots \times L_{p_m}(\mathbb{R}^n)$ provided by Theorem 2.1 and $(f_1^0, \dots, f_m^0) \in L_{p_1}(\mathbb{R}^n) \times \dots \times L_{p_m}(\mathbb{R}^n)$, $T_0(f_1^0, \dots, f_m^0)$ is well-defined. Next, we show that the second-term of the right-hand side defining $T(\vec{f})(x)$ converges absolutely for any $\vec{f} \in L_{p_1}^{loc}(\mathbb{R}^n) \times \dots \times L_{p_m}^{loc}(\mathbb{R}^n)$ and almost every $x \in \mathbb{R}^n$.

For the other terms, let us first deal with the case when $\beta_1 = \dots = \beta_m = \infty$.

When $x \in B$, $y_i \in \mathbb{C}(2B)$, we have $\frac{1}{2}|x_0 - y_i| \leq |x - y_i| \leq \frac{3}{2}|x_0 - y_i|$, and therefore,

$$\begin{aligned} \sum'_{\beta_1, \dots, \beta_m} \left| T(f_1^{\beta_1}, \dots, f_m^{\beta_m})(x) \right| &\lesssim \int_{(\mathbb{C}(2B))^m} \frac{|f_1(y_1) \dots f_m(y_m)|}{|(x_0 - y_1, \dots, x_0 - y_m)|^{mn}} d\vec{y} \\ &\lesssim \int_{(\mathbb{C}(2B))^m} \prod_{i=1}^m \frac{|f_i(y_i)|}{|x_0 - y_i|^n} dy_i. \end{aligned}$$

By Fubini's theorem we have

$$\begin{aligned} \int_{(\mathbb{C}(2B))^m} \prod_{i=1}^m \frac{|f_i(y_i)|}{|x_0 - y_i|^{mn}} dy_i &\approx \int_{(\mathbb{C}(2B))^m} \prod_{i=1}^m |f_i(y_i)| \int_{|x_0 - y_i|}^{\infty} \frac{dt}{t^{n+1}} dy_i \\ &\approx \int_{2r}^{\infty} \prod_{i=1}^m \int_{2r \leq |x_0 - y_i| < t} |f_i(y_i)| dy_i \frac{dt}{t^{n+1}} \lesssim \int_{2r}^{\infty} \prod_{i=1}^m \int_{B(x_0, t)} |f_i(y_i)| dy_i \frac{dt}{t^{n+1}}. \end{aligned}$$

Applying Hölder's inequality, we get

$$\begin{aligned} \int_{(\mathbb{C}(2B))^m} \prod_{i=1}^m \frac{|f_i(y_i)|}{|x_0 - y_i|^n} dy_i &\lesssim \int_{2r}^{\infty} \prod_{i=1}^m \|f_i\|_{L_{p_i}(B(x_0,t))} \|1\|_{L_{p'_i}(B(x_0,t))} \frac{dt}{t^{n+1}} \\ &\approx \int_{2r}^{\infty} t^{-\frac{n}{p}} \prod_{i=1}^m \|f_i\|_{L_{p_i}(B(x_0,t))} \frac{dt}{t}. \end{aligned} \quad (3.5)$$

We now consider the cases when exactly l of the β_i 's are ∞ for some $1 \leq l < m$. We only give the arguments for one of these cases. The rest are similar and can easily be obtained from the arguments below by permuting the indices. To this end we may assume that $\beta_1 = \dots = \beta_l = \infty$ and $\beta_{l+1} = \dots = \beta_m = 0$. Recall the fact that $|x_0 - y_i| \approx |x - y_i|$ for $x \in B$, $y_i \in \mathbb{C}(2B)$ and $1 \leq i \leq l$. We have

$$\begin{aligned} &|T(f_1^\infty, \dots, f_l^\infty, f_{l+1}^0, \dots, f_m^0)(x)| \\ &\lesssim \int_{(\mathbb{C}(2B))^l} \int_{(2B)^{m-l}} \frac{|f_1(y_1) \cdots f_m(y_m)| d\vec{y}}{(|x_0 - y_1| + \cdots + |x_0 - y_l|)^{mn}} \\ &\lesssim r^{-(m-l)n} \int_{(\mathbb{C}(2B))^l} \frac{|f_1(y_1) \cdots f_l(y_l)| dy_1 \cdots dy_l}{(|x_0 - y_1| + \cdots + |x_0 - y_l|)^{ln}} \\ &\quad \times \int_{(2B)^{m-l}} |f_1(y_{l+1}) \cdots f_m(y_m)| dy_{l+1} \cdots dy_m \\ &\lesssim \int_{(\mathbb{C}(2B))^l} \prod_{i=1}^l \frac{|f_i(y_i)|}{|x_0 - y_i|^n} dy_i r^{-n(m-l)} \int_{(2B)^{m-l}} \prod_{i=l+1}^m |f_i(y_i)| dy_i. \end{aligned}$$

Applying Hölder's inequality, we get

$$\begin{aligned} \int_{(\mathbb{C}(2B))^l} \prod_{i=1}^l \frac{|f_i(y_i)|}{|x_0 - y_i|^n} dy_i &\lesssim \int_{2r}^{\infty} \prod_{i=1}^l \|f_i\|_{L_{p_i}(B(x_0,t))} \|1\|_{L_{p'_i}(B(x_0,t))} \frac{dt}{t^{n+1}} \\ &\leq \int_{2r}^{\infty} t^{-\sum_{i=1}^l \frac{n}{p_i}} \prod_{i=1}^l \|f_i\|_{L_{p_i}(B(x_0,t))} \frac{dt}{t} \end{aligned} \quad (3.6)$$

and

$$\begin{aligned} r^{-n(m-l)} \int_{(2B)^{m-l}} \prod_{i=l+1}^m |f_i(y_i)| dy_i &\lesssim \prod_{i=l+1}^m \|f_i\|_{L_{p_i}(2B)} \|1\|_{L_{p'_i}(2B)} \int_{2r}^{\infty} \frac{dt}{t^{n+1}} \\ &\leq \int_{2r}^{\infty} \prod_{i=l+1}^m \|f_i\|_{L_{p_i}(B(x_0,t))} \|1\|_{L_{p'_i}(B(x_0,t))} \frac{dt}{t^{n+1}} \\ &\leq \int_{2r}^{\infty} t^{-\sum_{i=l+1}^m \frac{n}{p_i}} \prod_{i=l+1}^m \|f_i\|_{L_{p_i}(B(x_0,t))} \frac{dt}{t}. \end{aligned} \quad (3.7)$$

Therefore, from (3.5), (3.6) and (3.7) we get second-term of the right-hand side $\sum_{\beta_1, \dots, \beta_m} T(f_1^{\beta_1}, \dots, f_m^{\beta_m})(x)$

converges absolutely for any $\vec{f} \in L_{p_1}^{\text{loc}}(\mathbb{R}^n) \times \dots \times L_{p_m}^{\text{loc}}(\mathbb{R}^n)$, and almost every $x \in \mathbb{R}^n$, and therefore we get the right-hand side of (3.4) is finite.

We will show that the definition is independent of the choice of B . Let $\mathcal{B} = \{B(x, r) : x \in \mathbb{R}^n, r > 0\}$. For simplicity, let's assume that $m = 2$. That is, if $B_1, B_2 \in \mathcal{B}$ and $x \in B_1 \cap B_2$, then

$$\begin{aligned} & T_0(f_1\chi_{2B_1}, f_2\chi_{2B_1})(x) + T(f_1\chi_{2B_1}, f_2\chi_{\mathfrak{c}_{(2B_1)}})(x) \\ & + T(f_1\chi_{\mathfrak{c}_{(2B_1)}}, f_2\chi_{2B_1})(x) + T(f_1\chi_{\mathfrak{c}_{(2B_1)}}, f_2\chi_{\mathfrak{c}_{(2B_1)}})(x) \\ & = T_0(f_1\chi_{2B_2}, f_2\chi_{2B_2})(x) + T(f_1\chi_{2B_2}, f_2\chi_{\mathfrak{c}_{(2B_2)}})(x) \\ & + T(f_1\chi_{\mathfrak{c}_{(2B_2)}}, f_2\chi_{2B_2})(x) + T(f_1\chi_{\mathfrak{c}_{(2B_2)}}, f_2\chi_{\mathfrak{c}_{(2B_2)}})(x). \end{aligned} \quad (3.8)$$

Actually, let $B_3 \in \mathcal{B}$ be selected so that $2B_1 \cup 2B_2 \subset B_3$.

Since $f_1\chi_{2B_1}, f_1\chi_{B_3 \setminus 2B_1} \in L_{p_1, w_1}(\mathbb{R}^n)$, $f_2\chi_{2B_1}, f_2\chi_{B_3 \setminus 2B_1} \in L_{p_2, w_2}(\mathbb{R}^n)$, the linearity of T_0 on $L_{p_1, w_1}(\mathbb{R}^n) \times L_{p_2, w_2}(\mathbb{R}^n)$ yields

$$\begin{aligned} & T_0(f_1\chi_{2B_1}, f_2\chi_{2B_1})(x) + T(f_1\chi_{2B_1}, f_2\chi_{\mathfrak{c}_{(2B_1)}})(x) + T(f_1\chi_{\mathfrak{c}_{(2B_1)}}, f_2\chi_{2B_1})(x) \\ & + T(f_1\chi_{\mathfrak{c}_{(2B_1)}}, f_2\chi_{\mathfrak{c}_{(2B_1)}})(x) = T_0(f_1\chi_{2B_1}, f_2\chi_{2B_1})(x) + T(f_1\chi_{2B_1}, f_2\chi_{B_3 \setminus 2B_1})(x) \\ & + T(f_1\chi_{2B_1}, f_2\chi_{\mathfrak{c}_{B_3}})(x) + T(f_1\chi_{B_3 \setminus 2B_1}, f_2\chi_{2B_1})(x) + T(f_1\chi_{\mathfrak{c}_{B_3}}, f_2\chi_{2B_1})(x) \\ & + T(f_1\chi_{B_3 \setminus 2B_1}, f_2\chi_{B_3 \setminus 2B_1})(x) + T(f_1\chi_{\mathfrak{c}_{B_3}}, f_2\chi_{B_3 \setminus 2B_1})(x) \\ & + T(f_1\chi_{B_3 \setminus 2B_1}, f_2\chi_{\mathfrak{c}_{B_3}})(x) + T(f_1\chi_{\mathfrak{c}_{B_3}}, f_2\chi_{\mathfrak{c}_{B_3}})(x) \\ & = T_0(f_1\chi_{2B_1}, f_2\chi_{2B_1})(x) + T_0(f_1\chi_{2B_1}, f_2\chi_{B_3 \setminus 2B_1})(x) \\ & + T(f_1\chi_{2B_1}, f_2\chi_{\mathfrak{c}_{B_3}})(x) + T_0(f_1\chi_{B_3 \setminus 2B_1}, f_2\chi_{2B_1})(x) + T(f_1\chi_{\mathfrak{c}_{B_3}}, f_2\chi_{2B_1})(x) \\ & + T_0(f_1\chi_{B_3 \setminus 2B_1}, f_2\chi_{B_3 \setminus 2B_1})(x) + T(f_1\chi_{\mathfrak{c}_{B_3}}, f_2\chi_{B_3 \setminus 2B_1})(x) \\ & + T(f_1\chi_{B_3 \setminus 2B_1}, f_2\chi_{\mathfrak{c}_{B_3}})(x) + T(f_1\chi_{\mathfrak{c}_{B_3}}, f_2\chi_{\mathfrak{c}_{B_3}})(x) \\ & = T_0(f_1\chi_{B_3}, f_2\chi_{B_3})(x) + T(f_1\chi_{B_3}, f_2\chi_{\mathfrak{c}_{B_3}})(x) \\ & + T(f_1\chi_{\mathfrak{c}_{B_3}}, f_2\chi_{B_3})(x) + T(f_1\chi_{\mathfrak{c}_{B_3}}, f_2\chi_{\mathfrak{c}_{B_3}})(x). \end{aligned} \quad (3.9)$$

Similarly, we also get

$$\begin{aligned} & T_0(f_1\chi_{2B_2}, f_2\chi_{2B_2})(x) + T(f_1\chi_{2B_2}, f_2\chi_{\mathfrak{c}_{(2B_2)}})(x) + T(f_1\chi_{\mathfrak{c}_{(2B_2)}}, f_2\chi_{2B_2})(x) \\ & + T(f_1\chi_{\mathfrak{c}_{(2B_2)}}, f_2\chi_{\mathfrak{c}_{(2B_2)}})(x) = T_0(f_1\chi_{2B_2}, f_2\chi_{2B_2})(x) + T(f_1\chi_{2B_2}, f_2\chi_{B_3 \setminus 2B_2})(x) \\ & + T(f_1\chi_{2B_2}, f_2\chi_{\mathfrak{c}_{B_3}})(x) + T(f_1\chi_{B_3 \setminus 2B_2}, f_2\chi_{2B_2})(x) + T(f_1\chi_{\mathfrak{c}_{B_3}}, f_2\chi_{2B_2})(x) \\ & + T(f_1\chi_{B_3 \setminus 2B_2}, f_2\chi_{B_3 \setminus 2B_2})(x) + T(f_1\chi_{\mathfrak{c}_{B_3}}, f_2\chi_{B_3 \setminus 2B_2})(x) \\ & + T(f_1\chi_{B_3 \setminus 2B_2}, f_2\chi_{\mathfrak{c}_{B_3}})(x) + T(f_1\chi_{\mathfrak{c}_{B_3}}, f_2\chi_{\mathfrak{c}_{B_3}})(x) \\ & = T_0(f_1\chi_{2B_2}, f_2\chi_{2B_2})(x) + T_0(f_1\chi_{2B_2}, f_2\chi_{B_3 \setminus 2B_2})(x) \\ & + T(f_1\chi_{2B_2}, f_2\chi_{\mathfrak{c}_{B_3}})(x) + T_0(f_1\chi_{B_3 \setminus 2B_2}, f_2\chi_{2B_2})(x) + T(f_1\chi_{\mathfrak{c}_{B_3}}, f_2\chi_{2B_2})(x) \\ & + T_0(f_1\chi_{B_3 \setminus 2B_2}, f_2\chi_{B_3 \setminus 2B_2})(x) + T(f_1\chi_{\mathfrak{c}_{B_3}}, f_2\chi_{B_3 \setminus 2B_2})(x) \\ & + T(f_1\chi_{B_3 \setminus 2B_2}, f_2\chi_{\mathfrak{c}_{B_3}})(x) + T(f_1\chi_{\mathfrak{c}_{B_3}}, f_2\chi_{\mathfrak{c}_{B_3}})(x) \\ & = T_0(f_1\chi_{B_3}, f_2\chi_{B_3})(x) + T(f_1\chi_{B_3}, f_2\chi_{\mathfrak{c}_{B_3}})(x) \\ & + T(f_1\chi_{\mathfrak{c}_{B_3}}, f_2\chi_{B_3})(x) + T(f_1\chi_{\mathfrak{c}_{B_3}}, f_2\chi_{\mathfrak{c}_{B_3}})(x). \end{aligned} \quad (3.10)$$

Since T_m is an m -linear operator, we split $T_m(\vec{f})$ as follows:

$$\left| T_m(\vec{f})(y) \right| \leq \left| T_m(f_1^0, \dots, f_m^0)(y) \right| + \left| \sum'_{\beta_1, \dots, \beta_m} T_m(f_1^{\beta_1}, \dots, f_m^{\beta_m})(y) \right|,$$

where $\beta_1, \dots, \beta_m \in \{0, \infty\}$ and each term in \sum' contains at least one $\beta_i \neq 0$. Then,

$$\begin{aligned} \|T_m(\vec{f})\|_{L_p(B(x,r))} &\leq \|T_m(f_1^0, \dots, f_m^0)\|_{L_p(B(x,r))} \\ &+ \left\| \sum'_{\beta_1, \dots, \beta_m} T_m(f_1^{\beta_1}, \dots, f_m^{\beta_m}) \right\|_{L_p(B(x,r))}, \end{aligned}$$

and

$$\begin{aligned} \|T_m(\vec{f})\|_{WL_p(B(x,r))} &\leq \|T_m(f_1^0, \dots, f_m^0)\|_{WL_p(B(x,r))} \\ &+ \left\| \sum'_{\beta_1, \dots, \beta_m} T_m(f_1^{\beta_1}, \dots, f_m^{\beta_m}) \right\|_{WL_p(B(x,r))}. \end{aligned}$$

Applying Theorem 2.1 we get for $p_1, \dots, p_m \in (1, \infty)$, and $p \in (0, \infty)$ with $1/p = \sum_{i=1}^m 1/p_i$

$$\|T_m(\vec{f}^0)\|_{L_p(B(x,r))} \leq \|T_m(\vec{f}^0)\|_{L_p(\mathbb{R}^n)} \lesssim \prod_{i=1}^m \|f_i^0\|_{L_{p_i}(\mathbb{R}^n)} \lesssim \prod_{i=1}^m \|f_i\|_{L_{p_i}(B(x,2r))}$$

and for $p_1, \dots, p_m \in [1, \infty)$, $\min\{p_1, \dots, p_m\} = 1$, and $p \in (0, \infty)$ with $1/p = \sum_{i=1}^m 1/p_i$

$$\|T_m(\vec{f}^0)\|_{WL_p(B(x,r))} \leq \|T_m(\vec{f}^0)\|_{WL_p(\mathbb{R}^n)} \lesssim \prod_{i=1}^m \|f_i^0\|_{L_{p_i}(\mathbb{R}^n)} \lesssim \prod_{i=1}^m \|f_i\|_{L_{p_i}(B(x,2r))}.$$

On the other hand, we have

$$\begin{aligned} \prod_{i=1}^m \|f_i\|_{L_{p_i}(2B)} &\approx r^{\frac{n}{p}} \prod_{i=1}^m \|f_i\|_{L_{p_i}(2B)} \int_{2r}^{\infty} \frac{dt}{t^{\frac{n}{p_i}+1}} \\ &\leq r^{\frac{n}{p}} \int_{2r}^{\infty} t^{-\frac{n}{p}} \prod_{i=1}^m \|f_i\|_{L_{p_i}(B(x,t))} \frac{dt}{t}. \end{aligned} \quad (3.11)$$

Thus

$$\|T_m(\vec{f}^0)\|_{L_p(B(x,r))} \lesssim r^{\frac{n}{p}} \int_{2r}^{\infty} t^{-\frac{n}{p}} \prod_{i=1}^m \|f_i\|_{L_{p_i}(B(x,t))} \frac{dt}{t}. \quad (3.12)$$

For the other terms, let us first deal with the case when $\beta_1 = \dots = \beta_m = \infty$.

When $|x - y_i| \leq r$, $|z - y_i| \geq 2r$, we have $\frac{1}{2}|z - y_i| \leq |x - y_i| \leq \frac{3}{2}|z - y_i|$, and therefore,

$$\begin{aligned} |T_m(f_1^\infty, \dots, f_m^\infty)(z)| &\lesssim \int_{(\mathbb{C}_{B(x,2r)})^m} \frac{|f_1(y_1) \cdots f_m(y_m)|}{|(x - y_1, \dots, x - y_m)|^{mn}} dy_1 \cdots dy_m \\ &\lesssim \int_{(\mathbb{C}_{B(x,2r)})^m} \prod_{i=1}^m \frac{|f_i(y_i)|}{|x - y_i|^n} dy_i \end{aligned}$$

and

$$\begin{aligned} \|T_m(f_1^\infty, \dots, f_m^\infty)\|_{L_p(B(x,r))} &\leq \int_{(\mathbb{C}_{B(x,2r)})^m} \prod_{i=1}^m \frac{|f_i(y_i)|}{|x - y_i|^n} dy_i \|\chi_{B(x,r)}\|_{L_p} \\ &\lesssim r^{\frac{n}{p}} \int_{(\mathbb{C}_{B(x,2r)})^m} \prod_{i=1}^m \frac{|f_i(y_i)|}{|x - y_i|^n} dy_i. \end{aligned}$$

By Fubini's theorem we have

$$\begin{aligned} \int_{(\mathbb{C}_{B(x,2r)})^m} \prod_{i=1}^m \frac{|f_i(y_i)|}{|x - y_i|^n} dy_i &\approx \int_{(\mathbb{C}_{B(x,2r)})^m} \prod_{i=1}^m |f_i(y_i)| \int_{|x-y_i|}^{\infty} \frac{dt}{t^{n+1}} dy_i \\ &\approx \int_{2r}^{\infty} \prod_{i=1}^m \int_{2r \leq |x-y_i| < t} |f_i(y_i)| dy_i \frac{dt}{t^{n+1}} \lesssim \int_{2r}^{\infty} \prod_{i=1}^m \int_{B(x,t)} |f_i(y_i)| dy_i \frac{dt}{t^{n+1}}. \end{aligned}$$

Applying Hölder's inequality, we get

$$\begin{aligned} \int_{(\mathbb{C}_{B(x,2r)})^m} \prod_{i=1}^m \frac{|f_i(y_i)|}{|x - y_i|^n} dy_i &\lesssim \int_{2r}^{\infty} \prod_{i=1}^m \|f_i\|_{L_{p_i}(B(x,t))} \|1\|_{L_{p'_i}(B(x,t))} \frac{dt}{t^{n+1}} \\ &\leq \int_{2r}^{\infty} t^{-\frac{n}{p}} \prod_{i=1}^m \|f_i\|_{L_{p_i}(B(x,t))} \frac{dt}{t}. \end{aligned} \quad (3.13)$$

Moreover, for all $p_i \in [1, \infty)$, $i = 1, \dots, m$, the inequality

$$\|T_m(f_1^\infty, \dots, f_m^\infty)\|_{L_p(B(x,r))} \lesssim r^{\frac{n}{p}} \int_{2r}^{\infty} t^{-\frac{n}{p}} \prod_{i=1}^m \|f_i\|_{L_{p_i}(B(x,t))} \frac{dt}{t} \quad (3.14)$$

is valid.

We now consider the cases when exactly l of the β_i 's are ∞ for some $1 \leq l < m$. We only give the arguments for one of these cases. The rest are similar and can easily be obtained from the arguments below by permuting the indices. To this end we may assume that $\beta_1 = \dots = \beta_l = \infty$ and $\beta_{l+1} = \dots = \beta_m = 0$. Recall the fact that $|x - y_i| \approx |z - y_i|$

for $z \in B(x, r)$, $y_i \in {}^c B(x, 2r)$ and $1 \leq i \leq l$. We have

$$\begin{aligned}
& |T_m(f_1^\infty, \dots, f_l^\infty, f_{l+1}^0, \dots, f_m^0)(z)| \\
& \lesssim \int_{({}^c B(x, 2r))^l} \int_{(B(x, 2r))^{m-l}} \frac{|f_1(y_1) \cdots f_m(y_m)| dy_1 \cdots dy_m}{(|x - y_1| + \cdots + |x - y_l|)^{mn}} \\
& \lesssim r^{-(m-l)n} \int_{({}^c B(x, 2r))^l} \frac{|f_1(y_1) \cdots f_l(y_l)| dy_1 \cdots dy_l}{(|x - y_1| + \cdots + |x - y_l|)^{ln}} \\
& \times \int_{(B(x, 2r))^{m-l}} |f_1(y_{l+1}) \cdots f_m(y_m)| dy_{l+1} \cdots dy_m \\
& \lesssim \int_{({}^c B(x, 2r))^l} \prod_{i=1}^l \frac{|f_i(y_i)|}{|x - y_i|^n} dy_i r^{-n(m-l)} \int_{(B(x, 2r))^{m-l}} \prod_{i=l+1}^m |f_i(y_i)| dy_i.
\end{aligned}$$

Applying Hölder's inequality, we get

$$\begin{aligned}
& \int_{({}^c B(x, 2r))^l} \prod_{i=1}^l \frac{|f_i(y_i)|}{|x - y_i|^n} dy_i \lesssim \int_{2r}^\infty \prod_{i=1}^l \|f_i\|_{L_{p_i}(B(x, t))} \|1\|_{L_{p'_i}(B(x, t))} \frac{dt}{t^{n+1}} \\
& \leq \int_{2r}^\infty t^{-\sum_{i=1}^l \frac{n}{p_i}} \prod_{i=1}^l \|f_i\|_{L_{p_i}(B(x, t))} \frac{dt}{t} \tag{3.15}
\end{aligned}$$

and

$$\begin{aligned}
& r^{-n(m-l)} \int_{(B(x, 2r))^{m-l}} \prod_{i=l+1}^m |f_i(y_i)| dy_i \\
& \lesssim \prod_{i=l+1}^m \|f_i\|_{L_{p_i}(B(x, 2r))} \|1\|_{L_{p'_i}(B(x, 2r))} \int_{2r}^\infty \frac{dt}{t^{nl+1}} \\
& \leq \int_{2r}^\infty \prod_{i=l+1}^m \|f\|_{L_{p_i}(B(x, t))} \|1\|_{L_{p'_i}(B(x, t))} \frac{dt}{t^{nl+1}} \\
& \leq \int_{2r}^\infty \prod_{i=l+1}^m \|f\|_{L_{p_i}(B(x, t))} |B(x, t)|^{-\frac{1}{p_i}} \frac{dt}{t}. \tag{3.16}
\end{aligned}$$

From (3.15) and (3.16) we get

$$\begin{aligned}
& \|T_m(f_1^\infty, \dots, f_l^\infty, f_{l+1}^0, \dots, f_m^0)\|_{L_p(B(x, r))} \\
& \lesssim r^{\frac{n}{p}} \int_{({}^c B(x, 2r))^l} \prod_{i=1}^l \frac{|f_i(y_i)|}{|x - y_i|^n} dy_i r^{-n(m-l)} \int_{(B(x, 2r))^{m-l}} |f_i(y_i)| dy_i \\
& \lesssim r^{\frac{n}{p}} \int_{2r}^\infty t^{-\frac{n}{p}} \prod_{i=1}^m \|f\|_{L_{p_i}(B(x, t))} \frac{dt}{t}.
\end{aligned}$$

Thus we get the following inequality:

$$\left\| \sum_{\beta_1, \dots, \beta_m} T_m(f_1^{\beta_1}, \dots, f_m^{\beta_m}) \right\|_{L_p(B(x, r))} \lesssim r^{\frac{n}{p}} \int_{2r}^\infty t^{-\frac{n}{p}} \prod_{i=1}^m \|f\|_{L_{p_i}(B(x, t))} \frac{dt}{t}.$$

Consequently, the inequality (3.1) is valid.

Proof of Theorem 1.1. For $(p_1, \dots, p_m) \in (1, \infty)^m$ from Theorem 2.3 and Theorem 3.1 (see, inequality (3.1)) we get

$$\begin{aligned} \|T(\mathbf{f})\|_{M_{p, \varphi_2}^{\{x_0\}}} &\lesssim \sup_{r>0} \varphi_2(x_0, r)^{-1} \int_r^\infty t^{-\frac{n}{p}} \prod_{i=1}^m \|f_i\|_{L_{p_i}(B(x_0, t))} \frac{dt}{t} \\ &\lesssim \sup_{r>0} t^{-\frac{n}{p}} \prod_{i=1}^m \varphi_{1i}(x, r)^{-1} \|f_i\|_{L_{p_i}(B(x_0, r))} \lesssim \prod_{i=1}^m \|f_i\|_{M_{p_i, \varphi_{1i}}^{\{x_0\}}}. \end{aligned}$$

For $(p_1, \dots, p_m) \in [1, \infty)^m$, $\min\{p_1, \dots, p_m\} = 1$ from Theorem 2.3 and Theorem 3.1 (see, inequality (3.2)) we get

$$\begin{aligned} \|T(\mathbf{f})\|_{WM_{p, \varphi_2}^{\{x_0\}}} &\lesssim \sup_{r>0} \varphi_2(x_0, r)^{-1} \int_r^\infty t^{-\frac{n}{p}} \prod_{i=1}^m \|f_i\|_{L_{p_i}(B(x_0, t))} \frac{dt}{t} \\ &\lesssim \sup_{r>0} t^{-\frac{nm}{p}} \prod_{i=1}^m \varphi_{1i}(x_0, r)^{-1} \|f_i\|_{L_{p_i}(B(x_0, r))} \lesssim \prod_{i=1}^m \|f_i\|_{M_{p_i, \varphi_{1i}}^{\{x_0\}}}. \end{aligned}$$

This completes the proof of Theorem 1.1.

Now we prove a theorem below such that we prove Theorem 1.2 with the help of this theorem.

Theorem 3.2 *Let $m \geq 2$, $x_0 \in \mathbb{R}^n$, T be an m -linear ω -CZO, and ω satisfy (1.5). If $p_1, \dots, p_m \in (1, \infty)$ and $p \in (0, \infty)$ with $1/p = \sum_{i=1}^m 1/p_i$, then the inequality*

$$\begin{aligned} \|T_{II\mathbf{b}}(\mathbf{f})\|_{L_p(B(x_0, r))} &\lesssim \|\mathbf{b}\|_* r^{\frac{n}{p}} \\ &\times \int_{2r}^\infty \left(1 + \ln \frac{t}{r}\right)^m t^{-\frac{n}{p}} \prod_{i=1}^m \|f_i\|_{L_{p_i}(B(x_0, t))} \frac{dt}{t} \end{aligned} \quad (3.17)$$

holds for any balls $B(x_0, r)$ and for all $\vec{f} \in L_{p_1}^{loc}(\mathbb{R}^n) \times \dots \times L_{p_m}^{loc}(\mathbb{R}^n)$.

Proof. For simplicity of the expansion, we only present the case $m = 2$.

We represent f_i as $f_i = f_i^0 + f_i^\infty$, where $f_i^0 = f_i \chi_{B(x_0, 2r)}$, $i = 1, 2$, and $\chi_{B(x_0, 2r)}$ denotes the characteristic function of $B(x_0, 2s)$. Then,

$$\begin{aligned} \|T_{II\mathbf{b}}(\mathbf{f})\|_{L_p(B(x_0, r))} &\leq \left(\int_{B(x_0, r)} |T_{II\mathbf{b}}(f_1^0, f_2^0)(x)|^p dx \right)^{1/p} \\ &+ \left(\int_{B(x_0, r)} |T_{II\mathbf{b}}(f_1^0, f_2^\infty)(x)|^p dx \right)^{1/p} + \left(\int_{B(x_0, r)} |T_{II\mathbf{b}}(f_1^\infty, f_2^0)(x)|^p dx \right)^{1/p} \\ &+ \left(\int_{B(x_0, r)} |T_{II\mathbf{b}}(f_1^\infty, f_2^\infty)(x)|^p dx \right)^{1/p} = I + II + III + IV. \end{aligned} \quad (3.18)$$

Since $T_{II\mathbf{b}}$ bounded from $L_{p_1} \times L_{p_2}$ to L_p , we get

$$\left(\int_{B(x_0, r)} |T_{II\mathbf{b}}(f_1^0, f_2^0)(x)|^p dx \right)^{1/p} \lesssim \|\mathbf{b}\|_* \prod_{i=1}^m \|f_i\|_{L_{p_i}(B(x_0, 2r))}. \quad (3.19)$$

Then, by (3.11), we get

$$I \lesssim \|\mathbf{b}\|_* r^{\frac{n}{p}} \int_{2r}^{\infty} t^{-\frac{n}{p}} \prod_{i=1}^2 \|f_i\|_{L_{p_i}(B(x_0, t))} \frac{dt}{t}. \quad (3.20)$$

Owing to the symmetry of II and III, we only estimate II. Taking $\lambda_i = (b_i)_{B(x_0, s)}$, then

$$\begin{aligned} T_{II\mathbf{b}}(f_1^0, f_2^\infty)(x) &= (b_1(x) - \lambda_1)(b_2(x) - \lambda_2)T(f_1^0, f_2^\infty)(x) \\ &\quad - (b_1(x) - \lambda_1)T(f_1^0, (b_2 - \lambda_2)f_2^\infty)(x) - (b_2(x) - \lambda_2)T((b_1 - \lambda_1)f_1^0, f_2^\infty)(x) \\ &\quad + T((b_1 - \lambda_1)f_1^0, (b_2 - \lambda_2)f_2^\infty)(x) = II_1 + II_2 + II_3 + II_4. \end{aligned} \quad (3.21)$$

Similar to the estimate of (3.14), for any $x \in B(x_0, s)$, we can deduce

$$\sup_{x \in B(x_0, r)} |T(f_1^0, f_2^\infty)(x)| \lesssim \int_{2r}^{\infty} t^{-\frac{n}{p}} \prod_{i=1}^2 \|f_i\|_{L_{p_i}(B(x_0, t))} \frac{dt}{t}. \quad (3.22)$$

Applying Hölder's inequality and Lemma 2.1, we have

$$\begin{aligned} &\left(\int_{B(x_0, r)} |(b_1(x) - \lambda_1)(b_2(x) - \lambda_2)|^p dx \right)^{1/p} \\ &\lesssim \prod_{i=1}^2 \left(\int_{B(x_0, r)} |b_i(x) - \lambda_i|^{2p} dx \right)^{1/2p} \lesssim \|\mathbf{b}\|_*. \end{aligned} \quad (3.23)$$

Then, by (3.22) and (3.23) we have

$$\begin{aligned} &\left(\int_{B(x_0, r)} |II_1|^p dx \right)^{1/p} \leq \left(\int_{B(x_0, r)} |(b_1(x) - \lambda_1)(b_2(x) - \lambda_2)|^p dx \right)^{1/p} \\ &\quad \times \sup_{x \in B(x_0, r)} |T(f_1^0, f_2^\infty)(x)| \lesssim \|\mathbf{b}\|_* r^{\frac{n}{p}} \int_{2r}^{\infty} t^{-\frac{n}{p}} \prod_{i=1}^2 \|f_i\|_{L_{p_i}(B(x_0, t))} \frac{dt}{t}. \end{aligned} \quad (3.24)$$

Applying the size condition, we deduce that, for any $x \in B(x_0, r)$,

$$\begin{aligned} &|T(f_1^0, (b_2 - \lambda_2)f_2^\infty)(x)| \quad (3.25) \\ &\lesssim \int_{B(x_0, 2r)} \int_{\mathbb{R}^n \setminus B(x_0, 2r)} \frac{|f_1(y_1)(b_2(y_2) - \lambda_2)f_2(y_2)|}{(|x - y_1| + |x - y_2|)^{2n}} dy_1 dy_2 \\ &\lesssim \sum_{j=1}^{\infty} (2^{j+1}r)^{-2n} \int_{B(x_0, 2^{j+1}r)} |f_1(y_1)| dy_1 \int_{B(x_0, 2^{j+1}r)} |(b_2(y_2) - \lambda_2)f_2(y_2)| dy_2. \end{aligned}$$

Note that

$$\begin{aligned} &\int_{B(x_0, 2^{j+1}r)} |f_1(y_1)| dy_1 \leq C \|f_1\|_{L_{p_1}(B(x_0, 2^{j+1}r))} \|1\|_{L_{p'_1}(B(x_0, 2^{j+1}r))}, \\ &\int_{B(x_0, 2^{j+1}r)} |(b_2(y_2) - \lambda_2)f_2(y_2)| dy_2 \quad (3.26) \\ &\leq C \|f_2\|_{L_{p_2}(B(x_0, 2^{j+1}r))} \|b_2(\cdot) - \lambda_2\|_{L_{p'_2}(B(x_0, 2^{j+1}r))}. \end{aligned}$$

Then,

$$\begin{aligned} \sup_{x \in B(x_0, r)} |T(f_1^0, (b_2 - \lambda_2)f_2^\infty)(x)| &\leq C \int_{2r}^\infty \prod_{i=1}^2 \|f_i\|_{L_{p_i}(B(x_0, t))} \\ &\times \|1\|_{L_{p'_1}(B(x_0, t))} \|b_2(\cdot) - \lambda_2\|_{L_{p'_2}(B(x_0, t))} \frac{dt}{t^{n+1}}. \end{aligned} \quad (3.27)$$

From Lemma 2.1, we get

$$\begin{aligned} \|b_2(\cdot) - \lambda_2\|_{L_{p'_2}(B(x_0, r))} &\leq C \left(\int_{B(x_0, r)} |b_2(z) - \lambda_2|^{p'_2} dz \right)^{1/p'_2} \\ &\leq C \left(1 + \left| \ln \frac{t}{r} \right| \right) \|b_2\|_* |B(x_0, r)|^{1/p'_2}. \end{aligned} \quad (3.28)$$

Note that

$$\prod_{i=1}^2 \|1\|_{L_{p'_i}(B(x_0, r))} = |B(x_0, r)|^2 \prod_{i=1}^2 |B(x_0, r)|^{-\frac{1}{p_i}}. \quad (3.29)$$

From (3.27), (3.28), and (3.29), we can deduce

$$\begin{aligned} \sup_{x \in B(x_0, r)} |T(f_1^0, (b_2 - \lambda_2)f_2^\infty)(x)| \\ \lesssim \|b_2\|_* \int_{2r}^\infty \left(1 + \ln \frac{t}{r} \right) r^{-\frac{n}{p}} \prod_{i=1}^2 \|f_i\|_{L_{p_i}(B(x_0, t))} \frac{dt}{t}. \end{aligned} \quad (3.30)$$

Applying Lemma 2.1 and using Hölder's inequality, we have

$$\left(\int_{B(x_0, r)} |b_1(x) - \lambda_1|^p dx \right) \lesssim \|b_1\|_* \prod_{i=1}^2 |B(x_0, r)|^{\frac{1}{p_i}}. \quad (3.31)$$

Then, by (3.30),

$$\begin{aligned} \left(\int_{B(x_0, r)} |II_2|^p dx \right)^{1/p} &\leq \left(\int_{B(x_0, r)} |(b_1(x) - \lambda_1)|^p dx \right)^{1/p} \\ &\times \sup_{x \in B(x_0, r)} |T(f_1^0, (b_2 - \lambda_2)f_2^\infty)(x)| \lesssim \prod_{i=1}^2 \|b_i\|_* |B(x_0, r)|^{\frac{1}{p_i}} \\ &\times \int_{2r}^\infty \left(1 + \ln \frac{t}{r} \right) \left(\prod_{i=1}^2 \|f_i\|_{L_{p_i}(B(x_0, t))} |B(x_0, r)|^{-\frac{1}{p_i}} \right) \frac{dt}{t} \\ &\approx \|b\|_* r^{\frac{n}{p}} \int_{2r}^\infty t^{-\frac{n}{p}} \prod_{i=1}^2 \|f_i\|_{L_{p_i}(B(x_0, t))} \frac{dt}{t}. \end{aligned} \quad (3.32)$$

Similarly, we also have that

$$\begin{aligned}
& \left(\int_{B(x_0, r)} |II_3|^p dx \right)^{1/p} \leq C \prod_{i=1}^2 \left(\|b_i\|_* |B(x_0, r)|^{\frac{1}{p_i}} \right) \\
& \times \int_{2r}^{\infty} \left(1 + \ln \frac{t}{r} \right) \left(\prod_{i=1}^2 \|f_i\|_{L_{p_i}(B(x_0, t))} |B(x_0, t)|^{-\frac{1}{p_i}} \right) \frac{dt}{t} \\
& \approx \|b\|_* r^{\frac{n}{p}} \int_{2r}^{\infty} t^{-\frac{n}{p}} \prod_{i=1}^2 \|f_i\|_{L_{p_i}(B(x_0, t))} \frac{dt}{t}.
\end{aligned} \tag{3.33}$$

Applying the size condition again, for any $x \in B(x_0, r)$, with the same method of estimate for (3.30), we have

$$\begin{aligned}
|T((b_1 - \lambda_1)f_1^0, (b_2 - \lambda_2)f_2^\infty)(x)| & \lesssim \sum_{j=1}^{\infty} (2^{j+1}r) \prod_{i=1}^2 \int_{B(x_0, 2^{j+1}r)} |(b_i(y) - \lambda_i)f_i(y_i)| dy_i \\
& \lesssim \int_{2r}^{\infty} \left(\prod_{i=1}^2 \|f_i\|_{L_{p_i}(B(x_0, t))} \|b_i(\cdot) - \lambda_i\|_{L_{p'_i}(B(x_0, t))} \right) \frac{dt}{t^{n+1}} \\
& \lesssim \|b\|_* \int_{2r}^{\infty} \left(1 + \ln \frac{t}{r} \right)^2 t^{-\frac{n}{p}} \prod_{i=1}^2 \|f_i\|_{L_{p_i}(B(x_0, t))} \frac{dt}{t}.
\end{aligned} \tag{3.34}$$

Then,

$$\begin{aligned}
& \left(\int_{B(x_0, r)} |II_4|^p dx \right)^{1/p} \lesssim \sup_{x \in B(x_0, r)} |((b_1 - \lambda_1)f_1^0, (b_2 - \lambda_2)f_2^\infty)(x)| \\
& \lesssim \|b\|_* r^{\frac{n}{p}} \int_{2r}^{\infty} \left(1 + \ln \frac{t}{r} \right)^2 t^{-\frac{n}{p}} \prod_{i=1}^2 \|f_i\|_{L_{p_i}(B(x_0, t))} \frac{dt}{t}.
\end{aligned} \tag{3.35}$$

Then, combining (3.24), (3.32), (3.33), and (3.35), we get

$$\begin{aligned}
& \left(\int_{B(x_0, r)} |II|^p dx \right)^{1/p} \lesssim \|b\|_* r^{\frac{n}{p}} \\
& \times \int_{2r}^{\infty} \left(1 + \ln \frac{t}{r} \right)^2 t^{-\frac{n}{p}} \prod_{i=1}^2 \|f_i\|_{L_{p_i}(B(x_0, t))} \frac{dt}{t}.
\end{aligned} \tag{3.36}$$

Finally, we still decompose $T_{IIb}(f_1^\infty, f_2^\infty)(x)$ as follows:

$$\begin{aligned}
T_{IIb}(f_1^\infty, f_2^\infty)(x) & = (b_1(x) - \lambda_1)(b_2(x) - \lambda_2)T(f_1^\infty, f_2^\infty)(x) \\
& - (b_1(x) - \lambda_1)T(f_1^\infty, (b_2 - \lambda_2)f_2^\infty)(x) - (b_2(x) - \lambda_2)T((b_1 - \lambda_1)f_1^\infty, f_2^\infty)(x) \\
& + T((b_1 - \lambda_1)f_1^\infty, (b_2 - \lambda_2)f_2^\infty)(x) = IV_1 + IV_2 + IV_3 + IV_4.
\end{aligned} \tag{3.37}$$

Because each term IV_j is completely analogous to II_j , $j = 1, 2, 3, 4$ with a bit difference, we get the following estimate without details:

$$\begin{aligned}
& \left(\int_{B(x_0, r)} |IV|^p dx \right)^{1/p} \lesssim \|b\|_* r^{\frac{n}{p}} \\
& \times \int_{2r}^{\infty} \left(1 + \ln \frac{t}{r} \right)^2 t^{-\frac{n}{p}} \prod_{i=1}^2 \|f_i\|_{L_{p_i}(B(x_0, t))} \frac{dt}{t}.
\end{aligned} \tag{3.38}$$

Summing up the above estimates, (3.17) is proved.

Proof of Theorem 1.2. For $(p_1, \dots, p_m) \in (1, \infty)^m$ from Theorem 2.4 and Theorem 3.2 (see, inequality (3.17)) we get

$$\begin{aligned} \|T_{\Pi b}(\mathbf{f})\|_{M_{p, \varphi_2}^{\{x_0\}}} &\lesssim \sup_{r>0} \varphi_2(x_0, r)^{-1} \int_r^\infty \left(1 + \ln \frac{t}{r}\right)^m t^{-\frac{n}{p}} \prod_{i=1}^m \|f_i\|_{L_{p_i}(B(x_0, t))} \frac{dt}{t} \\ &\lesssim \sup_{r>0} t^{-\frac{n}{p}} \prod_{i=1}^m \varphi_{1i}(x, r)^{-1} \|f_i\|_{L_{p_i}(B(x_0, r))} \lesssim \prod_{i=1}^m \|f_i\|_{M_{p_i, \varphi_{1i}}^{\{x_0\}}}. \end{aligned}$$

This completes the proof of Theorem 1.2.

The proof of Theorem 1.3 can be done using the method in Theorem 1.2, with similar argument.

Acknowledgements.

The authors would like to express their gratitude to the referees for his very valuable comments and suggestions. The research of V. Guliyev and A. Serbetci was partially supported by grant of Cooperation Program 2532 TUBITAK - RFBR (RUSSIAN foundation for basic research) (Agreement number no. 119N455).

References

1. Ahmadli, A.A., Akbulut, A., Isayev, F.A., Serbetci, A.: *Multilinear commutators of parabolic Calderón-Zygmund operators on generalized weighted variable parabolic Morrey spaces*, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci. **41** (4), Mathematics, 3-16 (2021).
2. Coifman, R.R., Meyer, Y.: *On commutators of singular integrals and bilinear singular integrals*, Trans. Amer. Math. Soc. **212**, 315-331 (1975).
3. Coifman, R.R., Meyer, Y.: *Commutateurs d'intégrales singulières et opérateurs multilinéaires*, Ann. Inst. Fourier (Grenoble) **28** (3), 177-202 (1978).
4. Ekinçioğlu, I., Keskin, C., Serbetci, A.: *Multilinear commutators of Calderón-Zygmund operator on generalized variable exponent Morrey spaces*, Positivity **25** (4), 1551-1567 (2021).
5. Ekinçioğlu, I., Umarmhadzhiev, S.: *Oscillatory integrals with variable Calderón-Zygmund kernel on generalized weighted Morrey spaces*, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci. **42** (1), Mathematics, 99-110 (2022).
6. Grafakos, L., Torres, R.H.: *Multilinear Calderón-Zygmund theory*, Adv. Math. **165**, 124-164 (2002).
7. Guliyev, V.S.: *Integral operators on function spaces on the homogeneous groups and on domains in \mathbb{R}^n , [Russian Doctor's degree dissertation]. Moscow: Mat. Inst. Steklov; p. 329 (1994).*
8. Guliyev, V.S.: *Boundedness of the maximal, potential and singular operators in the generalized Morrey spaces*, J Inequal Appl. Art. ID 503948, 20 pp. (2009).
9. Guliyev, V.S.: *Generalized weighted Morrey spaces and higher order commutators of sublinear operators*, Eurasian Math. J. **3** (3), 33-61 (2012).
10. Guliyev, V.S.: *Generalized local Morrey spaces and fractional integral operators with rough kernel*, Problems in mathematical analysis. No. 71. J. Math. Sci. (N.Y.) **193** (2013), no. 2, 211-227.

11. Guliyev, V.S., Ismayilova, A.F.: *Multi-sublinear maximal operator and multilinear singular integral operators on generalized Morrey spaces*, Proc. Inst. Math. Mech. Natl. Acad. Sci. Azerb. **40** (2), 1-15 (2014).
12. Guliyev, V.S., Guliyev, R.V., Omarova, M.N.: *Riesz transforms associated with Schrödinger operator on vanishing generalized Morrey spaces*, Appl. Comput. Math. **17** (1), 56-71 (2018).
13. Ismayilova, A.F.: *Calderón-Zygmund operators with kernels of Dini's type and their multilinear commutators on generalized Morrey spaces*, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci. **41** (4), Mathematics, 100-111 (2021).
14. Ismayilova, A.F., Isayev, F.A.: *Multi-sublinear operators generated by multilinear Calderón-Zygmund operators on product generalized Morrey spaces*, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci. **40** (4), Mathematics, 110-117 (2020).
15. Kenig, C., Stein, E.M.: *Multilinear estimates and fractional integration*, Math. Res. Lett. **6** (1), 1-15 (1999).
16. Lerner, A.K., Ombrosi, S., Perez, C., Torres, R.H., Trujillo-Gonzalez, R.: *New maximal functions and multiple weights for the multilinear Calderón-Zygmund theory*, Adv. Math. **220** (4), 1222-1264 (2009).
17. Liu, Z., Lu, S.: *Endpoint estimates for commutators of Calderón-Zygmund type operators*, Kodai Math. J. **25** (1), 79-88 (2002).
18. Lu, G., Zhang, P.: *Multilinear Calderón-Zygmund operators with kernels of Dini's type and applications*, Nonlinear Analysis **107**, 92-117 (2014).
19. Maldonado, D., Naibo, V.: *Weighted norm inequalities for paraproducts and bilinear pseudodifferential operators with mild regularity*, J. Fourier Anal. Appl. **15**, 218-261 (2009).
20. Mizuhara, T.: *Boundedness of some classical operators on generalized Morrey spaces*, Harmonic Analysis (S. Igari, Editor), ICM 90 Satellite Proceedings. Springer - Verlag, Tokyo, 183-189 (1991).
21. Morrey, C.B.: *On the solutions of quasi-linear elliptic partial differential equations*, Trans. Amer. Math. Soc. **43**, 126-166 (1938).
22. Nakai, E.: *Hardy-Littlewood maximal operator, singular integral operators and Riesz potentials on generalized Morrey spaces*, Math. Nachr. **166**, 95-103 (1994).
23. Perez, C., Pradolini, G., Torres, R.H., Trujillo-Gonzalez, R.: *End-point estimates for iterated commutators of multilinear singular integrals*, Bull. London Math. Soc. **46**, 26-42 (2014).
24. Sawano, Y.: *A thought on generalized Morrey spaces*, J. Indonesian Math. Soc. **25** (3), 210-281 (2019).
25. Yabuta, K.: *Generalizations of Calderón-Zygmund operators*, Studia Math. **82** (1), 17-31 (1985).