

Fractional maximal operator and its higher order commutators on generalized weighted Morrey spaces

Afag F. Ismayilova *

Received: 21.01.2019 / Revised: 28.07.2019 / Accepted: 26.09.2019

Abstract. In this paper, we study the boundedness of the fractional maximal operator M_α and its k th order commutators $[b, M_\alpha]^k$ on generalized weighted Morrey spaces $M_{p,\varphi}(w)$. We find the sufficient conditions on the pair (φ_1, φ_2) with $b \in BMO(\mathbb{R}^n)$ and $w \in A_{pq}$ which ensures the boundedness of the operators M_α and $[b, M_\alpha]^k$ from $M_{p,\varphi_1}(w)$ to $M_{p,\varphi_2}(w)$ for $1 < p < \infty$. In all cases the conditions for the boundedness of the operators M_α , $[b, M_\alpha]^k$ are given in terms of supremal-type inequalities on (φ_1, φ_2) and w , which do not assume any assumption on monotonicity of $\varphi_1(x, r)$, $\varphi_2(x, r)$ in r . The main advance in comparison with the existing results is that we manage to obtain conditions for the boundedness not in integral terms but in less restrictive terms of supremal operators.

Keywords. Generalized weighted Morrey spaces; fractional maximal operator; A_p weights; commutator; BMO

Mathematics Subject Classification (2010): 42B20, 42B35

1 Introduction

The classical Morrey spaces were originally introduced by Morrey in [26] to study the local behavior of solutions to second order elliptic partial differential equations. For the properties and applications of classical Morrey spaces, we refer the readers to [5–9, 12]. Recently, Komori and Shirai [24] considered the weighted Morrey spaces $L^{p,\kappa}(w)$ and studied the boundedness of some classical operators such as the Hardy-Littlewood maximal operator, the Calderón-Zygmund operator on these spaces. Guliyev [13] gave a concept of generalized weighted Morrey space $M_{p,\varphi}(w)$ which could be viewed as extension of both generalized Morrey space $M_{p,\varphi}$ and weighted Morrey space $L^{p,\kappa}(w)$. In [13] Guliyev also studied the boundedness of the classical operators and its commutators in these spaces $M_{p,\varphi}(w)$, see also Guliyev et al. [10, 15–17].

Let $0 < \alpha < n$. The fractional maximal operator M_α and the Riesz potential operator I_α are defined by

$$M_\alpha f(x) = \sup_{t>0} |B(x, t)|^{-1+\frac{\alpha}{n}} \int_{B(x,t)} |f(y)| dy, \quad I_\alpha f(x) = \int_{\mathbb{R}^n} \frac{f(y)}{|x-y|^{n-\alpha}} dy.$$

* Corresponding author

If $\alpha = 0$, then $M \equiv M_0$ is the well known Hardy-Littlewood maximal operator. Recall that, for $0 < \alpha < n$,

$$M_\alpha f(x) \leq v_n^{\frac{\alpha}{n}-1} I_\alpha(|f|)(x).$$

The commutators generated by a suitable function b and the operator M_α is formally defined by

$$[b, M_\alpha]f = M_\alpha(bf) - bM_\alpha(f).$$

Given a measurable function b the fractional maximal commutator operators $M_{b,\alpha}$ is defined by

$$M_{b,\alpha}(f)(x) := \sup_{t>0} |B(x,t)|^{-1+\frac{\alpha}{n}} \int_{B(x,t)} |b(x) - b(y)||f(y)|dy.$$

If $\alpha = 0$, then $M_{b,0} \equiv M_b$ is the sublinear commutator of the Hardy-Littlewood maximal operator.

For a function b defined on \mathbb{R}^n , we denote

$$b^-(x) := \begin{cases} 0, & \text{if } b(x) \geq 0 \\ |b(x)|, & \text{if } b(x) < 0 \end{cases}$$

and $b^+(x) := |b(x)| - b^-(x)$. Obviously, $b^+(x) - b^-(x) = b(x)$.

The following relations between $[b, M_\alpha]$ and $M_{b,\alpha}$ are valid :

Let b be any non-negative locally integrable function. Then

$$|[b, M_\alpha]f(x)| \leq M_{b,\alpha}(f)(x), \quad x \in \mathbb{R}^n$$

holds for all $f \in L_{\text{loc}}^1(\mathbb{R}^n)$.

If b is any locally integrable function on \mathbb{R}^n , then

$$|[b, M_\alpha]f(x)| \leq M_{b,\alpha}(f)(x) + 2b^-(x)M_\alpha f(x), \quad x \in \mathbb{R}^n \quad (1.1)$$

holds for all $f \in L_{\text{loc}}^1(\mathbb{R}^n)$ (see, for example, [29]).

The main purpose of this paper is to study Spanne-Guliyev type boundedness of the operator M_α on the generalized weighted Morrey spaces, including weak versions. Also we study Spanne-Guliyev type boundedness of the k th order commutator operator $[b, M_\alpha]^k$ on the generalized weighted Morrey spaces. We find the sufficient conditions on the pair (φ_1, φ_2) with $b \in BMO(\mathbb{R}^n)$ and $w \in A_{pq}$ which ensures the boundedness of the operators $M_\alpha, [b, M_\alpha]^k$ from $M_{p,\varphi_1}(w)$ to $M_{p,\varphi_2}(w)$ for $1 < p < q < \infty$.

By $A \lesssim B$ we mean that $A \leq CB$ with some positive constant C independent of appropriate quantities. If $A \lesssim B$ and $B \lesssim A$, we write $A \approx B$ and say that A and B are equivalent.

2 Generalized weighted Morrey spaces

The classical Morrey spaces $M_{p,\lambda}$ were originally introduced by Morrey in [26] to study the local behavior of solutions to second order elliptic partial differential equations. For the properties and applications of classical Morrey spaces, we refer the readers to [11, 22, 25].

We recall that a weight function w is in the Muckenhoupt's class A_p [27], $1 < p < \infty$, if

$$\begin{aligned} [w]_{A_p} &:= \sup_B [w]_{A_p(B)} \\ &= \sup_B \left(\frac{1}{|B|} \int_B w(x)dx \right) \left(\frac{1}{|B|} \int_B w(x)^{1-p'} dx \right)^{p-1}, \end{aligned} \quad (2.1)$$

where the sup is taken with respect to all the balls B and $\frac{1}{p} + \frac{1}{p'} = 1$. Note that, for all balls B by Hölder's inequality

$$[w]_{A_p(B)}^{1/p} = |B|^{-1} \|w\|_{L_1(B)}^{1/p} \|w^{-1/p}\|_{L_{p'}(B)} \geq 1. \quad (2.2)$$

For $p = 1$, the class A_1 is defined by the condition $Mw(x) \leq Cw(x)$ with $[w]_{A_1} = \sup_{x \in \mathbb{R}^n} \frac{Mw(x)}{w(x)}$, and for $p = \infty$ $A_\infty = \bigcup_{1 \leq p < \infty} A_p$ and $[w]_{A_\infty} = \inf_{1 \leq p < \infty} [w]_{A_p}$.

A weight function w belongs to the Muckenhoupt-Wheeden class $A_{p,q}$ [28] for $1 < p, q < \infty$ if

$$\begin{aligned} [w]_{A_{p,q}} &:= \sup_B [w]_{A_{p,q}(B)} \\ &= \sup_B \left(\frac{1}{|B|} \int_B w(x)^q dx \right)^{1/q} \left(\frac{1}{|B|} \int_B w(x)^{-p'} dx \right)^{1/p'} < \infty, \end{aligned}$$

where the sup is taken with respect to all balls B . Note that, for all balls B by Hölder's inequality

$$[w]_{A_{p,q}(B)} = |B|^{\frac{1}{p} - \frac{1}{q} - 1} \|w\|_{L_q(B)} \|w^{-1}\|_{L_{p'}(B)} \geq 1. \quad (2.3)$$

If $p = 1$, w is in $A_{1,q}$ with $1 < q < \infty$ if

$$\begin{aligned} [w]_{A_{1,q}} &:= \sup_B [w]_{A_{1,q}(B)} \\ &= \sup_B \left(\frac{1}{|B|} \int_B w(x)^q dx \right)^{1/q} \left(\operatorname{ess\,sup}_{x \in B} \frac{1}{w(x)} \right) < \infty, \end{aligned}$$

where the sup is taken with respect to all balls B .

We define the generalized weighed Morrey spaces as follows.

Definition 2.1 Let $1 \leq p < \infty$, φ be a positive measurable function on $\mathbb{R}^n \times (0, \infty)$ and w be non-negative measurable function on \mathbb{R}^n . We denote by $M_{p,\varphi}(w)$ the generalized weighted Morrey space, the space of all functions $f \in L_{p,w}^{\text{loc}}(\mathbb{R}^n)$ with finite norm

$$\|f\|_{M_{p,\varphi}(w)} = \sup_{x \in \mathbb{R}^n, r > 0} \varphi(x, r)^{-1} w(B(x, r))^{-\frac{1}{p}} \|f\|_{L_{p,w}(B(x, r))},$$

where $L_{p,w}(B(x, r))$ denotes the weighted L_p -space of measurable functions f for which

$$\|f\|_{L_{p,w}(B(x, r))} \equiv \|f \chi_{B(x, r)}\|_{L_{p,w}(\mathbb{R}^n)} = \left(\int_{B(x, r)} |f(y)|^p w(y) dy \right)^{\frac{1}{p}}.$$

Furthermore, by $WM_{p,\varphi}(w)$ we denote the weak generalized weighted Morrey space of all functions $f \in WL_{p,w}^{\text{loc}}(\mathbb{R}^n)$ for which

$$\|f\|_{WM_{p,\varphi}(w)} = \sup_{x \in \mathbb{R}^n, r > 0} \varphi(x, r)^{-1} w(B(x, r))^{-\frac{1}{p}} \|f\|_{WL_{p,w}(B(x, r))} < \infty,$$

where $WL_{p,w}(B(x, r))$ denotes the weak $L_{p,w}$ -space of measurable functions f for which

$$\|f\|_{WL_{p,w}(B(x, r))} \equiv \|f \chi_{B(x, r)}\|_{WL_{p,w}(\mathbb{R}^n)} = \sup_{t > 0} t \left(\int_{\{y \in B(x, r) : |f(y)| > t\}} w(y) dy \right)^{\frac{1}{p}}.$$

Remark 2.1 (1) If $w \equiv 1$, then $M_{p,\varphi}(1) = M_{p,\varphi}$ is the generalized Morrey space.

(2) If $\varphi(x, r) \equiv w(B(x, r))^{\frac{\kappa-1}{p}}$, then $M_{p,\varphi}(w) = L_{p,\kappa}(w)$ is the weighted Morrey space.

(3) If $\varphi(x, r) \equiv v(B(x, r))^{\frac{\kappa}{p}} w(B(x, r))^{-\frac{1}{p}}$, then $M_{p,\varphi}(w) = L_{p,\kappa}(v, w)$ is the two weighted Morrey space.

(4) If $w \equiv 1$ and $\varphi(x, r) = r^{\frac{\lambda-n}{p}}$ with $0 < \lambda < n$, then $M_{p,\varphi}(w) = L_{p,\lambda}(\mathbb{R}^n)$ is the classical Morrey space and $WM_{p,\varphi}(w) = WL_{p,\lambda}(\mathbb{R}^n)$ is the weak Morrey space.

(5) If $\varphi(x, r) \equiv w(B(x, r))^{-\frac{1}{p}}$, then $M_{p,\varphi}(w) = L_{p,w}(\mathbb{R}^n)$ is the weighted Lebesgue space.

In the sequel $\mathfrak{M}(\mathbb{R}_+)$, $\mathfrak{M}^+(\mathbb{R}_+)$ and $\mathfrak{M}^+(\mathbb{R}_+; \uparrow)$ stand for the set of Lebesgue-measurable functions on \mathbb{R}_+ , and its subspaces of nonnegative and nonnegative non-decreasing functions, respectively. We also denote

$$\mathbb{A} = \{\varphi \in \mathfrak{M}^+(\mathbb{R}_+; \uparrow) : \lim_{t \rightarrow 0^+} \varphi = 0\}.$$

Let u be a continuous and non-negative function on \mathbb{R}_+ . We define the supremal operator \bar{S}_u by

$$(\bar{S}_u g)(t) := \|ug\|_{L_\infty(t, \infty)}, \quad t \in (0, \infty),$$

The following theorem was proved in [4].

Theorem 2.1 [4] *Suppose that v_1 and v_2 are nonnegative measurable functions such that $0 < \|v_1\|_{L_\infty(0, \cdot)} < \infty$ for every $t > 0$. Let u be a continuous nonnegative function on \mathbb{R} . Then the operator \bar{S}_u is bounded from $L_{\infty, v_1}(\mathbb{R}_+)$ to $L_{\infty, v_2}(\mathbb{R}_+)$ on the cone \mathbb{A} if and only if*

$$\left\| v_2 \bar{S}_u(\|v_1\|_{L_\infty(\cdot, \infty)}^{-1}) \right\|_{L_\infty(\mathbb{R}_+)} < \infty.$$

3 Fractional maximal operator in the spaces $M_{p,\varphi}(\mathbb{R}^n, w)$

The following Guliyev weighted local estimates are valid (see [13]).

Theorem 3.1 *Let $1 \leq p < q < \infty$, $0 \leq \alpha < \frac{n}{p}$, $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$, and $\omega \in A_{p,q}(\mathbb{R}^n)$. Then, for $p > 1$ the inequality*

$$\|M_\alpha f\|_{L_{q,w^q}(B(x,r))} \lesssim (w^q(B(x,r)))^{\frac{1}{q}} \sup_{t \geq r} \|f\|_{L_{p,w^p}(B(x,t))} (w^q(B(x,t)))^{-\frac{1}{q}} \quad (3.1)$$

holds for any ball $B(x, r)$ and for all $f \in L_{p,w}^{loc}(\mathbb{R}^n)$.

Moreover, for $p = 1$ the inequality

$$\|M_\alpha f\|_{WL_{q,w^q}(B(x,r))} \lesssim (w^q(B(x,r)))^{\frac{1}{q}} \sup_{t \geq r} \|f\|_{L_{1,w}(B(x,t))} (w^q(B(x,t)))^{-\frac{1}{q}} \quad (3.2)$$

holds for any ball $B(x, r)$ and for all $f \in L_{1,w}^{loc}(\mathbb{R}^n)$.

Proof. Let $1 < p < q < \infty$, $0 < \alpha < \frac{n}{p}$, $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$, and $w \in A_{p,q}(\mathbb{R}^n)$. For arbitrary $x \in \mathbb{R}^n$ and $r > 0$, set $B = B(x, r)$, $2B = B(x, 2r)$.

We present f as

$$f = f_1 + f_2, \quad f_1(y) = f(y) \chi_{2B}(y), \quad f_2(y) = f(y) \chi_{\complement_{(2B)}}(y),$$

and have

$$\|M_\alpha f\|_{L_{q,w^q}(B)} \leq \|M_\alpha f_1\|_{L_{q,w^q}(B)} + \|M_\alpha f_2\|_{L_{q,w^q}(B)}.$$

Since $f_1 \in L_{p,w^p}(\mathbb{R}^n)$, $M_\alpha f_1 \in L_{q,w^q}(\mathbb{R}^n)$ and from the boundedness of M_α from $L_{p,w^p}(\mathbb{R}^n)$ to $L_{q,w^q}(\mathbb{R}^n)$ (see [28]) it follows that:

$$\|M_\alpha f_1\|_{L_{q,w^q}(B)} \leq \|M_\alpha f_1\|_{L_{q,w^q}} \lesssim \|f_1\|_{L_{p,w^p}} = \|f\|_{L_{p,w^p}(2B)}. \quad (3.3)$$

From (3.3) we obtain

$$\|M_\alpha f_1\|_{L_{q,w^q}(B)} \lesssim (w^q(B(x,r)))^{\frac{1}{q}} \sup_{t \geq r} \|f\|_{L_{p,w^p}(B(x,t))} (w^q(B(x,t)))^{-\frac{1}{q}}. \quad (3.4)$$

Let z be an arbitrary point in $B \equiv B(x, r)$. If $B(z, t) \cap \complement_{B(x, 2r)} \neq \emptyset$, then $t > r$. Indeed, if $y \in B(z, t) \cap \complement_{B(x, 2r)}$, then we get $t > |y - z| \geq |x - y| - |x - z| > 2r - r = r$.

On the other hand, $B(z, t) \cap \complement_{B(x, 2r)} \subset B(x, 2t)$. Indeed, if $y \in B(z, t) \cap \complement_{B(x, 2r)}$, then we get $|x - y| \leq |y - z| + |x - z| < t + r < 2t$. Hence, for all $z \in B$

$$\begin{aligned} M_\alpha f_2(z) &= \sup_{t > 0} |B(z, t)|^{-1 + \frac{\alpha}{n}} \int_{B(z, t)} |f_2(y)| dy \\ &\leq \sup_{t > r} |B(x, 2t)|^{-1 + \frac{\alpha}{n}} \int_{B(z, t) \cap \complement_{B(x, 2r)}} |f(y)| dy \\ &\leq \sup_{t > r} |B(x, 2t)|^{-1 + \frac{\alpha}{n}} \int_{B(x, 2t)} |f(y)| dy \\ &= \sup_{t > 2r} |B(x, t)|^{-1 + \frac{\alpha}{n}} \int_{B(x, t)} |f(y)| dy. \end{aligned}$$

By applying Hölder's inequality, for all $z \in B$ we get

$$\begin{aligned} M_\alpha f_2(z) &\leq \sup_{t > 2r} |B(x, t)|^{-1 + \frac{\alpha}{n}} \int_{B(x, t)} |f(y)| dy \\ &\lesssim \sup_{t > r} |B(x, t)|^{-1 + \frac{\alpha}{n}} \|f\|_{L_{p,w^p}(B(x,t))} \|w^{-1}\|_{L_{p'}(B(x,t))} \\ &\lesssim \sup_{t > r} \|f\|_{L_{p,w^p}(B(x,t))} (w^q(B(x,t)))^{-\frac{1}{q}}. \end{aligned}$$

Thus, the function $M_\alpha f_2(z)$, with fixed x and r , is dominated by the expression not depending on z . Then

$$\|M_\alpha f_2\|_{L_{q,w^q}(B(x,r))} \lesssim (w^q(B(x,r)))^{\frac{1}{q}} \sup_{t > r} \|f\|_{L_{p,w^p}(B(x,t))} (w^q(B(x,t)))^{-\frac{1}{q}}. \quad (3.5)$$

We then obtain (3.1) from (3.4) and (3.5).

Let $p = 1$, and $w \in A_{1,q}(\mathbb{R}^n)$.

Then,

$$\|M_\alpha f\|_{WL_{q,w^q}(B)} \leq \|M_\alpha f_1\|_{WL_{q,w^q}(B)} + \|M_\alpha f_2\|_{WL_{q,w^q}(B)}.$$

Since $f_1 \in L_{1,w}(\mathbb{R}^n)$, $M_\alpha f_1 \in WL_{q,w^q}(\mathbb{R}^n)$ and from the boundedness of M_α from $L_{1,w}(\mathbb{R}^n)$ to $WL_{q,w^q}(\mathbb{R}^n)$ (see [28]) it follows that:

$$\|M_\alpha f_1\|_{WL_{q,w^q}(B)} \leq \|M_\alpha f_1\|_{WL_{q,w^q}} \lesssim \|f_1\|_{L_{1,w}} = \|f\|_{L_{1,w}(2B)}. \quad (3.6)$$

From (3.6) we obtain

$$\|M_\alpha f_1\|_{WL_{q,w^q}(B)} \lesssim (w^q(B(x,r)))^{\frac{1}{q}} \sup_{t>r} \|f\|_{L_{1,w}(B(x,t))} (w^q(B(x,t)))^{-\frac{1}{q}}. \quad (3.7)$$

On the other hand,

$$\begin{aligned} \|M_\alpha f_2\|_{WL_{q,w^q}(B(x,r))} &\leq \|M_\alpha f_2\|_{L_{q,w^q}(B(x,r))} \\ &\lesssim (w^q(B(x,r)))^{\frac{1}{q}} \sup_{t>r} \|f\|_{L_{1,w}(B(x,t))} (w^q(B(x,t)))^{-\frac{1}{q}}. \end{aligned} \quad (3.8)$$

We then obtain (3.2) from (3.7) and (3.8).

For the operator M_α the following Spanne-Guliyev type result on the space $M_{p,\varphi}(w)$ is valid.

Theorem 3.2 *Let $1 \leq p < q < \infty$, $0 < \alpha < \frac{n}{p}$, $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$, $w \in A_{p,q}(\mathbb{R}^n)$, and (φ_1, φ_2) satisfy the condition*

$$\sup_{t>r} \frac{\operatorname{ess\,inf}_{t<s<\infty} \varphi_1(x,s) (w^p B((x,s)))^{1/p}}{(w^q(B(x,t)))^{1/q}} \leq C \varphi_2(x,r), \quad (3.9)$$

where C does not depend on u and r . Then the operator M_α is bounded from $M_{p,\varphi_1}(w^p)$ to $M_{q,\varphi_2}(w^q)$ for $p > 1$ and from $M_{1,\varphi_1}(w)$ to $WM_{q,\varphi_2}(w^q)$ for $p = 1$. Moreover, for $p > 1$

$$\|M_\alpha f\|_{M_{q,\varphi_2}(w^q)} \lesssim \|f\|_{M_{p,\varphi_1}(w^p)},$$

and for $p = 1$

$$\|M_\alpha f\|_{WM_{q,\varphi_2}(w^q)} \lesssim \|f\|_{M_{1,\varphi_1}(w)}.$$

Proof. For $p > 1$ from Theorem 2.1 and Theorem 3.1 we get

$$\begin{aligned} \|M_\alpha f\|_{M_{q,\varphi_2}(w^q)} &\lesssim \sup_{x \in \mathbb{R}^n, r > 0} \varphi_2(x,r)^{-1} \sup_{t>r} \|f\|_{L_{p,w^p} w(B(x,t))}^{-\frac{1}{q}} \\ &\lesssim \sup_{x \in \mathbb{R}^n, r > 0} \varphi_1(x,r)^{-1} \|w\|_{L_q(B(x,r))}^{-1} \|f\|_{L_{p,w^p}} \\ &\lesssim \|f\|_{M_{p,\varphi_1}(w^p)} \end{aligned}$$

and for $p = 1$

$$\begin{aligned} \|M_\alpha f\|_{WM_{q,\varphi_2}(w^q)} &\lesssim \sup_{x \in \mathbb{R}^n, r > 0} \varphi_2(x,r)^{-1} \sup_{t>r} \|f\|_{L_{1,w} w(B(x,t))}^{-\frac{1}{q}} \\ &\lesssim \sup_{x \in \mathbb{R}^n, r > 0} \varphi_1(x,r)^{-1} \|w\|_{L_q(B(x,r))}^{-1} \|f\|_{L_{1,w}} \\ &\lesssim \|f\|_{M_{p,\varphi_1}(w^p)}. \end{aligned}$$

Remark 3.1 Note that, in the case $w \equiv 1$, Theorems 3.1 and 3.2 were proved in [14], see also [1, 2, 18–20].

4 Commutators of fractional maximal operators in the spaces $M_{p,\varphi}(\mathbb{R}^n, w)$

We recall the definition of the space of $BMO(\mathbb{R}^n)$.

Definition 4.1 Suppose that $b \in L_1^{\text{loc}}(\mathbb{R}^n)$, and let

$$\|b\|_* = \sup_{x \in \mathbb{R}^n, r > 0} \frac{1}{|B(x, r)|} \int_{B(x, r)} |b(y) - b_{B(x, r)}| dy < \infty,$$

where

$$b_{B(x, r)} = \frac{1}{|B(x, r)|} \int_{B(x, r)} b(y) dy.$$

Define

$$BMO(\mathbb{R}^n) = \{b \in L_1^{\text{loc}}(\mathbb{R}^n) : \|b\|_* < \infty\}.$$

Modulo constants, the space $BMO(\mathbb{R}^n)$ is a Banach space with respect to the norm $\|\cdot\|_*$.

Lemma 4.1 [28] Let $w \in A_\infty$. Then the norm $\|\cdot\|_*$ is equivalent to the norm

$$\|b\|_{*,w} = \sup_{x \in \mathbb{R}^n, r > 0} \frac{1}{w(B(x, r))} \int_{B(x, r)} |b(y) - b_{B(x, r), w}| w(y) dy,$$

where

$$b_{B(x, r), w} = \frac{1}{w(B(x, r))} \int_{B(x, r)} b(y) w(y) dy.$$

The following lemma is proved in [13].

Lemma 4.2 1 Let $w \in A_\infty$ and $b \in BMO(\mathbb{R}^n)$. Let also $1 \leq p < \infty$, $x \in \mathbb{R}^n$, $k > 0$ and $r_1, r_2 > 0$. Then,

$$\left(\frac{1}{w(B(x, r_1))} \int_{B(x, r_1)} |b(y) - b_{B(x, r_2), w}|^{kp} w(y) dy \right)^{\frac{1}{p}} \leq C \left(1 + \left| \ln \frac{r_1}{r_2} \right| \right)^k \|b\|_*^k,$$

where $C > 0$ is independent of f , w , x , r_1 and r_2 .

2 Let $w \in A_p$ and $b \in BMO(\mathbb{R}^n)$. Let also $1 < p < \infty$, $x \in \mathbb{R}^n$, $k > 0$ and $r_1, r_2 > 0$. Then,

$$\begin{aligned} \left(\frac{1}{w^{1-p'}(B(x, r_1))} \int_{B(x, r_1)} |b(y) - b_{B(x, r_2), w}|^{kp'} w(y)^{1-p'} dy \right)^{\frac{1}{p'}} \\ \leq C \left(1 + \left| \ln \frac{r_1}{r_2} \right| \right)^k \|b\|_*^k, \end{aligned}$$

where $C > 0$ is independent of b , w , x , r_1 and r_2 .

Remark 4.1 (1) Let $b \in BMO(\mathbb{R}^n)$. Then

$$\|b\|_* \approx \sup_{x \in \mathbb{R}^n, r > 0} \left(\frac{1}{|B(x, r)|} \int_{B(x, r)} |b(y) - b_{B(x, r)}|^p dy \right)^{\frac{1}{p}} \quad (4.1)$$

for $1 < p < \infty$.

(2) Let $b \in BMO(\mathbb{R}^n)$. Then there is a constant $C > 0$ such that

$$|b_{B(x, r)} - b_{B(x, \tau)}| \leq C \|b\|_* \log \frac{\tau}{r} \quad \text{for } 0 < 2r < \tau, \quad (4.2)$$

where C is independent of f , x , r and τ .

The commutator generated by $b \in L_1^{loc}(\mathbb{R}^n)$ and the operator M_α is defined by

$$M_{b,\alpha}(f)(x) = \sup_{r>0} |B(x,r)|^{-1+\frac{\alpha}{Q}} \int_{B(x,r)} |b(x) - b(y)| |f(y)| dy. \quad (4.3)$$

For the k th-order fractional maximal commutator operator $M_{b,\alpha,k}$ (see [14])

$$M_{b,\alpha,k}(f)(x) = \sup_{r>0} |B(x,r)|^{-1+\frac{\alpha}{Q}} \int_{B(x,r)} |b(x) - b(y)|^k |f(y)| dy$$

the following Guliyev weighted local estimates are valid (see [13]).

Theorem 4.1 *Let $1 < p < q < \infty$, $0 < \alpha < \frac{Q}{p}$, $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{Q}$, $b \in BMO(\mathbb{R}^n)$, and $w \in A_{p,q}(\mathbb{R}^n)$. Then the inequality*

$$\begin{aligned} & \|M_{b,\alpha,k} f\|_{L_{q,w^q}(B(x,r))} \\ & \lesssim \|b\|_*^k w^q(B(x,r))^{\frac{1}{q}} \sup_{t>2r} \ln^k \left(e + \frac{t}{r} \right) w^q(B(x,t))^{-\frac{1}{q}} \|f\|_{L_{p,w^p}(B(x,t))} \end{aligned}$$

holds for any ball $B(x,r)$ and for all $f \in L_{p,w^p}^{loc}(\mathbb{R}^n)$.

Proof. Let $1 < p < q < \infty$, $0 < \alpha < \frac{Q}{p}$, $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{Q}$. We write f as

$$f = f_1 + f_2, \quad f_1(y) = f(y) \chi_{2B}(y), \quad f_2(y) = f(y) \chi_{\mathring{c}B}(y).$$

Hence,

$$\|M_{b,\alpha,k} f\|_{L_{q,w^q}(B)} \leq \|M_{b,\alpha,k} f_1\|_{L_{q,w^q}(B)} + \|M_{b,\alpha,k} f_2\|_{L_{q,w^q}(B)}$$

From the boundedness of $M_{b,\alpha,k}$ from $L_{p,w^p}(\mathbb{R}^n)$ to $L_{q,w^q}(\mathbb{R}^n)$ (see [3]) it follows that:

$$\|M_{b,\alpha,k} f_1\|_{L_{q,w^q}(B)} \leq \|M_{b,\alpha,k} f_1\|_{L_{q,w^q}} \lesssim \|b\|_*^k \|f_1\|_{L_{p,w^p}} = \|b\|_*^k \|f\|_{L_{p,w^p}(2B)}. \quad (4.4)$$

From (4.4) we obtain

$$\|M_{b,\alpha,k} f_1\|_{L_{q,w^q}(B)} \lesssim \|b\|_*^k (w^q(B(x,r)))^{\frac{1}{q}} \sup_{t \geq r} \|f\|_{L_{p,w^p}(B(x,t))} (w^q(B(x,t)))^{-\frac{1}{q}}. \quad (4.5)$$

Let z be an arbitrary point in $B \equiv B(x,r)$. If $B(z,t) \cap \mathring{c}B(x,2r) \neq \emptyset$, then $t > r$. Indeed, if $y \in B(z,t) \cap \mathring{c}B(x,2r)$, then we get $t > |y-z| \geq |x-y| - |x-z| > 2r - r = r$.

On the other hand, $B(z,t) \cap \mathring{c}B(x,2r) \subset B(x,2t)$. Indeed, if $y \in B(z,t) \cap \mathring{c}B(x,2r)$, then we get $|x-y| \leq |y-z| + |x-z| < t + r < 2t$. Hence, for all $z \in B$

$$\begin{aligned} M_{b,\alpha,k} f_2(z) &= \sup_{t>0} |B(z,t)|^{-1+\frac{\alpha}{n}} \int_{B(z,t)} |b(y) - b(z)|^k |f_2(y)| dy \\ &= \sup_{t>0} |B(z,t)|^{-1+\frac{\alpha}{n}} \int_{B(z,t) \cap \mathring{c}B(x,2r)} |b(y) - b(z)|^k |f(y)| dy \\ &\lesssim \sup_{t>r} |B(x,2t)|^{-1+\frac{\alpha}{n}} \int_{B(x,2t)} |b(y) - b(z)|^k |f(y)| dy \\ &= \sup_{t>2r} |B(x,2t)|^{-1+\frac{\alpha}{n}} \int_{B(x,t)} |b(y) - b(z)|^k |f(y)| dy. \end{aligned}$$

Therefore, for all $z \in B$ we have

$$M_{b,\alpha,k}f_2(z) \lesssim \sup_{t>2r} |B(x, 2t)|^{-1+\frac{\alpha}{n}} \int_{B(x,t)} |b(y) - b(z)|^k |f(y)| dy.$$

Thus, the function $M_{\alpha}f_2(z)$, with fixed u and r , is dominated by the expression not depending on z . Then

$$\begin{aligned} & \|M_{b,\alpha,k}f_2\|_{L_{q,w^q}(B)} \\ & \lesssim \left(\int_B \left(\sup_{t>2r} |B(x, t)|^{-1+\frac{\alpha}{n}} \int_{B(x,t)} |b(y) - b(z)|^k |f(y)| dy \right)^q w^q(z) dz \right)^{\frac{1}{n}} \\ & \lesssim \left(\int_B \left(\sup_{t>2r} |B(x, t)|^{-1+\frac{\alpha}{n}} \int_{B(x,t)} |b(y) - b_{B(x,r),w}|^k |f(y)| dy \right)^q w^q(z) dz \right)^{\frac{1}{n}} \\ & + \left(\int_B \left(\sup_{t>2r} |B(x, t)|^{-1+\frac{\alpha}{n}} \int_{B(x,t)} |b(z) - b_{B(x,r),w}|^k |f(y)| dy \right)^q w^q(z) dz \right)^{\frac{1}{n}} \\ & = J_1 + J_2. \end{aligned}$$

Let us estimate J_1 . Applying Hölder's inequality and by Lemma 4.2 we get

$$\begin{aligned} J_1 & = w^q(B(x, r))^{\frac{1}{q}} \sup_{t>2r} |B(x, t)|^{-1+\frac{\alpha}{n}} \int_{B(x,t)} |b(y) - b_{B(x,r),w}|^k |f(y)| dy \\ & \approx w^q(B(x, r))^{\frac{1}{q}} \sup_{t>2r} t^{\alpha-Q} \int_{B(x,t)} |b(y) - b_{B(x,r),w}|^k |f(y)| dy \\ & \leq (w^q(B(x, r)))^{\frac{1}{q}} \sup_{t>2r} t^{\alpha-Q} \left(\int_{B(x,t)} |b(y) - b_{B(x,r),w}|^{kp'} w(y)^{-p'} dy \right)^{\frac{1}{p'}} \|f\|_{L_{p,w^p}(B(x,t))} \\ & \lesssim \|b\|_*^k w^q(B(x, r))^{\frac{1}{q}} \sup_{t>2r} t^{\alpha-Q} \left(1 + \ln \frac{t}{r} \right)^k \|w^{-1}\|_{L_{p'}(B(x,t))} \|f\|_{L_{p,w^p}(B(x,t))} \\ & \lesssim \|b\|_*^k w^q(B(x, r))^{\frac{1}{n}} \sup_{t>2r} t^{\alpha-Q} \left(1 + \ln \frac{t}{r} \right)^k (w^q(B(x, t)))^{-\frac{1}{n}} t^{\frac{n}{p} + \frac{n}{p'}} \|f\|_{L_{p,w^p}(B(x,t))} \\ & = \|b\|_*^k w^q(B(x, r))^{\frac{1}{q}} \sup_{t>2r} \ln^k \left(e + \frac{t}{r} \right) w^q(B(x, t))^{-\frac{1}{q}} \|f\|_{L_{p,w^p}(B(x,t))}. \end{aligned}$$

In order to estimate I_2 we get

$$\begin{aligned} J_2 & = \left(\int_B \left(\sup_{t>2r} |B(x, t)|^{-1+\frac{\alpha}{n}} \int_{B(x,t)} |b(z) - b_{B(x,r),w}|^k |f(y)| dy \right)^q w^q(z) dz \right)^{\frac{1}{q}} \\ & \approx \left(\int_B |b(z) - b_{B(x,r),w}|^{kq} w^q(z) dz \right)^{\frac{1}{q}} \sup_{t>2r} t^{\alpha-Q} \int_{B(x,t)} |f(y)| dy. \end{aligned}$$

According to the first part of Lemma 4.2, we get

$$\begin{aligned} J_2 & \lesssim \|b\|_*^k \left(1 + \ln \frac{r}{r} \right)^k w^q(B(x, r))^{\frac{1}{q}} \sup_{t>2r} t^{\alpha-Q} \int_{B(x,t)} |f(y)| dy \\ & \leq \|b\|_*^k w^q(B(x, r))^{\frac{1}{q}} \sup_{t>2r} t^{\alpha-Q} \|f\|_{L_{p,w^p}(B(x,t))} \|w^{-1}\|_{L_{p'}(B(x,t))} \\ & = \|b\|_*^k w^q(B(x, r))^{\frac{1}{q}} \sup_{t>2r} \ln^k \left(e + \frac{t}{r} \right) w^q(B(x, t))^{-\frac{1}{q}} \|f\|_{L_{p,w^p}(B(x,t))}. \end{aligned}$$

Summing up J_1 and J_2 , for all $p \in (1, \infty)$ we get

$$\begin{aligned} \|M_{b,\alpha,k}f_2\|_{L_{q,w^q}(B)} &\lesssim \|b\|_*^k w^q(B(x,r))^{\frac{1}{q}} \\ &\quad \times \sup_{t>2r} \ln^k \left(e + \frac{t}{r} \right) w^q(B(x,t))^{-\frac{1}{q}} \|f\|_{L_{p,w^p}(B(x,t))}. \end{aligned} \quad (4.6)$$

Finally, from (4.5) and (4.6) we get

$$\begin{aligned} \|M_{b,\alpha,k}f\|_{L_{q,w^q}(B)} &\lesssim \|b\|_*^k (w^q(B(x,r)))^{\frac{1}{q}} \sup_{t \geq r} \|f\|_{L_{p,w^p}(B(x,t))} (w^q(B(x,t)))^{-\frac{1}{q}} \\ &\quad + \|b\|_*^k w^q(B(x,r))^{\frac{1}{q}} \sup_{t>2r} \ln^k \left(e + \frac{t}{r} \right) w^q(B(x,t))^{-\frac{1}{q}} \|f\|_{L_{p,w^p}(B(x,t))} \\ &\lesssim \|b\|_*^k w^q(B(x,r))^{\frac{1}{q}} \sup_{t>2r} \ln^k \left(e + \frac{t}{r} \right) w^q(B(x,t))^{-\frac{1}{q}} \|f\|_{L_{p,w^p}(B(x,t))}. \end{aligned}$$

For the operator $M_{b,\alpha,k}$ the following Spanne-Guliyev type result on the space $M_{p,\varphi}(w)$ is valid.

Theorem 4.2 *Let $1 < p < q < \infty$, $0 < \alpha < \frac{n}{p}$, $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$, $w \in A_{p,q}(\mathbb{R}^n)$, $b \in BMO(\mathbb{R}^n)$ and (φ_1, φ_2) satisfy the condition*

$$\sup_{t>r} \ln^k \left(e + \frac{t}{r} \right) \frac{\operatorname{ess\,inf}_{t<s<\infty} \varphi_1(x,s) (w^p B((x,s)))^{1/p}}{(w^q(B(x,t)))^{1/q}} \leq C \varphi_2(x,r), \quad (4.7)$$

where C does not depend on u and r . Then the operator $M_{b,\alpha,k}$ is bounded from $M_{p,\varphi_1}(w^p)$ to $M_{q,\varphi_2}(w^q)$. Moreover,

$$\|M_{b,\alpha,k}f\|_{M_{q,\varphi_2}(w^q)} \lesssim \|b\|_*^k \|f\|_{M_{p,\varphi_1}(w^p)}.$$

Proof. Using the Theorem 2.1 and the Theorem 4.1 we have

$$\begin{aligned} \|M_{b,\alpha,k}f\|_{M_{q,\varphi_2}(w^q)} &= \sup_{x \in \mathbb{R}^n, r>0} \varphi_2(x,r)^{-1} w^q(B(x,r))^{\frac{1}{q}} \|M_{b,\alpha,k}f\|_{L_{q,w^q}B(x,r)} \\ &\lesssim \|b\|_*^k \sup_{x \in \mathbb{R}^n, r>0} \varphi_2(x,r)^{-1} \sup_{t>2r} \ln^k \left(e + \frac{t}{r} \right) w^q(B(x,t))^{-\frac{1}{q}} \|f\|_{L_{p,w^p}(B(x,t))} \\ &\lesssim \|b\|_*^k \sup_{x \in \mathbb{R}^n, r>0} \varphi_1(x,r)^{-1} (w^p(B(x,r)))^{-\frac{1}{p}} \|f\|_{L_{p,w^p}(B(x,r))} \\ &= \|b\|_*^k \|f\|_{M_{p,\varphi_1}(w^p)}. \end{aligned}$$

Corollary 4.1 *Let $1 < p < q < \infty$, $0 < \alpha < \frac{n}{p}$, $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$, $w \in A_{p,q}(\mathbb{R}^n)$, $b \in BMO(\mathbb{R}^n)$, $b^- \in L^\infty(\mathbb{R}^n)$ and (φ_1, φ_2) satisfy the condition (4.7). Then the operator $[b, M_\alpha]^k$ is bounded from $M_{p,\varphi_1}(w^p)$ to $M_{q,\varphi_2}(w^q)$.*

Remark 4.2 Note that, in the case $w \equiv 1$, Theorems 4.1 and 4.2 were proved in [14], see also [19, 21, 23].

Acknowledgements. The author would like to express their gratitude to the referees for his very valuable comments and suggestions.

References

1. Akbulut, A., Guliyev, V.S., Mustafayev, R.: *On the boundedness of the maximal operator and singular integral operators in generalized Morrey spaces*, Math. Bohem. **137** (1), 27-43 (2012).
2. Akbulut, A., Hamzayev, V.H., Safarov, Z.V.: *Rough fractional multilinear integral operators on generalized weighted Morrey spaces*, Azerb. J. Math. **6** (2), 128-142 (2016).
3. Bernardis, A., Hartzstein, S., Pradolini, G.: *Weighted inequalities for commutators of fractional integrals on spaces of homogeneous type*, J. Math. Anal. Appl. **322**, 825-846 (2006).
4. Burenkov, V., Gogatishvili, A., Guliyev, V., Mustafayev, R.: *Boundedness of the fractional maximal operator in local Morrey-type spaces*, Complex Var Elliptic Equ. **55** (8-10), 739-758 (2010).
5. Chiarenza, F., Frasca, M.: *Morrey spaces and Hardy-Littlewood maximal function*, Rend Mat. **7**, 273-279 (1987).
6. Chiarenza, F., Frasca, M., Longo, P.: *$W^{2,p}$ -solvability of Dirichlet problem for non-divergence elliptic equations with VMO coefficients*, Trans. Amer. Math. Soc. **336**, 841-853 (1993).
7. Di Fazio, G., Ragusa, M.A.: *Interior estimates in Morrey spaces for strong solutions to nondivergence form equations with discontinuous coefficients*, J. Funct. Anal. **112**, 241-256 (1993).
8. Fan, D., Lu, S., Yang, D.: *Boundedness of operators in Morrey spaces on homogeneous spaces and its applications*, Acta Math. Sinica (N. S.) **14**, suppl. 625-634 (1998).
9. Eroglu, A., Omarova, M.N., Muradova, Sh.A.: *Elliptic equations with measurable coefficients in generalized weighted Morrey spaces*, Proc. Inst. Math. Mech. Natl. Acad. Sci. Azerb. **43** (2), 197-213 (2017).
10. Deringoz, F.: *Parametric Marcinkiewicz integral operator and its higher order commutators on generalized weighted Morrey spaces*, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci. **37** (4), Mathematics, 24-32 (2017).
11. Giaquinta, M.: *Multiple integrals in the calculus of variations and nonlinear elliptic systems*, Princeton Univ. Press, Princeton, NJ, 1983.
12. Guliyev, V.S.: *Boundedness of the maximal, potential and singular operators in the generalized Morrey spaces*, J. Inequal. Appl. 2009, Art. ID 503948, 20 pp.
13. Guliyev, V.S.: *Generalized weighted Morrey spaces and higher order commutators of sublinear operators*, Eurasian Math. J. **3** (3), 33-61 (2012).
14. Guliyev, V.S., Akbulut, A., Mammadov, Y.Y.: *Boundedness of fractional maximal operator and their higher order commutators in generalized Morrey spaces on Carnot groups*, Acta Math. Sci. Ser. B Engl. Ed. **33** (5), 1329-1346 (2013).
15. Guliyev, V.S., Karaman, T., Mustafayev, R.Ch., Serbetci, A.: *Commutators of sublinear operators generated by Calderón-Zygmund operator on generalized weighted Morrey spaces*, Czechoslovak Math. J. **64(139)** (2), 365-386 (2014).
16. Guliyev, V.S., Ismayilova, A.F.: *Marcinkiewicz integral and their commutators on generalized weighted Morrey spaces*, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci. **34** (4), Mathematics and Mechanics, 53-64 (2014).
17. Guliyev, V.S., Omarova, M.N.: *Multilinear commutators of vector-valued intrinsic square functions on vector-valued generalized weighted Morrey spaces*, J. Inequal. Appl. 2014, 2014:258.
18. Guliyev, V.S., Balakishiyev, A.S.: *Parabolic fractional maximal and integral operators with rough kernels in parabolic generalized Morrey spaces*, J. Math. Inequal. **9** (1), 257-276 (2015).
19. Guliyev, V.S., Deringoz, F.: *Boundedness of fractional maximal operator and its commutators on generalized Orlicz-Morrey spaces*, Complex Anal. Oper. Theory **9** (6),

- 1249-1267 (2015).
20. Guliyev, V.S., Omarova, M.N.: *Multilinear singular and fractional integral operators on generalized weighted Morrey spaces*, Azerb. J. Math. **5** (1), 104-132 (2015).
 21. Hamzayev, V.H.: *Sublinear operators with rough kernel generated by Calderon-Zygmund operators and their commutators on generalized weighted Morrey spaces*, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci. **38** (1), Mathematics, 79-94 (2018).
 22. Guliyev, V.S., Omarova, M.N., Softova L.: *The Dirichlet problem in a class of generalized weighted Morrey spaces*, Proc. Inst. Math. Mech. Natl. Acad. Sci. Azerb., **45** (2), 1-19 (2019).
 23. Ismayilova, A.F.: *Commutators of the Marcinkiewicz integral on generalized weighted Morrey spaces*, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci. **38** (4), Mathematics, 79-92 (2018).
 24. Komori, Y., Shirai, S.: *Weighted Morrey spaces and a singular integral operator*, Math. Nachr. **282** (2), 219-231 (2009).
 25. Kufner, A., John, O., Fučík, S.: *Function Spaces*. Noordhoff International Publishing: Leyden, Publishing House Czechoslovak Academy of Sciences: Prague, 1977.
 26. Morrey, C.B.: *On the solutions of quasi-linear elliptic partial differential equations*, Trans. Amer. Math. Soc. **43**, 126-166 (1938).
 27. Muckenhoupt, B.: *Weighted norm inequalities for the Hardy maximal function*, Trans. Amer. Math. Soc. **165**, 207-226 (1972).
 28. Muckenhoupt, B., Wheeden, R.: *Weighted norm inequalities for fractional integrals*, Trans. Amer. Math. Soc. **192**, 261-274 (1974).
 29. Zhang, P., Wu, J.: *Commutators of the fractional maximal function on variable exponent Lebesgue spaces*. Czechoslovak Math. J. **64** (139) (1), 183-197 (2014).