

Absence of global solutions of a semilinear biharmonic equation with a singular potential in exterior domains

Shirmail G. Bagirov* · Abdurrahim F. Guliyev

Received: 03.11.2021 / Revised: 13.04.2022 / Accepted: 16.05.2022

Abstract. In the domain $\{x \in \mathbb{R}^n : |x| > R\} \times \{y \in \mathbb{R}^m : |y| > R\}$ the problem of the absence of global solutions of a semilinear biharmonic equation $\Delta_x (|x|^\alpha \Delta_x u) + \Delta_y^2 u - \frac{C_1}{|x|^{4-\alpha}} u - \frac{C_2}{|y|^4} u - |x|^{\sigma_1} |y|^{\sigma_2} |u|^q = 0$ with are conditions $u \geq 0, \Delta u \leq 0$ on $\{|x| = R\} \times \{|y| > R\} \cup \{|x| > R\} \times \{|y| = R\}$ is investigated, where $q > 1, \alpha < 4, n > 4 - \alpha, m > 4, 0 \leq C_1 < \left(\frac{(n-\alpha)(n+\alpha-4)}{4}\right)^2, 0 \leq C_2 < \left(\frac{m(m-4)}{4}\right)^2$. A sufficient condition for the absence of global solutions is obtained. The proof is based on the test function method.

Keywords. Semilinear biharmonic equation, global solution, singular potential, critical exponent, test function method.

Mathematics Subject Classification (2010): 35A01, 35B33, 35J61, 35J30

1 Introduction

We induuce the following designations:

$$\begin{aligned}x &= (x_1, \dots, x_n) \in \mathbb{R}^n, \quad y = (y_1, \dots, y_m) \in \mathbb{R}^m, \quad |x| = \sqrt{\sum_{i=1}^n x_i^2}, \quad |y| = \sqrt{\sum_{i=1}^m y_i^2}, \\B_x(r) &= \{x \in \mathbb{R}^n : |x| < r\}, \quad B_y(r) = \{y \in \mathbb{R}^m : |y| < r\}, \\S_x(r) &= \{x \in \mathbb{R}^n : |x| = r\}, \quad S_y(r) = \{y \in \mathbb{R}^m : |y| = r\}, \\B(r) &= B_x(r) \times B_y(r), \quad B_x(r_1, r_2) = \{x \in \mathbb{R}^n : r_1 < |x| < r_2\}, \\B_y(r_1, r_2) &= \{y \in \mathbb{R}^m : r_1 < |y| < r_2\}, \\B(r_1, r_2) &= B_x(r_1, r_2) \times B_y(r_1, r_2), \quad B'_x(R) = \mathbb{R}^n \setminus B_x(R), \quad B'_y(R) = \mathbb{R}^m \setminus B_y(R), \\B'(R) &= B'_x(R) \times B'_y(R), \quad \partial B'(R) = S_x(R) \times B'_y(R) \cup B'_x(R) \times S_y(R),\end{aligned}$$

* Corresponding author

Sh.G. Bagirov,
Baku State University, Baku, Azerbaijan
Institute of Mathematics and Mechanics of NAS of Azerbaijan Baku, Azerbaijan
E-mail: sh_bagirov@yahoo.com

A.F. Guliyev
Institute of Mathematics and Mechanics of NAS of Azerbaijan Baku, Azerbaijan
E-mail: abdurrahim_57@yahoo.com

$$B'_x(1) = B'_x, B'_y(1) = B'_y, B'(1) = B'.$$

In $B'(R)$ we consider the equation

$$\Delta_x(|x|^\alpha \Delta_x u) + \Delta_y^2 u - \frac{C_1}{|x|^{4-\alpha}} u - \frac{C_2}{|y|^4} u - |x|^{\sigma_1} |y|^{\sigma_2} |u|^q = 0, \quad (1.1)$$

where

$$\alpha < 4, \sigma_1, \sigma_2 \in R, q > 1, 0 \leq C_1 < \left(\frac{(n-\alpha)(n+\alpha-4)}{4} \right)^2, \\ 0 \leq C_2 < \left(\frac{m-(m-4)}{4} \right)^2, \Delta_x = \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2}, \Delta_y^2 = \Delta_y(\Delta_y), \Delta_y = \sum_{i=1}^m \frac{\partial^2}{\partial y_i^2}.$$

We will consider the existence of the solution of an equation satisfying the following conditions:

$$u \geq 0, \Delta u \leq 0 \text{ on } \partial B'(R). \quad (1.2)$$

We will understand the solution of problem (1.1),(1.2) in the classic sense, namely, the function $u(x, y) \in C^4(B'_R) \cup C^3(B'_R \cup \partial B'_R)$ satisfying equation (1.1) at each point of $B'(R)$ and condition (1.2) on $\partial B'(R)$ will be called global solutions of problem (1.1),(1.2).

The problem of non- existence of global solutions for a different class of differential equations and inequalities play an important role in theory and applications; therefore, constantly draw attention of mathematicians and there is a large number of works devoted to them. A survey of such results was given in the monograph [15]. It is well known such statements are a direct analogs of the Liouville's theorem from the theory of functions of a complex variable.

The issues considered in this paper were earlier studied in the works of Mitidieri and Pokhozhaev [15], Gidas and Spruck [9], Brezis, Dupaigne and Tesei [6], Bidaut-Veron and Pohozaev [5], Serrin and Zou [18], Serrin [19], Konkov [10,11,12], Bagirov [1,2], Bagirov, Guliyev [4], Mamedov [14]. Similar issues for weakly nonlinear equations with a biharmonic operator were considered by various authors as Laptev [13], Volodin [20,21], Xu [22], Ghergu and Taliaferro [8], Carriao, Demarque and Miyagaki [7], Yao, Wang, Shen [23], Bagirov [3].

In the paper [21] for $\alpha = 0, C_1 = C_2 = 0$ is considered in $B'(R)$ with different boundary conditions on S_R and critical exponents of the absence of the solutions of the considered problems were found. In the paper [3] for $\alpha = 0$ the problem (1.1), (1.2) is considered in $B'(R)$ and critical exponent of the absence global solutions are also found.

In this paper the problem (1.1), (1.2) is consider for $\alpha < 4, n > 4 - \alpha, m > 4, 0 \leq C_1 < \left(\frac{(n-\alpha)(n+\alpha-4)}{4} \right)^2, 0 \leq C_2 < \left(\frac{(m-4)}{4} \right)^2$ and exact estimates of the absence global solutions are obtained. The technique of the test functions developed by Mitidieri and Pohozhayev in the paper [15,16,17] is used.

2 Main result and its proof

We introduce the following designation:

$$D_1 = \sqrt{\left(\frac{(n-2)(\alpha-2)}{2} \right)^2 + C_1}, \quad D_2 = \sqrt{(m-2)^2 + C_2} \\ \alpha_{\pm} = \sqrt{\left(\frac{n-2}{2} \right)^2 + \left(\frac{\alpha-2}{2} \right)^2} \mp D_1 \quad \text{for } 2 \leq \alpha < 4,$$

$$\alpha_{\pm} = \sqrt{\left(\frac{n-2}{2}\right)^2 + \left(\frac{\alpha-2}{2}\right)^2} \pm D_1 \quad \text{for } \alpha < 2,$$

$$\beta_{\pm} = \sqrt{\left(\frac{m-2}{2}\right)^2 + 1} \pm \sqrt{D_2}, \quad \gamma = \frac{2\sqrt{D_1}}{|\alpha-2|} - \alpha_{+},$$

$$a = \frac{m+4}{2} + \beta_{-} - \sigma_2 \frac{1}{q-1} = \frac{m+4}{2} + \beta_{-} - \sigma_2(q'-1),$$

$$b = \frac{n+4-\alpha}{2} + \alpha_{-} - \sigma_1(q'-1),$$

where $\frac{1}{q} + \frac{1}{q'} = 1$.

Let us consider the following functions

$$\xi(x) = \xi(|x|) = \frac{1}{2}(1+\gamma)|x|^{-\frac{n+\alpha-4}{2}+\alpha_{-}} + \frac{1}{2}(1-\gamma)|x|^{-\frac{n+\alpha-4}{2}-\alpha_{-}} - |x|^{-\frac{n+\alpha-4}{2}-\alpha_{+}} \quad \text{for } \alpha \neq 2,$$

$$\xi(x) = \frac{\alpha_{-}}{\alpha_{+}} \left(|x|^{-\frac{n-2}{2}+\alpha_{-}} - |x|^{-\frac{n-2}{2}-\alpha_{-}} \right) + |x|^{-\frac{n-2}{2}+\alpha_{+}} - |x|^{-\frac{n-2}{2}-\alpha_{+}} \quad \text{for } \alpha = 2,$$

$$\eta(y) = \eta(|y|) = \frac{1}{2} \left(1 + \frac{\sqrt{D_2} - \beta_{+}}{\beta_{-}} \right) |y|^{-\frac{m-4}{2}+\beta_{-}} + \frac{1}{2} \left(1 - \frac{\sqrt{D_2} - \beta_{+}}{\beta_{-}} \right) |y|^{-\frac{m-4}{2}-\beta_{-}} - |y|^{-\frac{m-4}{2}-\beta_{+}}.$$

We can show that $\xi(x)$ is the solution of the equation

$$\Delta_x(|x|^{\alpha} \Delta_x u) - \frac{C_1}{|x|^{4-\alpha}} u = 0, \quad (2.1)$$

$\eta(y)$ is the solution of the equation

$$\Delta_y^2 u - \frac{C_2}{|y|^4} u = 0, \quad (2.2)$$

and

$$\xi(x)|_{|x|=1} = 0, \quad \frac{\partial \xi}{\partial r} \Big|_{|x|=1} > 0, \quad \Delta_x \xi|_{|x|=1} = 0, \quad \frac{\Delta_x \xi}{\partial r} \Big|_{|x|=1} \leq 0,$$

$$\eta(x)|_{|y|=1} = 0, \quad \frac{\partial \eta}{\partial r} \Big|_{|y|=1} > 0, \quad \Delta_y \eta|_{|y|=1} = 0, \quad \frac{\Delta_y \eta}{\partial r} \Big|_{|y|=1} \leq 0.$$

We consider the following functions $\varphi(x), \psi(y)$ such that $\varphi(x) \in C^{\infty}(\mathbb{R}^n)$, $\varphi(x) = 1$ for $|x| \leq \rho$, $\varphi(x) = 0$ for $|x| \geq 2\rho$, $\psi(y) \in C^{\infty}(\mathbb{R}^m)$, $\psi(y) = 1$ for $|y| \leq \rho^{\varepsilon}$, $\psi(y) = 0$ for $|y| \geq 2\rho^{\varepsilon}$.

The following theorem is the main result of this paper.

Theorem 2.1 Let $\alpha < 4, n > 4 - \alpha, m > 4, 0 \leq C_1 < \left(\frac{(n-\alpha)(n+\alpha-4)}{4}\right)^2, 0 \leq C_2 < \left(\frac{m(m-4)}{4}\right)^2$ and

$$a) \sigma_1 > 0, \sigma_2 > 0, 1 < q \leq \min \left\{ 1 + \frac{\sigma_1}{\frac{n-\alpha+4}{2} + \alpha_-}, 1 + \frac{\sigma_2}{\frac{m+4}{2} + \beta_-} \right\},$$

$$b) \sigma_1 > 0, \sigma_2 > -4, q > \max \left\{ 1, 1 + \frac{\sigma_2}{\frac{m+4}{2} + \beta_-} \right\} \text{ and } q < 1 + \frac{\sigma_1}{\frac{n-\alpha+4}{2} + \alpha_-},$$

$$q \leq 1 + \frac{\sigma_2 + 4}{\frac{m+4}{2} + \beta_- - 4} \text{ or } q = 1 + \frac{\sigma_1}{\frac{n-\alpha+4}{2} + \alpha_-}, \quad q < 1 + \frac{\sigma_2 + 4}{\frac{m+4}{2} + \beta_- - 4},$$

$$c) \sigma_1 > \alpha - 4, \sigma_2 > 0, \quad q > \max \left\{ 1, 1 + \frac{\sigma_1}{\frac{n-\alpha+4}{2} + \alpha_-} \right\} \text{ and } q < 1 + \frac{\sigma_2}{\frac{m+4}{2} + \beta_-},$$

$$q \leq 1 + \frac{\sigma_1 + 4 - \alpha}{\frac{n+\alpha-4}{2} + \alpha_-} \text{ or } q = 1 + \frac{\sigma_2}{\frac{m+4}{2} + \beta_-}, \quad q < 1 + \frac{\sigma_1 + 4 - \alpha}{\frac{n+\alpha-4}{2} + \alpha_-},$$

$$d) 4 - \alpha + \sigma_1 + \frac{4-\alpha}{4}\sigma_2 > 0,$$

$$\max \left\{ 1, 1 + \frac{\sigma_1}{\frac{n-\alpha+4}{2} + \alpha_-}, 1 + \frac{\sigma_2}{\frac{m+4}{2} + \beta_-} \right\} < q \leq 1 + \frac{4 - \alpha + \sigma_1 + \frac{4-\alpha}{4}\sigma_2}{\frac{n-\alpha+4}{2} + \alpha_- + \frac{4-\alpha}{4} \left(\frac{m-4}{2} + \beta_- \right)}.$$

Then if $u(x, y)$ a nontrivial solution of the problem (1.1), (1.2), then $u \equiv 0$.

Proof. For simplicity of the notation we take $R = 1$. We multiply equation (1.1) by the function $f(x, y) = \xi(x)\eta(y)\varphi(x)\psi(y)$ and integrate with respect to the domain B' . We get the following equality

$$\begin{aligned} Q &\equiv \iint_{B'} |x|^{\sigma_1} |y|^{\sigma_2} |u|^q f dx dy = \iint_{B'} \Delta_x (|x|^\alpha \Delta_x u) f dx dy \\ &+ \iint_{B'} \Delta_y^2 u f dx dy - \iint_{B'} \frac{C_1}{|x|^{4-\alpha}} u f dx dy - \iint_{B'} \frac{C_2}{|y|^4} u f dx dy. \end{aligned}$$

Integrating by parts, we get

$$\begin{aligned} Q &= \int \int_{S_x B'_y} \frac{\partial}{\partial \nu_x} (|x|^\alpha \Delta_x u) \xi \eta \varphi \psi d\sigma_x dy \\ &- \int \int_{B'} (\nabla_x (|x|^\alpha \Delta_x u), \nabla_x (\xi \varphi)) \eta \psi dx dy \\ &+ \int \int_{B'_x S_y} \frac{\partial}{\partial \nu_y} (\Delta_y u) \xi \eta \varphi \psi dx d\sigma_y \\ &- \int \int_{B'} (\nabla_y (\Delta_y u), \nabla_y (\eta \psi)) \xi \varphi dx dy \end{aligned}$$

$$\begin{aligned}
& - \iint_{B'} \frac{C_1}{|x|^{4-\alpha}} u \xi \eta \varphi \psi dx dy - \iint_{B'} \frac{C_2}{|y|^4} u \xi \eta \varphi \psi dx dy = \\
& = \Gamma_x^1 - \Gamma_x^2 + \Gamma_x^3 - \Gamma_x^4 + \Gamma_y^1 - \Gamma_y^2 + \Gamma_y^3 - \Gamma_y^4 + \\
& + \iint_{B'} u \Delta_x (|x|^\alpha \Delta_x (\xi \varphi)) \eta \psi dx dy + \iint_{B'} u \Delta_y^2 (\eta \psi) \xi \varphi dx dy, \quad (2.3)
\end{aligned}$$

where $\Gamma_x^i, \Gamma_y^i, i = 1, 4$ are appropriate surface integrals on $S_x \times B'_y, B'_x \times S_y$ and σ_x, σ_y are the elements of the surfaces S_x, S_y .

Using the properties of the functions $\xi(x), \eta(y), \varphi(x), \psi(y)$ we show that $(-1)^{i-1} \Gamma_x^i, (-1)^{i-1} \Gamma_y^i \leq 0, i = 1, 4$.

Obviously all surface integrals on $S_x(2\rho) \times B'_y$ and $B'_x \times S_y(2\rho^\varepsilon)$ are equal to zero.

$$\Gamma_x^1 = \int \int_{S_x B'_y} \frac{\partial}{\partial \nu_x} (|x|^\alpha \Delta_x u) \xi \eta \varphi \psi d\sigma_x dy = 0,$$

since $\xi|_{|x|=1} = 0$.

$$\begin{aligned}
\Gamma_x^2 & = \int \int_{S_x B'_y} |x|^\alpha \Delta_x u \frac{\partial(\xi \varphi)}{\partial \nu_x} \eta \psi d\sigma_x dy \\
& = \int \int_{S_x B'_y} |x|^\alpha \Delta_x u \left(\frac{\partial \xi}{\partial \nu_x} \varphi + \xi \frac{\partial \varphi}{\partial \nu_x} \right) \eta \psi d\sigma_x dy \\
& = - \int \int_{S_x B'_y} |x|^\alpha \Delta_x u \frac{\partial \xi}{\partial r} \eta \psi d\sigma_x dy \geq 0,
\end{aligned}$$

by write of the condition (1.2) and that on S_x $\xi = 0, \frac{\partial \xi}{\partial \nu_x} = -\frac{\partial \xi}{\partial r} \leq 0, \varphi = 1$.

$$\begin{aligned}
\Gamma_x^3 & = \int \int_{S_x B'_y} \frac{\partial u}{\partial \nu_x} |x|^\alpha \Delta_x (\xi \varphi) \eta \psi d\sigma_x dy \\
& = \int \int_{S_x B'_y} \frac{\partial u}{\partial \nu_x} |x|^\alpha (\Delta_x \xi \varphi + 2(\nabla_x \xi, \nabla_x \varphi) + \xi \Delta_x \varphi) \eta \psi d\sigma_x dy = 0,
\end{aligned}$$

since on S_x $\xi = 0, \Delta_x \xi = 0, \nabla_x \varphi = 0$.

$$\begin{aligned}
\Gamma_x^4 & = \int \int_{S_x B'_y} u \frac{\partial}{\partial \nu_x} (|x|^\alpha \Delta_x (\varphi \xi)) \eta \psi d\sigma_x dy \\
& = \int \int_{S_x B'_y} u [\alpha |x|^{\alpha-2} (\Delta_x \varphi \xi + 2(\nabla_x \varphi, \nabla_x \xi) + \varphi \Delta_x \xi)(x, \nu_x) \\
& + |x|^\alpha \frac{\partial}{\partial \nu_x} (\Delta_x \varphi \xi + 2(\nabla_x \varphi, \nabla_x \xi) + \varphi \Delta_x \xi)] \eta \psi d\sigma_x dy
\end{aligned}$$

$$= \int \int_{S_x B'_y} u \frac{\partial}{\partial \nu_x} (\Delta_x \xi) \eta \psi d\sigma_x dy \geq 0.$$

Here we use the fact that for $|x| = 1$ $u \geq 0$, the function φ and all its derivatives equal zero, $\Delta_x \xi = 0$ and $\frac{\partial}{\partial \nu_x} \Delta_x \xi \geq 0$.

It the same way, we can show that all $(-1)^{i-1} \Gamma_y^i \leq 0, i = 1, 4$.

As a result, from (2.3) we get the following inequality

$$\begin{aligned} Q &= \iint_{B'} u \Delta_x (|x|^\alpha (\Delta_x \varphi \xi + 2(\nabla_x \xi, \nabla_x \varphi) + \varphi \Delta_x \xi)) \eta \psi dx dy \\ &+ \iint_{B'} u \Delta_y (\Delta_y \eta \psi + 2(\nabla_y \eta, \nabla_y \psi) + \eta \Delta_y \psi) \xi \varphi dx dy \\ &- \iint_{B'} \frac{C_1}{|x|^{4-\alpha}} u \xi \eta \varphi \psi dx dy - \iint_{B'} \frac{C_2}{|y|^4} u \xi \eta \varphi \psi dx dy \\ &= \iint_{B'} u \left(\Delta_x (|x|^\alpha \Delta_x \xi) - \frac{C_1}{|x|^{4-\alpha}} \xi \right) \eta \varphi \psi dx dy + \iint_{B'} \left(\Delta_y^2 \eta - \frac{C_2}{|y|^4} \eta \right) \xi \varphi \psi dx dy \\ &+ \iint_{B'} u [2(\nabla_x (|x|^\alpha \Delta_x \xi), \nabla_x \varphi) + |x|^\alpha \Delta_x \xi \Delta_x \varphi + 2\Delta_x (|x|^\alpha (\nabla_x \xi, \nabla_x \varphi)) \\ &+ \Delta_x (|x|^\alpha \xi \Delta_x \varphi)] \eta \psi dx dy + \iint_{B'} u [2(\nabla_y (\Delta_y \eta), \nabla_y \psi) \\ &+ \Delta_y \eta \Delta_y \psi + 2\Delta_y (\nabla_y \eta, \nabla_y \psi) + \Delta_y (\eta \Delta_y \psi)] \xi \varphi dx dy \\ &= \iint_{B'} u \eta \psi G_1(\xi, \varphi) dx dy + \iint_{B'} u \xi \varphi G_2(\eta, \psi) dx dy, \end{aligned} \quad (2.4)$$

where $G_1(\xi, \varphi), G_2(\eta, \psi)$ are the expressions in the square brackets in the first and second integral. Here we take into account that $\xi(x)$ is the solution of the equation (2.1), $\eta(x)$ is the solution of equation (2.2). As a result, we have

$$\begin{aligned} Q &\leq \iint_{B'} u \eta \psi G_1(\xi, \varphi) dx dy + \iint_{B'} u \xi \varphi G_2(\eta, \psi) dx dy \\ &\leq \left(\iint_{B_x(\rho, 2\rho) \times B_y(1, 2\rho^\varepsilon)} |x|^{\sigma_1} |y|^{\sigma_2} |u|^q \xi \eta \varphi \psi dx dy \right)^{\frac{1}{q}} \\ &\times \left(\iint_{B_x(\rho, 2\rho) \times B_y(1, 2\rho^\varepsilon)} \frac{|G_1|^{q'} \eta \psi}{|x|^{\sigma_1(q'-1)} |y|^{\sigma_2(q'-1)} \xi^{q'-1} \varphi^{q'-1}} dx dy \right)^{\frac{1}{q'}} \end{aligned}$$

$$\begin{aligned}
& + \left(\iint_{B_x(1,2\rho) \times B_y(\rho,2\rho^\varepsilon)} |x|^{\sigma_1} |y|^{\sigma_2} |u|^q \xi \eta \varphi \psi dx dy \right)^{\frac{1}{q}} \\
& \times \left(\iint_{B_x(1,2\rho) \times B_y(\rho,2\rho^\varepsilon)} \frac{|G_2|^{q'} \xi \varphi}{|x|^{\sigma_1(q'-1)} |y|^{\sigma_2(q'-1)} \eta^{q'-1} \psi^{q'-1}} dx dy \right)^{\frac{1}{q'}} \\
& \leq \left(\iint_{B'} |x|^{\sigma_1} |y|^{\sigma_2} |u|^q \xi \eta \varphi \psi dx dy \right)^{\frac{1}{q}} \left(J_1^{\frac{1}{q'}} + J_2^{\frac{1}{q'}} \right), \tag{2.5}
\end{aligned}$$

where $\frac{1}{q} + \frac{1}{q'} = 1$ and J_1, J_2 one the second integrals in the first and second summed.

Hence

$$Q \leq C (J_1 + J_2). \tag{2.6}$$

Estimate the integrals J_1, J_2 separately.

As first consider G_1 and G_2 .

$$\begin{aligned}
G_1 & = 2 (\nabla_x (|x|^\alpha \Delta_x \xi), \nabla_x \varphi) + |x|^\alpha \Delta_x \xi \Delta_x \varphi + 2 \Delta_x (|x|^\alpha (\nabla_x \xi, \nabla_x \varphi)) \\
& + \Delta_x (|x|^\alpha \xi \Delta_x \varphi) = 2 \frac{d}{dr} \left(r^\alpha \left(\frac{d^2 \xi}{dr^2} + \frac{n-1}{r} \frac{d\xi}{dr} \right) \right) \frac{d\varphi}{dr} \\
& \quad + r^\alpha \left(\frac{d^2 \xi}{dr^2} + \frac{n-1}{r} \frac{d\xi}{dr} \right) \left(\frac{d^2 \varphi}{dr^2} + \frac{n-1}{r} \frac{d\varphi}{dr} \right) \\
& \quad + \left(\frac{d^2}{dr^2} + \frac{n-1}{r} \frac{d}{dr} \right) \left(r^\alpha \xi \left(\frac{d^2 \varphi}{dr^2} + \frac{n-1}{r} \frac{d\varphi}{dr} \right) \right) \\
& + 2 \left(\frac{d^2}{dr^2} + \frac{n-1}{r} \frac{d}{dr} \right) \left(r^\alpha \frac{d\xi}{dr} \frac{d\varphi}{dr} \right) = \left(2\alpha r^{\alpha-1} \left(\frac{d^2 \xi}{dr^2} + \frac{n-1}{r} \frac{d\xi}{dr} \right) \right. \\
& \quad \left. + 2r^\alpha \left(\frac{d^3 \xi}{dr^3} + \frac{n-1}{r} \frac{d^2 \xi}{dr^2} - \frac{n-1}{r^2} \frac{d\xi}{dr} \right) \right) \frac{d\varphi}{dr} \\
& + r^\alpha \left(\frac{d^2 \xi}{dr^2} + \frac{n-1}{r} \frac{d\xi}{dr} \right) \frac{d^2 \varphi}{dr^2} + (n-1)r^{\alpha-1} \left(\frac{d^2 \xi}{dr^2} + \frac{n-1}{r} \frac{d\xi}{dr} \right) \frac{d\varphi}{dr} \\
& + \left(\alpha(\alpha-1)r^{\alpha-2} \xi + 2\alpha r^{\alpha-1} \frac{d\xi}{dr} + r^\alpha \frac{d^2 \xi}{dr^2} \right) \left(\frac{d^2 \varphi}{dr^2} + \frac{n-1}{r} \frac{d\varphi}{dr} \right) \\
& + 2 \left(\alpha r^{\alpha-1} \xi + r^\alpha \frac{d\xi}{dr} \right) \left(\frac{d^3 \varphi}{dr^3} + \frac{n-1}{r} \frac{d^2 \varphi}{dr^2} - \frac{n-1}{r^2} \frac{d\varphi}{dr} \right) \\
& + r^\alpha \xi \left(\frac{d^4 \varphi}{dr^4} + \frac{n-1}{r} \frac{d^3 \varphi}{dr^3} - 2 \frac{n-1}{r^2} \frac{d^2 \varphi}{dr^2} + 2 \frac{n-1}{r^3} \frac{d\varphi}{dr} \right) \\
& \quad + \frac{n-1}{r} \left(\alpha r^{\alpha-1} \xi + r^\alpha \frac{d\xi}{dr} \right) \left(\frac{d^2 \varphi}{dr^2} + \frac{n-1}{r} \frac{d\varphi}{dr} \right) \\
& + (n-1)r^{\alpha-1} \xi \left(\frac{d^3 \varphi}{dr^3} + \frac{n-1}{r} \frac{d^2 \varphi}{dr^2} - \frac{n-1}{r^2} \frac{d\varphi}{dr} \right) + 2 \frac{d^2}{dr^2} \left(r^\alpha \frac{d\xi}{dr} \right) \frac{d\varphi}{dr}
\end{aligned}$$

$$\begin{aligned}
& +4 \frac{d}{dr} \left(\alpha r^{\alpha-1} \frac{d\xi}{dr} + r^\alpha \frac{d^2\xi}{dr^2} \right) \frac{d^2\varphi}{dr^2} + 2r^\alpha \frac{d\xi}{dr} \frac{d^3\varphi}{dr^3} \\
& + 2 \frac{n-1}{r} \left(\alpha r^{\alpha-1} \frac{d\xi}{dr} + r^\alpha \frac{d^2\xi}{dr^2} \right) \frac{d\varphi}{dr} + 2(n-1)r^{\alpha-1} \frac{d\xi}{dr} \frac{d^2\varphi}{dr^2} \\
& = \left[2\alpha r^{\alpha-1} \left(\frac{d^2\xi}{dr^2} + \frac{n-1}{r} \frac{d\xi}{dr} \right) + 2r^\alpha \left(\frac{d^3\xi}{dr^3} + \frac{n-1}{r} \frac{d^2\xi}{dr^2} - \frac{n-1}{r^2} \frac{d\xi}{dr} \right) \right. \\
& + (n-1)r^{\alpha-1} \left(\frac{d^2\xi}{dr^2} + \frac{n-1}{r} \frac{d\xi}{dr} \right) + \left(\alpha(\alpha-1)r^{\alpha-2}\xi + 2\alpha r^{\alpha-1} \frac{d\xi}{dr} + r^\alpha \frac{d^2\xi}{dr^2} \right) \frac{n-1}{r} \\
& - 2 \frac{n-1}{r^2} \left(\alpha r^{\alpha-1}\xi + r^\alpha \frac{d\xi}{dr} \right) + 2(n-1)r^{\alpha-3}\xi + \frac{(n-1)^2}{r^2} \left(\alpha r^{\alpha-1}\xi + r^\alpha \frac{d\xi}{dr} \right) \\
& - (n-1)^2 r^{\alpha-3}\xi + 2 \left(\alpha(\alpha-1)r^{\alpha-2} \frac{d\xi}{dr} + 2\alpha r^{\alpha-1} \frac{d^2\xi}{dr^2} + r^\alpha \frac{d^3\xi}{dr^3} \right) \\
& + 2 \frac{n-1}{r} \left(\alpha r^{\alpha-1} \frac{d\xi}{dr} + r^\alpha \frac{d^2\xi}{dr^2} \right) \left. \right] \frac{d\varphi}{dr} + \left[r^\alpha \left(\frac{d^2\xi}{dr^2} + \frac{n-1}{r} \frac{d\xi}{dr} \right) \right. \\
& + \alpha(\alpha-1)r^{\alpha-2}\xi + 2\alpha r^{\alpha-1} \frac{d\xi}{dr} + r^\alpha \frac{d^2\xi}{dr^2} + \frac{2(n-1)}{r} \left(\alpha r^{\alpha-1}\xi + r^\alpha \frac{d\xi}{dr} \right) \\
& - 2(n-1)r^{\alpha-2}\xi + \frac{n-1}{r} \left(\alpha r^{\alpha-1}\xi + r^\alpha \frac{d\xi}{dr} \right) + (n-1)^2 r^{\alpha-2}\xi \\
& + 4\alpha r^{\alpha-1} \frac{d\xi}{dr} + 4r^\alpha \frac{d^2\xi}{dr^2} + 2(n-1)r^{\alpha-1} \frac{d\xi}{dr} \left. \right] \frac{d^2\varphi}{dr^2} \\
& + \left[2\alpha r^{\alpha-1}\xi + 2r^\alpha \frac{d\xi}{dr} + 2(n-1)r^{\alpha-1}\xi + 2r^\alpha \frac{d\xi}{dr} \right] \frac{d^3\varphi}{dr^3} + r^\alpha \xi \frac{d^4\varphi}{dr^4}. \quad (2.7)
\end{aligned}$$

Since for $|x| > 1$

$$\left| \frac{d^i \xi}{dr^i} \right| \leq C r^{-\frac{n+\alpha-4}{2} + \alpha - i}, \quad i = 0, 1, 2, 3, 4,$$

then from (2.7) we get

$$|G_1| \leq C r^{-\frac{n+\alpha-4}{2} + \alpha - \alpha - 4} \sum_{i=1}^4 r^i \left| \frac{d^i \varphi}{dr^i} \right|. \quad (2.8)$$

In a similar way, we get

$$|G_2| \leq C r^{-\frac{m-4}{2} + \beta - 4} \sum_{i=1}^4 r^i \left| \frac{d^i \psi}{dr^i} \right|. \quad (2.9)$$

Using (2.8),(2.9) and making substitution $s = \frac{r}{\rho}, \tau = \frac{r}{\rho^\varepsilon}, \tilde{\varphi}(s) = \varphi(s\rho), \tilde{\psi}(\tau) = \psi(\tau\rho^\varepsilon)$ for J_1, J_2 we get the following estimaties.

$$|J_1| \leq \iint_{B_x(\rho, 2\rho) \times B_y(1, 2\rho^\varepsilon)} \frac{|G_1|^{q'} \eta \psi}{|x|^{\sigma_1(q'-1)} |y|^{\sigma_2(q'-1)} \xi^{q'-1} \varphi^{q'-1}} dx dy$$

$$\begin{aligned}
&\leq \int_{B_x(\rho, 2\rho)} \frac{|G_1|^{q'}}{|x|^{\sigma_1(q'-1)} \xi^{q'-1} \varphi^{q'-1}} dx \int_{B_y(1, 2\rho^\varepsilon)} \frac{\eta \psi}{|y|^{\sigma_2(q'-1)}} dy \\
&\leq C \int_\rho^{2\rho} \frac{r^{(-\frac{n+\alpha-4}{2} + \alpha_- + \alpha - 4)q'} \left(\sum_{i=1}^4 r^i \left| \frac{d^i \varphi}{dr^i} \right| \right)^{q'} r^{n-1}}{r^{\sigma_1(q'-1)} r^{(-\frac{n+\alpha-4}{2} + \alpha_-)(q'-1)} \varphi^{q'-1}} dr \\
&\times \int_1^{2\rho^\varepsilon} \frac{r^{-\frac{m-4}{2} + \beta_-} r^{m-1}}{r^{\sigma_2(q'-1)}} dr \leq c \rho^{(\alpha-4)q'+b} \int_1^2 \frac{s^{(-\frac{n+\alpha-4}{2} + \alpha_- + \alpha - 4) \left(\sum_{i=1}^4 s^i \left| \frac{d^i \tilde{\varphi}}{ds^i} \right| \right)^{q'}}}{s^{\sigma_1(q'-1)} s^{(-\frac{n+\alpha-4}{2} + \alpha_-)(q'-1) - n + 1} \tilde{\varphi}^{q'-1}} ds \\
&\times \begin{cases} r^a |1|^{2\rho^\varepsilon}, & \text{if } a \neq 0, \\ \ln r |1|^{2\rho^\varepsilon}, & \text{if } a = 0 \end{cases} \leq C \tilde{J}_1 \begin{cases} \rho^{(\alpha-4)q'+b+\varepsilon a}, & \text{if } a > 0, \\ \rho^{(\alpha-4)q'+b} \ln 2\rho^\varepsilon, & \text{if } a = 0, \\ \rho^{(\alpha-4)q'+b}, & \text{if } a < 0. \end{cases} \quad (2.10)
\end{aligned}$$

$$\begin{aligned}
|J_2| &\leq \iint_{B_x(1, 2\rho) \times B_y(\rho^\varepsilon, 2\rho^\varepsilon)} \frac{|G_2|^{q'} \xi \varphi}{|x|^{\sigma_1(q'-1)} |y|^{\sigma_2(q'-1)} \eta^{q'-1} \psi^{q'-1}} dx dy \\
&\leq \int_{B_x(1, 2\rho)} \frac{\xi \varphi}{|x|^{\sigma_1(q'-1)}} dx \int_{B_y(\rho^\varepsilon, 2\rho^\varepsilon)} \frac{|G_2|^{q'}}{|y|^{\sigma_2(q'-1)} \eta^{q'-1} \psi^{q'-1}} dy \\
&\leq C \int_1^{2\rho} r^{-\frac{n+\alpha-4}{2} + \alpha_- - \sigma_1(q'-1) + n - 1} dr \int_{\rho^\varepsilon}^{2\rho^\varepsilon} \frac{r^{(-\frac{m-4}{2} + \beta_- - 4)q'} \left(\sum_{i=1}^4 r^i \left| \frac{d^i \psi}{dr^i} \right| \right)^{q'} r^{m-1}}{r^{\sigma_2(q'-1)} r^{(-\frac{m-4}{2} + \beta_-)(q'-1)} \psi^{q'-1}} dr \\
&\leq C \rho^{\varepsilon(-4q'+a)} \int_1^2 \frac{\tau^{(-\frac{m-4}{2} + \beta_- - 4)q'} \left(\sum_{i=1}^4 \tau^i \left| \frac{d^i \tilde{\psi}}{d\tau^i} \right| \right)^{q'} \tau^{m-1}}{\tau^{\sigma_2(q'-1)} \tau^{(-\frac{m-4}{2} + \beta_-)(q'-1)} \tilde{\psi}} d\tau \\
&\times \begin{cases} \rho^b, & \text{if } b > 0, \\ \ln 2\rho, & \text{if } b = 0, \\ 1, & \text{if } b < 0 \end{cases} \leq C \tilde{J}_2 \begin{cases} \rho^{-4\varepsilon q' + \varepsilon a + b}, & \text{if } b > 0, \\ \rho^{-4\varepsilon q' + \varepsilon a} \ln 2\rho, & \text{if } b = 0, \\ \rho^{-4\varepsilon q' + \varepsilon a}, & \text{if } b < 0. \end{cases} \quad (2.11)
\end{aligned}$$

By \tilde{J}_1, \tilde{J}_2 we denote the last integral in the inequalities (2.8), (2.9). It is known that under appropriate choice of functions φ and ψ the integrals \tilde{J}_1, \tilde{J}_2 converge (see [15]).

Now we take such ε that

$$-4\varepsilon q' + \varepsilon a + b = (\alpha - 4)q' + b + \varepsilon a.$$

Hence

$$-4\varepsilon = (\alpha - 4) \Rightarrow \varepsilon = \frac{4 - \alpha}{4}.$$

Then, taking into account (2.10),(2.11) in (2.6) we get the followings

$$Q \leq \begin{cases} C\rho^{(\alpha-4)q'+b+\frac{4-\alpha}{4}a} & \text{if } a > 0, & b > 0, \\ C\rho^{(\alpha-4)q'} \ln \rho & \text{if } a = 0, & b = 0, \\ C\rho^{(\alpha-4)q'+\frac{4-\alpha}{4}a}(c_1 + c_2 \ln \rho) & \text{if } a > 0, & b = 0, \\ C\rho^{(\alpha-4)q'+b} (c_1 + c_2 \ln \rho) & \text{if } a = 0, & b > 0, \\ C\rho^{(\alpha-4)q'} & \text{if } a < 0, & b < 0, \\ C\rho^{(\alpha-4)q'} \ln \rho & \text{if } a = 0, b < 0 \text{ or } a < 0, b = 0, \\ C\rho^{(\alpha-4)q'+\frac{4-\alpha}{4}a} & \text{if } a > 0, & b < 0, \\ C\rho^{(\alpha-4)q'+b} & \text{if } a < 0, & b > 0. \end{cases} \quad (2.12)$$

Let us consider separate cases.

1) Let at first $a \leq 0, b \leq 0$. Since $\alpha - 4 < 0$, then it follows from (2.12) that when tending of $\rho \rightarrow +\infty, Q \leq 0$.

So,

$$\iint_{B'} |x|^{\sigma_1} |y|^{\sigma_2} |u|^q \xi \eta dx dy \leq 0.$$

Hence we get $u \equiv 0$. From the inequality $a \leq 0, b \leq 0$ it follows

$$\frac{m+4}{2} + \beta_- - \sigma_2(q'-1) \leq 0, \quad \frac{n-\alpha+4}{2} + \alpha_- - \sigma_1(q'-1) \leq 0.$$

From here

$$q \leq 1 + \frac{\sigma_2}{\frac{m+4}{2} + \beta_-}, \quad q \leq 1 + \frac{\sigma_1}{\frac{n-\alpha+4}{2} + \alpha_-}.$$

As a result, we get that if $\sigma_1, \sigma_2 > 0$ and $1 < q \leq \min \left\{ 1 + \frac{\sigma_1}{\frac{n-\alpha+4}{2} + \alpha_-}, 1 + \frac{\sigma_2}{\frac{m+4}{2} + \beta_-} \right\}$,

then equation (1.1) has no global solutions.

2) Let $a > 0, b \leq 0$. This means that

$$q > 1 + \frac{\sigma_2}{\frac{m+4}{2} + \beta_-}, \quad q \leq 1 + \frac{\sigma_1}{\frac{n-\alpha+4}{2} + \alpha_-}.$$

From (2.12) we get

$$Q \leq C\rho^{(\alpha-4)q'+\frac{4-\alpha}{4}(\frac{m+4}{2}+\beta_- - \sigma_2(q'-1))} \text{ for } a > 0, b < 0, \quad (2.13)$$

$$Q \leq C\rho^{(\alpha-4)q'+\frac{4-\alpha}{4}(\frac{m+4}{2}+\beta_- - \sigma_2(q'-1))} \ln \rho \text{ for } a > 0, b = 0. \quad (2.14)$$

If

$$\begin{aligned} & (\alpha-4)q' + \frac{4-\alpha}{4} \left(\frac{m+4}{2} + \beta_- - \sigma_2(q'-1) \right) \\ &= \left(\alpha-4 - \sigma_2 \frac{4-\alpha}{4} \right) (q'-1) + \frac{4-\alpha}{4} \left(\frac{m+4}{2} + \beta_- - 4 \right) < 0, \end{aligned}$$

i.e.

$$q < 1 + \frac{4-\alpha + \frac{4-\alpha}{4}\sigma_2}{\frac{4-\alpha}{4} \left(\frac{m+4}{2} + \beta_- - 4 \right)} = 1 + \frac{4 + \sigma_2}{\frac{m+4}{2} + \beta_- - 4},$$

then when tending $\rho \rightarrow +\infty$ from (2.13), (2.14) we get $Q \leq 0$. Hence, as in the first case we get $u \equiv 0$. What if $q = 1 + \frac{4+\sigma_2}{\frac{m+4}{2}+\beta_-}$, then from (2.10),(2.11),(2.12) it follows that $|J_1| \leq C, |J_2| \leq C$ and $Q \leq C$. Tending $\rho \rightarrow +\infty$ from (2.13) we have

$$\iint_{B'} |x|^{\sigma_1} |y|^{\sigma_2} |u|^q \xi \eta dx dy \leq C. \quad (2.15)$$

From (2.15) and from the property of the integral it follows, that as $\rho \rightarrow +\infty$

$$\iint_{B_x(\rho, 2\rho) \times B_y(1, 2\rho^\varepsilon)} |x|^{\sigma_1} |y|^{\sigma_2} |u|^q \xi \eta dx dy \rightarrow 0, \quad (2.16)$$

$$\iint_{B_x(1, 2\rho) \times B_y(\rho^\varepsilon, 2\rho^\varepsilon)} |x|^{\sigma_1} |y|^{\sigma_2} |u|^q \xi \eta dx dy \rightarrow 0. \quad (2.17)$$

From (2.5) we get that

$$\begin{aligned} \iint_{B_x(1, \rho) \times B_y(1, 2\rho^\varepsilon)} |x|^{\sigma_1} |y|^{\sigma_2} |u|^q \xi \eta dx dy &\leq \iint_{B_x(1, 2\rho) \times B_y(1, 2\rho^\varepsilon)} |x|^{\sigma_1} |y|^{\sigma_2} |u|^q \xi \eta \psi dx dy \\ &\leq \left(\iint_{B_x(\rho, 2\rho) \times B_y(1, 2\rho^\varepsilon)} |x|^{\sigma_1} |y|^{\sigma_2} |u|^q \xi \eta \psi dx dy \right)^{\frac{1}{q}} J_1^{\frac{1}{q'}} \\ &\quad + \left(\iint_{B_x(1, 2\rho) \times B_y(\rho^\varepsilon, 2\rho^\varepsilon)} |x|^{\sigma_1} |y|^{\sigma_2} |u|^q \xi \eta \psi dx dy \right)^{\frac{1}{q}} J_2^{\frac{1}{q'}} \\ &\leq C \left(\iint_{B_x(\rho, 2\rho) \times B_y(1, 2\rho^\varepsilon)} |x|^{\sigma_1} |y|^{\sigma_2} |u|^q \xi \eta dx dy \right)^{\frac{1}{q}} \\ &\quad + C \left(\iint_{B_x(1, 2\rho) \times B_y(\rho^\varepsilon, 2\rho^\varepsilon)} |x|^{\sigma_1} |y|^{\sigma_2} |u|^q \xi \eta dx dy \right)^{\frac{1}{q}}. \end{aligned}$$

Hence, by virtue of (2.16),(2.17) tending $\rho \rightarrow +\infty$ we get

$$\iint_{B'} |x|^{\sigma_1} |y|^{\sigma_2} |u|^q \xi \eta dx dy \leq 0.$$

This means that in this case too $u \equiv 0$. As a result, we got that if $\sigma_1 > 0, \sigma_2 > -4$, $q > \max \left\{ 1, 1 + \frac{\sigma_2}{\frac{m+4}{2}+\beta_-} \right\}$, $q < \left\{ 1 + \frac{\sigma_1}{\frac{n-\alpha+4}{2}+\alpha_-} \right\}$, $q \leq \left\{ 1 + \frac{\sigma_2+4}{\frac{m+4}{2}+\beta_-} \right\}$ and $q > \max \left\{ 1, 1 + \frac{\sigma_2}{\frac{m+4}{2}+\beta_-} \right\}$,

$q = \left\{ 1 + \frac{\sigma_1}{\frac{n-\alpha+4}{2} + \alpha_-} \right\}$, $q < \left\{ 1 + \frac{\sigma_2+4}{\frac{m+4}{2} + \beta_- - 4} \right\}$ the equation (1.1) has no trivial global solutions.

3) Let $a \leq 0, b > 0$. This means that

$$q \leq 1 + \frac{\sigma_2}{\frac{m+4}{2} + \beta_-}, \quad q > 1 + \frac{\sigma_1}{\frac{n-\alpha+4}{2} + \alpha_-}.$$

From (2.12) we get

$$Q \leq C\rho^{(\alpha-4)q'+b} \text{ for } a < 0, b > 0, \quad (2.18)$$

$$Q \leq C\rho^{(\alpha-4)q'+b} \ln \rho \text{ for } a = 0, b > 0. \quad (2.19)$$

If

$$\begin{aligned} (\alpha-4)q' + b &= (\alpha-4)(q'-1) + \alpha - 4 + \\ &+ \frac{n-\alpha+4}{2} + \alpha_- - \sigma_1(q'-1) < 0, \end{aligned}$$

i.e.

$$q < 1 + \frac{\sigma_1 + 4 - \alpha}{\frac{n-\alpha+4}{2} + \alpha_- + \alpha - 4},$$

then tendig $\rho \rightarrow +\infty$ from (2.18), (2.19) we get, $Q \leq 0$ and respectiely $u \equiv 0$.

Similar in the case 2) we can show that for $a < 0, b > 0$ and in the case $q = 1 + \frac{\sigma_1+4-\alpha}{\frac{n-\alpha+4}{2} + \alpha_- + \alpha - 4}$ too $u \equiv 0$.

As a result we get that if $\sigma_2 > 0, \sigma_1 + 4 - \alpha > 0$, $q > \max \left\{ 1, 1 + \frac{\sigma_1}{\frac{n-\alpha+4}{2} + \alpha_-} \right\}$, then at $q < 1 + \frac{\sigma_2}{\frac{m+4}{2} + \beta_-}$, $q \leq 1 + \frac{\sigma_1+4-\alpha}{\frac{n-\alpha+4}{2} + \alpha_- + \alpha - 4}$ or $q = 1 + \frac{\sigma_2}{\frac{m+4}{2} + \beta_-}$, $q < 1 + \frac{\sigma_1+4-\alpha}{\frac{n-\alpha+4}{2} + \alpha_- + \alpha - 4}$ equation (1.1) has no non-trivial global solutions.

4) Let now $a > 0, b > 0$. This means that

$$q > 1 + \frac{\sigma_1}{\frac{n-\alpha+4}{2} + \alpha_-}, \quad q > 1 + \frac{\sigma_2}{\frac{m+4}{2} + \beta_-}.$$

In this case if

$$\begin{aligned} (\alpha-4)q' + b + \frac{4-\alpha}{4}a &= (\alpha-4)(q'-1) + \alpha - 4 \\ &+ \frac{n-\alpha+4}{2} + \alpha_- - \sigma_1(q'-1) + \frac{4-\alpha}{4} \left(\frac{m+4}{2} + \beta_- - \sigma_2(q'-1) \right) \\ &= - \left(4 - \alpha + \sigma_1 + \frac{4-\alpha}{4}\sigma_2 \right) (q'-1) + \alpha - 4 \\ &+ \frac{n-\alpha+4}{2} + \alpha_- + \frac{4-\alpha}{4} \left(\frac{m+4}{2} + \beta_- \right) < 0, \end{aligned} \quad (2.20)$$

then tending $\rho \rightarrow +\infty$ from (2.12) we get $Q \leq 0$. Hence as carlier it follows that $u \equiv 0$. But if $(\alpha-4)q' + b + \frac{4-\alpha}{4}a = 0$, from (2.10),(2.11),(2.12) it follows that $|J_1| \leq C, |J_2| \leq C, Q \leq C$. As in the case 2) from (2.5) we get that in this case also $Q \leq 0$ and respectiely $u \equiv 0$.

We can write inequality (2.20) in the following form

$$q < 1 + \frac{4 - \alpha + \sigma_1 + \frac{4-\alpha}{4}\sigma_2}{\alpha - 4 + \frac{n-\alpha+4}{2} + \alpha_- + \frac{4-\alpha}{4} \left(\frac{m+4}{2} + \beta_-\right)}.$$

As a result, we get that if $4 - \alpha + \sigma_1 + \frac{4-\alpha}{4}\sigma_2 > 0$,

$$\max \left\{ 1 + \frac{\sigma_1}{\frac{n-\alpha+4}{2} + \alpha_-}, 1 + \frac{\sigma_2}{\frac{m+4}{2} + \beta_-} \right\}$$

$$< q \leq 1 + \frac{4 - \alpha + \sigma_1 + \frac{4-\alpha}{4}\sigma_2}{\alpha - 4 + \frac{n-\alpha+4}{2} + \alpha_- + \frac{4-\alpha}{4} \left(\frac{m+4}{2} + \beta_-\right)},$$

then equation (1.1) has no non-trivial global solutions.

This completely prunes the theorem.

References

1. Bagirov Sh. G.: Absence of positive solutions of a second-order semilinear parabolic equation with time-periodic coefficients, Translation of *Differ. Uravn.* **50**(4), 551-555 (2014). *Differ. Equ.* 50 (4), 548-553 (2014).
2. Bagirov Sh.H., Aliyev M.J.: On absence of solutions of a semi-linear elliptic equation with biharmonic operator in the exterior of a ball, *Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci. Mathematics* **36** (4), 63-69 (2016).
3. Bagirov Sh.G., Nonexistence of solutions of a semilinear biharmonic equation with singular potential, *Math. Notes* **103**(1), 23-32 (2018).
4. Bagirov Sh. G., Guliyev K.A.: Non-existence of global solution of a semi-linear parabolic equation with a singular potential, *Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci. Mathematics*, **40**(1), 79-87 (2020).
5. Bidaut-Veron M.F., Pohozaev S.: Nonexistence results and estimates for some nonlinear elliptic problems, *J. Anal. Math.* **84**, 1-49 (2001).
6. Brezis H., Dupaigne Z., Tesei A.: On a semilinear elliptic equation with inverse-square potential, *Selecta Math. (N. S.)* **11**(1), 1-7 (2005).
7. Carriao P.C., Demarque R., Miyagaki O.H.: Nonlinear biharmonic problems with singular potentials, *Commun. Pure Appl. Anal.* **13**(6), 2141-2154 (2014).
8. Ghergu M., Taliaferro S.D.: Nonexistence of positive supersolutions of nonlinear biharmonic equations without the maximum principle, *Comm. Partial Differential Equations* **40** (6), 1029-1069 (2015).
9. Gidas B., Spruck J., Global and local behavior of positive solutions of linear elliptic equations, *Comm. Pure Appl. Math.* **34**(4), 525-598 (1981).
10. Kon'kov A.A.: On the behavior at infinity of the solutions of a class of second-order nonlinear equations, (Russian) *Mat. Zametki* **60**(1), 30-39 (1996); translation in *Math. Notes* **60** (1-2), 22-28 (1996).
11. Kon'kov A.A.: On solutions of quasilinear elliptic inequalities containing terms with lower-order derivatives, *Nonlinear Anal.* **90**, 121-134 (2013).
12. Kon'kov A.A.: On properties of solutions of quasilinear second-order elliptic inequalities, *Nonlinear Anal.* 123-124, 89-114 (2015).
13. Laptev G.G.: On the absence of solutions for a class of singular semilinear differential inequalities, (Russian) *Tr. Mat. Inst. Steklova* Funkts. Prostran., Garmon. Anal., *Differ. Uravn.* **232**, 223-235 (2001); translation in *Proc. Steklov Inst. Math.* **232**, 216-228 (2001).
14. Mamedov F.: A Poincare's inequality with non-uniformly dergenerating gradient, *Monatshefte fur Mathematik*, **194**(1), 151-165 (2021).

15. Mitidieri E., Pokhozhaev S.I.: A priori estimates and blow-up of solutions to nonlinear partial differential equations and inequalities, (Russian) *Tr. Mat. Inst. Steklova*, **234**, 1-384 (2001); translation in *Proc. Steklov Inst. Math.* **234**, 1-362 (2001).
16. Mitidieri E., Pokhozhaev S.I.: Absence of global positive solutions of quasilinear elliptic inequalities, (Russian) *Dokl. Akad. Nauk* **359**(4), 456-460 (1998).
17. Mitidieri E., Pokhozhaev S.I.: Absence of positive solutions for quasilinear elliptic problems in \mathbb{R}^N , (Russian) *Tr. Mat. Inst. Steklova*, Issled. po Teor. Differ. Funkts. Mnogikh Perem. i ee Prilozh. **227**, 192-222 (1999); translation in *Proc. Steklov Inst. Math.* **227**, 186-216 (1998).
18. Serrin J., Zou H.: Nonexistence of positive solutions of Lane-Emden system, *Differential Integral Equations* **9** (4), 635-653 (1996).
19. Serrin J.: Positive solutions of prescribed mean curvature problem, *Calculus of variations and partial differential equations (Trento, 1986)*, *Lect. Notes in Math.* (Springer-Verlag, Berlin), **1340**, 248-255 (1988).
20. Volodin Yu.V.: On the critical exponents of certain nonlinear boundary value problems with biharmonic operator in the exterior of a ball, (Russian) *Mat. Zametki* **79**(2), 201-212 (2006); translation in *Math. Notes* **79**(1-2), 185-195 (2006).
21. Volodin Yu.V.: The critical exponents of semilinear boundary-value problems with biharmonic operator in the exterior of a ball with boundary conditions of first type, *Uchen. Zap. Ross. Gos. Sots. Univ.* (8), 208-215 (2009).
22. Xu X.: Uniqueness theorem for the entire positive solutions of biharmonic equations in R^n , *Proc. Roy. Soc. Edinburgh Sec. A* **130**(3), 651-670 (2000).
23. Yao Y., Wang R., Shen Y.: Nontrivial solution for the class of semilinear biharmonic equation involving critical exponents, *Acta Math. Sci. Ser. B Engl. Ed.* **27**(3), 509-514 (2007).