

## On Riesz property and equivalent basis property of the system of root vector functions of Dirac-type operator

Vali M. Kurbanov · Elchin J. Ibadov\*

Received: 31.09.2023 / Revised: 28.01.2024 / Accepted: 15.02.2024

**Abstract.** Dirac-type operator is considered on the finite interval  $G = (a, b)$ . It is assumed that its coefficient (potential) is a complex-valued matrix function summable on  $G = (a, b)$ . Riesz property criterion for a system of root vector functions is established and theorem on equivalent basis property in  $L_p^2(G)$ ,  $1 < p < \infty$ , is proved.

**Keywords.** Dirac-type operator · root vector function · Riesz property · equivalent basis property.

**Mathematics Subject Classification (2010):** 34L10, 34L20, 34B05, 34B24

### 1 Main concepts and statement of results

Riesz and basis properties of the systems of root vector functions of Dirac-type operator are studied in this work. Root vector functions are considered in generalized sense, i.e. regardless of boundary conditions (see [1]). With such a generalization, V.A. Il'in in [1] found the necessary and sufficient conditions for unconditional basis property (Riesz basis property) of the systems of root vector functions of the operator  $L = -d^2/dx^2 + q(x)$  for  $L_2$ . The work [1] served as a starting point for many mathematicians to study the Bessel, unconditional basis and basis properties of the systems of root vector functions of higher order differential operators.

For a Dirac operator with a potential from the class  $L_2$ , Bessel property and unconditional basis property criteria have been established in [2]. Componentwise uniform equiconvergence on a compact, uniform convergence, Riesz property of the systems of root vector functions of Dirac operator and unconditional basis property for Dirac-type operator have been considered in [3-7].

Basis property and other spectral properties of root vector functions of Dirac operator (with boundary conditions) have been treated in [8-16] and the references therein. In [8], the Riesz basis property for Dirac operator with a potential from the class  $L_2$  and separated boundary conditions has been established. Dirac operator with a potential from the class  $L_2$  and general regular conditions has been studied in [9], where the Riesz basis property

---

\* Corresponding author

V.M. Kurbanov  
Ministry of Science and Education, Institute of Mathematics and Mechanics, Baku, Azerbaijan  
E-mail: q.vali@yahoo.com

E.J. Ibadov  
Azerbaijan State Pedagogical University, Baku, Azerbaijan  
E-mail: e.c.ibadov@yahoo.com

of subspaces and, in case of strongly regular boundary conditions, the Riesz basis property have been proved. The case where the potential belongs to the class  $L_p, p \geq 1$ , has been considered in [10, 11], where the Riesz basis property (with strongly regular boundary conditions) and the Riesz basis property of subspaces (with regular boundary conditions) have been established. For Dirac-type operator with a potential from  $L_1$  and strongly regular conditions, the Riesz basis property has been proved in [12].

Consider one-dimensional Dirac-type operator

$$Dy = B \frac{dy}{dx} + P(x)y, \quad y(x) = (y_1(x), y_2(x))^T,$$

where  $B = \begin{pmatrix} 0 & b_1 \\ b_2 & 0 \end{pmatrix}$ ,  $b_2 < 0 < b_1$ ,  $P(x) = \text{diag}(p_1(x), p_2(x))$ ,

and  $p_1(x), p_2(x)$  are complex-valued summable functions on the arbitrary finite interval  $G = (a, b)$  of the real axis.

Following [1], by the eigen vector function of the operator  $D$  corresponding to the complex eigenvalue  $\lambda$ , we will mean any complex-valued vector function  $\overset{\circ}{u}(x)$  not identically zero, which is absolutely continuous on every closed subinterval of  $G$  and satisfies the equation  $D \overset{\circ}{u} = \lambda \overset{\circ}{u}$  almost everywhere in  $G$ .

Similarly, by the associated vector function of degree  $l$ ,  $l \geq 1$ , corresponding to the same  $\lambda$  and the same eigenfunction  $\overset{\circ}{u}(x)$ , we will mean any complex-valued vector function  $\overset{l}{u}(x)$ , which is absolutely continuous on every closed subinterval of  $G$  and satisfies the equation  $D \overset{l}{u} = \lambda \overset{l}{u} + \overset{l-1}{u}$  almost everywhere in  $G$ .

Let  $\{u_k(x)\}_{k=1}^{\infty}$  be an arbitrary system of root (eigen- and associated) vector functions of the operator  $D$ , and  $\{\lambda_k\}_{k=1}^{\infty}$  be the corresponding system of eigenvalues. In the sequel we will assume that every vector function  $u_k(x)$  belongs to the system  $\{u_k(x)\}_{k=1}^{\infty}$  together with all corresponding associated functions of a lesser degree, and the lengths of the chains of root vector functions are uniformly bounded. This means, in particular, that every vector function  $u_k(x)$  satisfies the equation

$$Du_k = \lambda_k u_k + \theta_k u_{k-1}$$

almost everywhere in  $G$ , where  $\theta_k$  is equal to either 0 (in this case,  $u_k(x)$  is an eigen vector function) or 1 (in this case,  $u_k(x)$  is an associated vector function,  $\lambda_k = \lambda_{k-1}$ ).

Let  $L_p^2(G)$ ,  $p \geq 1$ , be a space of two-component vector functions  $f(x) = (f_1(x), f_2(x))^T$  with the norm

$$\|f\|_{p,2} = \left[ \int_G (|f_1(x)|^2 + |f_2(x)|^2)^{p/2} dx \right]^{1/p}.$$

In case  $p = \infty$ , the norm in this space is defined by the equality  $\|f\|_{\infty,2} = \sup_{x \in \overline{G}} \text{vrai } |f(x)|$ .

Obviously, for the vector functions  $f(x) \in L_p^2(G)$ ,  $g(x) \in L_q^2(G)$ ,  $p^{-1} + q^{-1} = 1$ ,  $p \geq 1$ , the "scalar product"

$$(f, g) = \int_a^b \sum_{j=1}^2 f_j(x) \overline{g_j(x)} dx$$

is defined.

**Definition 1.1.** A system  $\{\varphi_k(x)\}_{k=1}^{\infty} \subset L_q^2(G)$ ,  $q \geq 2$ , is called a Riesz system, or a system which satisfies the Riesz property, if there exists a constant  $M = M(p)$  such that the inequality

$$\sum_{k=1}^{\infty} |(f, \varphi_k)|^q \leq M \|f\|_{p,2}^q$$

holds for an arbitrary function  $f(x) \in L_p^2(G)$ , where  $p^{-1} + q^{-1} = 1$ .

**Definition 1.2.** A system  $\{\varphi_k(x)\}_{k=1}^{\infty} \subset L_p^2(G)$ ,  $p \geq 1$ , is called  $p$ -close to the system  $\{\psi_k(x)\}_{k=1}^{\infty} \subset L_p^2(G)$  in  $L_p^2(G)$  if the relation

$$\sum_{k=1}^{\infty} \|\varphi_k - \psi_k\|_{p,2}^p < \infty$$

holds.

**Definition 1.3.** Two sequences of elements in the Banach space  $X$  are called equivalent if there exists a bounded, linear and boundedly invertible operator in  $X$ , which maps one of these sequences into another.

The following theorems are proved in this work.

**Theorem 1.1** (Criterion of Reizs property). Let  $P(x) \in L_1(G)$  and there exist a constant  $C_0$  such that

$$|Im\lambda_k| \leq C_0, \quad k = 1, 2, \dots \quad (1.1)$$

Then, for the system  $\{u_k(x) \|u_k\|_{q,2}^{-1}\}_{k=1}^{\infty} \subset L_q^2(G)$  to be Riesz, it is necessary and sufficient that there exists a constant  $M_1$  such that the inequality

$$\sum_{|Re\lambda_k - \nu| \leq 1} 1 \leq M_1 \quad (1.2)$$

holds for every real number  $\nu$ .

Let  $D^*$  be a formal adjoint operator of  $D$ , i.e.  $D^* = -B^* \frac{d}{dx} + P^*(x)$ , where  $P^*(x)$  is an adjoint matrix function of  $P(x)$ , and  $B^*$  is an adjoint matrix of  $B$ . Denote by  $\{v_k(x)\}_{k=1}^{\infty}$  a biorthogonal adjoint system of  $\{u_k(x)\}_{k=1}^{\infty}$  and assume that it consists of root vector functions of the operator  $D^*$ , i.e.  $D^*v_k = \lambda_k v_k + \theta_{k+1} v_{k+1}$ .

**Theorem 1.2** (On equivalent basis property). Let  $1 < p \leq 2$ ,  $P(x) \in L_1(G)$ , the lengths of the chains of root vector functions be uniformly bounded, conditions (1.1), (1.2) be satisfied, there exist a constant  $M_2$  such that

$$\|u_k\|_{2,2} \|v_k\|_{2,2} \leq M_2, \quad k = 1, 2, \dots, \quad (1.3)$$

and the system  $\{u_k(x) \|u_k\|_{p,2}^{-1}\}_{k=1}^{\infty}$  be  $p$ -close to some basis  $\{\psi_k(x)\}_{k=1}^{\infty}$  in  $L_p^2(G)$ . Then the systems  $\{u_k(x) \|u_k\|_{p,2}^{-1}\}$  and  $\{v_k(x) \|u_k\|_{p,2}\}_{k=1}^{\infty}$  are the bases in  $L_p^2(G)$  and  $L_q^2(G)$ , respectively, and these systems are equivalent to the basis  $\{\psi_k(x)\}_{k=1}^{\infty}$  and its biorthogonal adjoint, respectively.

**Remark 1.1.** If in Theorem 1.2 the systems  $\{u_k(x)\}_{k=1}^{\infty}$  and  $\{v_k(x)\}_{k=1}^{\infty}$  are interchanged, then we get the basicity of the system  $\{u_k(x)\}_{k=1}^{\infty}$  in  $L_p^2(G)$  for  $p \geq 2$ .

## 2 Auxiliary statements.

Statements below will be used to prove the above theorems.

**Statement 2.1** (see [7]). *If the functions  $p_1(x)$  and  $p_2(x)$  belong to the class  $L_1^{loc}(G)$  and the points  $x - t, x, x + t$ , lie in the interval  $G$ , then the following formulas are true for the root vector function  $u_k(x)$ :*

$$u_k(x \pm t) = \left[ \cos \frac{\lambda_k t}{\sqrt{|b_1 b_2|}} I \mp \sin \frac{\lambda_k t}{\sqrt{|b_1 b_2|}} \frac{B}{\sqrt{|b_1 b_2|}} \right] u_k(x) \\ \pm B^{-1} \int_x^{x \pm t} \left( \sin \frac{\lambda_k(t-|\xi-x|)}{\sqrt{|b_1 b_2|}} \frac{B}{\sqrt{|b_1 b_2|}} \mp \cos \frac{\lambda_k(t-|\xi-x|)}{\sqrt{|b_1 b_2|}} I \right) \quad (2.1)$$

$$\times [P(\xi)u_k(\xi) - \theta_k u_{k-1}(\xi)] d\xi,$$

$$u_k(x - t) + u_k(x + t) = 2u_k(x) \cos \frac{\lambda_k t}{\sqrt{|b_1 b_2|}} \\ + B^{-1} \int_{x-t}^{x+t} \left( \sin \frac{\lambda_k(t-|\xi-x|)}{\sqrt{|b_1 b_2|}} \frac{B}{\sqrt{|b_1 b_2|}} - \text{sign}(\xi - x) \right) \quad (2.2) \\ \times \cos \frac{\lambda_k(t-|\xi-x|)}{\sqrt{|b_1 b_2|}} I [P(\xi)u_k(\xi) - \theta_k u_{k-1}(\xi)] d\xi,$$

where  $I$  is a unit matrix function.

**Statement 2.2** (see [7]). *Let the functions  $p_1(x)$  and  $p_2(x)$  belong to the class  $L_1(G)$ . Then there exist the constants  $C_i(n_k, G, b_1, b_2)$ ,  $i = 1, 2$ , independent of  $\lambda_k$  such that*

$$\|\theta_k u_{k-1}\|_{\infty, G} \leq C_1(n_k, G, b_1, b_2)(1 + |Im \lambda_k|) \|u_k\|_{\infty, G}, \quad (2.3)$$

$$\|u_k\|_{\infty, G} \leq C_2(n_k, G, b_1, b_2)(1 + |Im \lambda_k|)^{1/r} \|u_k\|_{r, G}, \quad (2.4)$$

where  $n_k$  is a degree of the root vector function  $u_k(x)$ ,  $r \geq 1$ .

## 3 Proof of the Riesz property criterion.

In this section, we prove Theorem 1.1 (On the Riesz property of the systems of root vector functions of the operator  $D$ ).

**Necessity.** Consider any real number  $\nu$ . Introduce an index set  $I_\nu = \{k : |Re \lambda_k - \nu| \leq 1, |Im \lambda_k| \leq C_0\}$ , where  $C_0$  is a constant appearing in the condition (1.1). Let's choose the positive numbers  $R$  and  $R^*$  such that  $R \leq R^*$  and the inequality  $\omega(R) \leq L^{-1}$  holds for every set  $E \subset \overline{G}$ ,  $mes G \leq 2R^*$ , where  $L$  is a positive number to be defined later and

$$\omega(R) = \sup_{E \subset \overline{G}} \left\{ \|P\|_{1, E} \right\}, \quad \|P\|_{1, E} = \int_E (|p_1(x)| + |p_2(x)|) dx.$$

Let  $x \in [a, \frac{a+b}{2}]$ . Let's write the mean value formula (2.2) for the points  $x, x+t, x+2t$ , where  $t \in [0, R]$ :

$$u_k(x) = 2u_k(x+t) \cos \frac{\lambda_k t}{\sqrt{|b_1 b_2|}} - u_k(x+2t) \\ + B^{-1} \int_x^{x+2t} \left\{ \sin \frac{\lambda_k(t-|x+t-\xi|)}{\sqrt{|b_1 b_2|}} \frac{B}{\sqrt{|b_1 b_2|}} - \text{sign}(\xi - x - t) \right.$$

$$- \cos \frac{\lambda_k(t - |x + t - \xi|)}{\sqrt{|b_1 b_2|}} I \left\{ [P(\xi)u_k(\xi) - \theta_k u_{k-1}(\xi)] d\xi \right\}.$$

Add and subtract the function  $2u_k(x+t) \cos \frac{\nu t}{\sqrt{|b_1 b_2|}}$  on the right-hand side of this equality and perform the operation  $R^{-1} \int_0^R dt$ . Then we get

$$\begin{aligned} u_k(x) &= 2R^{-1} \int_0^R u_k(x+t) \cos \frac{\nu t}{\sqrt{|b_1 b_2|}} dt - R^{-1} \int_0^R u_k(x+2t) dt \\ &\quad + 4R^{-1} \int_0^R u_k(x+t) \sin \frac{\lambda_k + \nu}{2\sqrt{|b_1 b_2|}} \sin \frac{\nu - \lambda_k}{2\sqrt{|b_1 b_2|}} dt \\ &\quad + R^{-1} B^{-1} \int_0^R \int_x^{x+2t} \left\{ \sin \frac{\lambda_k(t - |x + t - \xi|)}{\sqrt{|b_1 b_2|}} \frac{B}{\sqrt{|b_1 b_2|}} \right. \\ &\quad \left. - \operatorname{sign}(\xi - x - t) \cos \frac{\lambda_k(t - |x + t - \xi|)}{\sqrt{|b_1 b_2|}} I \right\} \\ &\quad \times [P(\xi)u_k(\xi) - \theta_k u_{k-1}(\xi)] d\xi. \end{aligned}$$

Using formula (2.1) in the third term, we get

$$\begin{aligned} u_k(x) &= R^{-1} \int_G u_k(z)V(z)dz + 4R^{-1} \int_0^R \left( \cos \frac{\lambda_k t}{\sqrt{|b_1 b_2|}} I \right. \\ &\quad \left. - \sin \frac{\lambda_k t}{\sqrt{|b_1 b_2|}} \frac{B}{\sqrt{|b_1 b_2|}} \right) \sin \frac{(\lambda_k + \nu)t}{2\sqrt{|b_1 b_2|}} \sin \frac{(\nu - \lambda_k)t}{2\sqrt{|b_1 b_2|}} dt u_k(x) \\ &\quad + 4R^{-1} B^{-1} \int_0^R \int_x^{x+t} \left\{ \sin \frac{\lambda_k(t - |\xi - x|)}{\sqrt{|b_1 b_2|}} \frac{B}{\sqrt{|b_1 b_2|}} + \cos \frac{\lambda_k(t - |\xi - x|)}{\sqrt{|b_1 b_2|}} I \right\} \\ &\quad \times [P(\xi)u_k(\xi) - \theta_k u_{k-1}(\xi)] \sin \frac{(\nu + \lambda_k)t}{2\sqrt{|b_1 b_2|}} \sin \frac{(\nu - \lambda_k)t}{2\sqrt{|b_1 b_2|}} dt \\ &\quad + R^{-1} B^{-1} \int_0^R \int_x^{x+2t} \left\{ \sin \frac{\lambda_k(t - |x + t - \xi|)}{\sqrt{|b_1 b_2|}} \frac{B}{\sqrt{|b_1 b_2|}} + \cos \frac{\lambda_k(t - |x + t - \xi|)}{\sqrt{|b_1 b_2|}} I \right\} \\ &\quad \times [P(\xi)u_k(\xi) - \theta_k u_{k-1}(\xi)] d\xi dt = R^{-1} \int_G u_k(z)V(z)dz + J_1 + J_2 + J_3, \quad (3.1) \end{aligned}$$

where  $V(z) = 2 \cos \frac{\nu(x-z)}{\sqrt{|b_1 b_2|}} - \frac{1}{2}$  for  $x \leq z \leq x+R$ ,  $V(z) = -\frac{1}{2}$  for  $x+R < z \leq x+2R$ , and  $V(z) = 0$  for  $z \notin [x, x+2R]$ .

Let  $k \in I_\nu$ . Let's estimate the integrals  $J_i$ ,  $i = \overline{1, 3}$ . Using the inequalities

$$|\sin z| \leq 2, \quad |\cos z| \leq 2, \quad |\sin z| \leq 2|z|, \quad (3.2)$$

which hold for  $|Imz| \leq 1$ , we obtain

$$|J_1| \leq 8R \left( \frac{2}{\sqrt{|b_1 b_2|}} + \frac{|b_1| + |b_2|}{|b_1 b_2|} \right) |\nu - \lambda_k| |u_k(x)| \leq 8R \left( \frac{2}{\sqrt{|b_1 b_2|}} + \frac{|b_1| + |b_2|}{|b_1 b_2|} \right)$$

$$\times (1 + |Im\lambda_k|) |u_k(x)| \leq 8R \left( \frac{2}{\sqrt{|b_1 b_2|}} + \frac{|b_1| + |b_2|}{|b_1 b_2|} \right) (1 + C_0) \|u_k\|_{\infty,2}.$$

Applying the inequalities (3.2) and the Holder inequality for  $p = 1, q = \infty$ , we find

$$|J_2| \leq 32 \left( \frac{2}{\sqrt{|b_1 b_2|}} + \frac{|b_1| + |b_2|}{|b_1 b_2|} \right) \left( \omega(R) \|u_k\|_{\infty,2} + \frac{R}{2} \|\theta_k u_{k-1}\|_{\infty,2} \right);$$

$$|J_3| \leq 2 \left( \frac{2}{\sqrt{|b_1 b_2|}} + \frac{|b_1| + |b_2|}{|b_1 b_2|} \right) \left( \omega(R) \|u_k\|_{\infty,2} + R \|\theta_k u_{k-1}\|_{\infty,2} \right).$$

Considering these estimates in the equality (3.1), we obtain

$$|u_k(x)| \leq R^{-1} \left| \int_G u_k(z) V(z) dz \right| + 8 \left( \frac{2}{\sqrt{|b_1 b_2|}} + \frac{|b_1| + |b_2|}{|b_1 b_2|} \right) \times (R(1 + C_0) + 5\omega(R)) \|u_k\|_{\infty,2} + 18R \left( \frac{2}{\sqrt{|b_1 b_2|}} + \frac{|b_1| + |b_2|}{|b_1 b_2|} \right) \|\theta_k u_{k-1}\|_{\infty,2}. \quad (3.3)$$

The inequality (3.3) can be proved similarly in case  $x \in [\frac{a+b}{2}, b]$ . In this case,  $V(z) = -\frac{1}{2}$  for  $x - 2R \leq z < x - R$ ,  $V(z) = 2 \cos \frac{\nu(x-z)}{\sqrt{|b_1 b_2|}} - \frac{1}{2}$  for  $x - R \leq z \leq x$ , and  $V(z) = 0$  for  $z \notin [x - 2R, x]$ .

Consequently, the inequality (3.3) is true for every  $x \in \bar{G}$ .

Applying the estimates (2.3), (2.4) and taking into account the relation  $1 + |Im\lambda_k| \leq 1 + C_0$ , from (3.3) we obtain

$$|u_k(x)| \leq R^{-1} \left| \int_G u_k(z) V(z) dz \right| + 8 \left( \frac{2}{\sqrt{|b_1 b_2|}} + \frac{|b_1| + |b_2|}{|b_1 b_2|} \right) \left\{ 5\omega(R) C_2(n_k, G, b_1, b_2) (1 + C_0)^{1/q} + RC_2(n_k, G, b_1, b_2) (1 + C_0)^{1+1/q} + 18RC_1(n_k, G, b_1, b_2) C_2(n_k, G, b_1, b_2) \theta_k (1 + C_0)^{1+1/q} \right\} \|u_k\|_{q,2}.$$

Due to the uniform boundedness of the lengths of the chains, we have

$$40 \left( \frac{2}{\sqrt{|b_1 b_2|}} + \frac{|b_1| + |b_2|}{|b_1 b_2|} \right) C_2(n_k, G, b_1, b_2) \leq \gamma_1 = const,$$

$$144 \left( \frac{2}{\sqrt{|b_1 b_2|}} + \frac{|b_1| + |b_2|}{|b_1 b_2|} \right) C_2(n_k, G, b_1, b_2) C_2(n_k, G, b_1, b_2) \leq \gamma_2 = const.$$

Consequently,

$$|u_k(x)| \leq R^{-1} \left| \int_G u_k(z) V(z) dz \right| + \left\{ \omega(R) \gamma_1 (1 + C_0)^{1/q} + \gamma_1 R (1 + C_0)^{1+1/q} + R \gamma_2 \theta_k (1 + C_0)^{1+1/q} \right\} \|u_k\|_{q,2}.$$

Multiplying both sides of this inequality by  $\|u_k\|_{q,2}^{-1}$ , raising to a degree  $q$  and applying the inequality  $\left| \sum_{i=1}^n a_i \right|^q \leq n^{q-1} \sum_{i=1}^n |a_i|^q$ , we find

$$\begin{aligned} & |u_k(x)|^q \|u_k\|_{q,2}^{-q} \leq 3^{q-1} R^{-q} \left\{ \left| \int_G u_k^1(z) V(z) dz \right|^q + \left| \int_G u_k^2(z) V(z) dz \right|^q \right\} \\ & \times \|u_k\|_{q,2}^{-q} + 3^{q-1} \left\{ \gamma_1 L^{-1} (1 + C_0)^{1/q} + R\gamma_1 (1 + C_0)^{1+1/q} + R\gamma_2 \theta_k (1 + C_0)^{1+1/q} \right\}^q, \end{aligned}$$

where  $u_k(x) = (u_k^1(x), u_k^2(x))^T$ .

By virtue of Riesz inequality and  $\|V\|_p^q \leq 3^q R^{q/p}$ , we obtain

$$\begin{aligned} & \sum_{k \in J} |u_k(x)|^q \|u_k\|_{q,2}^{-q} \leq 2 \cdot 3^{2q-1} M R^{q(\frac{1}{p}-1)} \\ & + 3^{q-1} \left\{ \gamma_1 L^{-1} (1 + C_0)^{1/q} + R\gamma_1 (1 + C_0)^{1+1/q} + R\gamma_2 \theta_k (1 + C_0)^{1+1/q} \right\} \sum_{k \in J} 1, \end{aligned}$$

where  $J \subset I_\nu$  is an arbitrary finite set of indices  $k$ , which correspond to the root functions  $u_k(x)$ . Integrating this inequality over  $x \in G$  and choosing  $R$  (consequently, the number  $L^{-1}$  too) small enough to have an estimate

$$3^{q-1} \left\{ \gamma_1 L^{-1} (1 + C_0)^{1/q} + R\gamma_1 (1 + C_0)^{1+1/q} + R\gamma_2 \theta_k (1 + C_0)^{1+1/q} \right\}^q < \frac{1}{2mesG},$$

we arrive at the inequality

$$\sum_{k \in J} 1 \leq 4 \cdot 3^{2q-1} M R^{-1} mesG,$$

which, due to the arbitrariness of the finite set  $J$  and the uniform boundedness of the chains of root vector functions, implies the necessity of the inequality (1.2).

**Sufficiency.** For simplicity we consider  $G = (0, 2\pi)$ . Note that in this case it suffices for us to establish the Bessel property of the system  $\left\{ u_k(x) \|u_k\|_{2,2}^{-1} \right\}_{k=1}^\infty$  in  $L_2^2(0, 2\pi)$ . In fact, due to the estimate (2.4) and the condition (1.1), for every vector function  $f(x) \in L_2^2(0, 2\pi)$  we have

$$\sup_k \left| \int_0^{2\pi} \left( f(x), u_k(x) \|u_k\|_{2,2}^{-1} \right) dx \right| \leq const \|f\|_{1,2}.$$

Therefore, by Riesz-Thorin interpolation theorem, (see, e.g., [17, p.144]), the system  $\left\{ u_k(x) \|u_k\|_{2,2}^{-1} \right\}_{k=1}^\infty$  is Riesz.

On the other hand, by (2.4) and (1.1), (1.2), we have

$$\|u_k\|_{2,2} \|u_k\|_{q,2}^{-1} \leq (2\pi)^{\frac{1}{2}} \|u_k\|_{\infty,2} \|u_k\|_{q,2}^{-1} \leq const, \quad k = 1, 2, \dots$$

Consequently, for every  $f(x) \in L_p^2(0, 2\pi)$ ,  $1 < p \leq 2$ , the estimate

$$\begin{aligned} & \sum_{k=1}^\infty \left| \left( f, u_k \|u_k\|_{2,2}^{-1} \right) \right|^q = \sum_{k=1}^\infty \|u_k\|_{2,2}^q \|u_k\|_{q,2}^{-q} \left| \left( f, u_k \|u_k\|_{2,2}^{-1} \right) \right|^q \\ & \leq const \sum_{k=1}^\infty \left| \left( f, u_k \|u_k\|_{2,2}^{-1} \right) \right|^q \leq M_3 \|f\|_{p,2}^q \end{aligned}$$

is true.

So, we have to prove the Bessel property of the system  $\left\{u_k(x) \|u_k\|_{2,2}^{-1}\right\}_{k=1}^{\infty}$  in  $L_2^2(0, 2\pi)$ . Considering the shift formula (2.1) for  $u_k(x+t)$  as  $x=0$  and then multiplying it scalarly by the vector function  $f(t) = (f_1(t), f_2(t))^T \in L_2^2(0, 2\pi)$ , we conclude that to prove the Besselness of the system  $\varphi_k(t) = u_k(t) \|u_k\|_{2,2}^{-1}, k = 1, 2, \dots$  in  $L_2^2(0, 2\pi)$  it suffices to get the validity of the following inequalities:

$$\sum_{k=1}^{\infty} \left| \int_0^{2\pi} \overline{f_i(t)} \cos \frac{\lambda_k t}{\sqrt{|b_1 b_2|}} dt \right|^2 |\varphi_k^i(0)|^2 \leq C \|f\|_{2,2}^2, i = 1, 2; \quad (3.4)$$

$$\sum_{k=1}^{\infty} \left| \int_0^{2\pi} \overline{f_i(t)} \sin \frac{\lambda_k t}{\sqrt{|b_1 b_2|}} dt \right|^2 |\varphi_k^{3-i}(0)|^2 \leq C \|f\|_{2,2}^2, i = 1, 2; \quad (3.5)$$

$$\sum_{k=1}^{\infty} \left| \int_0^{2\pi} \overline{f_1(t)} \int_0^t p_1(\xi) \varphi_k^1(\xi) \sin \frac{\lambda_k(t-\xi)}{\sqrt{|b_1 b_2|}} d\xi dt \right|^2 \leq C \|f\|_{2,2}^2, \quad (3.6)$$

$$\sum_{k=1}^{\infty} \left| \int_0^{2\pi} \overline{f_1(t)} \int_0^t p_2(\xi) \varphi_k^2(\xi) \cos \frac{\lambda_k(t-\xi)}{\sqrt{|b_1 b_2|}} d\xi dt \right|^2 \leq C \|f\|_{2,2}^2, \quad (3.7)$$

$$\sum_{k=1}^{\infty} \left| \int_0^{2\pi} \overline{f_2(t)} \int_0^t p_1(\xi) \varphi_k^1(\xi) \cos \frac{\lambda_k(t-\xi)}{\sqrt{|b_1 b_2|}} d\xi dt \right|^2 \leq C \|f\|_{2,2}^2, \quad (3.8)$$

$$\sum_{k=1}^{\infty} \left| \int_0^{2\pi} \overline{f_2(t)} \int_0^t p_2(\xi) \varphi_k^2(\xi) \sin \frac{\lambda_k(t-\xi)}{\sqrt{|b_1 b_2|}} d\xi dt \right|^2 \leq C \|f\|_{2,2}^2, \quad (3.9)$$

$$\sum_{k=1}^{\infty} \left| \theta_k \int_0^{2\pi} \overline{f_i(t)} \int_0^t \frac{u_{k-1}^i(\xi)}{\|u_k\|_{2,2}} \sin \frac{\lambda_k(t-\xi)}{\sqrt{|b_1 b_2|}} d\xi dt \right|^2 \leq C \|f\|_{2,2}^2, i = 1, 2; \quad (3.10)$$

$$\sum_{k=1}^{\infty} \left| \theta_k \int_0^{2\pi} \overline{f_i(t)} \int_0^t \frac{u_{k-1}^{3-i}(\xi)}{\|u_k\|_{2,2}} \cos \frac{\lambda_k(t-\xi)}{\sqrt{|b_1 b_2|}} d\xi dt \right|^2 \leq C \|f\|_{2,2}^2, i = 1, 2; \quad (3.11)$$

where  $\varphi_k^i(\xi) = u_k^i(\xi) \|u_k\|_{2,2}^{-1}$ .

Let's prove the estimate (3.4). By the estimate (2.4) and the conditions (1.1),(1.2), we have

$$|\varphi_k^i(0)| = |u_k^i(0)| \|u_k\|_{2,2}^{-1} \leq \|u_k\|_{\infty,2} \|u_k\|_{2,2}^{-1}$$

$$\leq C_2(n_k, G, b_1, b_2)(1 + C_0)^{1/2} \|u_k\|_{2,2} \|u_k\|_{2,2}^{-1} \leq C_2(n_k, G, b_1, b_2)(1 + C_0)^{1/2} = const,$$

because the sequence  $C_2(n_k, G, b_1, b_2)$  is bounded due to the condition (2.4). Therefore, for (3.4) to be valid it suffices that the inequality

$$\sum_{k=1}^{\infty} \left| \int_0^{2\pi} \overline{f_i(t)} \cos \frac{\lambda_k t}{\sqrt{|b_1 b_2|}} dt \right|^2 \leq C \|f\|_{2,2}^2, i = 1, 2, \quad (3.12)$$

holds.

Under conditions (1.1) and (1.2) with  $\nu \geq 1$ , the validity of the inequality (3.12) has been proved in [1]. Hence it follows the validity of (3.12) for  $Re\lambda_k \in (-\infty, +\infty), |Im\lambda_k| \leq C_0$ ,



because, by the condition of Theorem 1.1, the condition (1.2) holds for any  $\nu \in (-\infty, +\infty)$ . The inequality (3.5) can be proved in the same way.

Let's verify the inequalities (3.6)-(3.9). They all are proved similarly, so we will only prove (3.6). Denote

$$g_i(t, \xi) = \begin{cases} f_i(t + \xi), & 0 \leq t \leq 2\pi - \xi, \\ 0, & 2\pi - \xi < t \leq 2\pi, \end{cases}$$

where  $\xi \in [0, 2\pi]$ ,  $i = 1, 2$ . Then, by the estimate (2.4) for  $r = 2$  and the conditions (1.1), (1.2), we obtain

$$\begin{aligned} T_k &= \left| \int_0^{2\pi} \overline{f_1(t)} \int_0^t p_1(\xi) \varphi_k^1(\xi) \sin \frac{\lambda_k(t - \xi)}{\sqrt{|b_1 b_2|}} d\xi dt \right|^2 \\ &= \int_0^{2\pi} \overline{f_1(t)} \int_0^t p_1(\xi) \varphi_k^1(\xi) \sin \frac{\lambda_k(t - \xi)}{\sqrt{|b_1 b_2|}} d\xi dt \times \\ &\quad \times \int_0^{2\pi} f_1(t) \int_0^t \overline{p_1(\xi) \varphi_k^1(\xi) \sin \frac{\lambda_k(t - \xi)}{\sqrt{|b_1 b_2|}}} d\xi dt \\ &= \int_0^{2\pi} p_1(\xi) \varphi_k^1(\xi) \int_0^{2\pi} \overline{g_1(t, \xi)} \sin \frac{\lambda_k t}{\sqrt{|b_1 b_2|}} dt d\xi \times \\ &\quad \times \int_0^{2\pi} \overline{p(\tau) \varphi_k^1(\tau)} \int_0^{2\pi} g_1(r, \tau) \sin \frac{\lambda_k r}{\sqrt{|b_1 b_2|}} dr d\tau \\ &= \int_0^{2\pi} \int_0^{2\pi} p_1(\xi) \overline{p_1(\tau)} \varphi_k^1(\xi) \overline{\varphi_k^1(\tau)} \int_0^{2\pi} \overline{g_1(t, \xi)} \sin \frac{\lambda_k t}{\sqrt{|b_1 b_2|}} dt \times \\ &\quad \times \int_0^{2\pi} g_1(r, \tau) \sin \frac{\lambda_k r}{\sqrt{|b_1 b_2|}} dr d\xi d\tau \\ &\leq C_2^2(n_k, G, b_1, b_2) (1 + C_0) \int_0^{2\pi} \int_0^{2\pi} |p_1(\xi)| |p_1(\tau)| \left| \int_0^{2\pi} \overline{g_1(t, \xi)} \sin \frac{\lambda_k t}{\sqrt{|b_1 b_2|}} dt \right| \\ &\quad \times \left| \int_0^{2\pi} g_1(r, \tau) \sin \frac{\lambda_k r}{\sqrt{|b_1 b_2|}} dr \right| d\xi d\tau \\ &\leq \text{const} \int_0^{2\pi} \int_0^{2\pi} |p_1(\xi)| |p_1(\tau)| \left| \int_0^{2\pi} \overline{g_1(t, \xi)} \sin \frac{\lambda_k t}{\sqrt{|b_1 b_2|}} dt \right| \\ &\quad \times \left| \int_0^{2\pi} g_1(r, \tau) \sin \frac{\lambda_k r}{\sqrt{|b_1 b_2|}} dr \right| d\xi d\tau. \end{aligned}$$

Then, for arbitrary positive integer  $N$  we obtain

$$\begin{aligned} \sum_{k=1}^N T_k &\leq \text{const} \int_0^{2\pi} \int_0^{2\pi} |p_1(\xi)| |p_1(\tau)| \\ &\quad \times \left( \sum_{k=1}^N \left| \int_0^{2\pi} \overline{g_1(t, \xi)} \sin \frac{\lambda_k t}{\sqrt{|b_1 b_2|}} dt \right| \left| \int_0^{2\pi} g_1(r, \tau) \sin \frac{\lambda_k r}{\sqrt{|b_1 b_2|}} dr \right| \right) d\xi d\tau \end{aligned}$$

$$\leq \text{const} \int_0^{2\pi} \int_0^{2\pi} |p_1(\xi)| |p_1(\tau)| \|g_1(\cdot, \xi)\|_2 \|g_1(\cdot, \tau)\|_2 d\xi d\tau.$$

As the inequality  $\|g_1(\cdot, \xi)\|_2 \leq \|f_1\|_2$  holds for every fixed  $\xi \in [0, 2\pi]$ , we get

$$\sum_{k=1}^N T_k \leq \text{const} \|p_1\|_1^2 \|f_1\|_2^2 \leq \text{const} \|f\|_{2,2}^2.$$

Hence, due to the arbitrariness of the number  $N$ , we get the validity of the inequality (3.6).

Now let's prove (3.10). By (2.3), (2.4) and (1.1), (1.2), we have

$$\begin{aligned} \theta_k |u_{k-1}^i(\xi)| \|u_k\|_{2,2}^{-1} &\leq \theta_k C_1(n_k, G, b_1, b_2) C_2(n_k, G, b_1, b_2) (1 + C_0)^{\frac{2}{3}} \\ &\times \|u_k\|_{2,2} \|u_k\|_{2,2}^{-1} \leq C = \text{const} \end{aligned}$$

After changing the order of integration, the left-hand side of the inequality (3.10) is majorized from above by the series

$$C \sum_{k=1}^{\infty} \int_0^{2\pi} \left| \int_0^{2\pi} \frac{g_i(t, \xi) \sin \frac{\lambda_k t}{\sqrt{|b_1 b_2|}} dt \right|^2 d\xi.$$

This series converges due to the Bessel property of the system  $\left\{ \sin \frac{\lambda_k t}{\sqrt{|b_1 b_2|}} \right\}_{k=1}^{\infty}$ , and its sum is bounded from above by  $\text{const} \|f\|_{2,2}^2$ .

The inequality (3.10) is proved. The inequality (3.11) is proved similarly.

#### 4 Proof of the Theorem 1.2.

As the system  $\{v_k(x)\}_{k=1}^{\infty}$  consists of root vector functions of the operator  $D^*$  (formal adjoint of  $D$ ), by Theorem 1.1, the conditions (1.1) and (1.2) provide the Riesz property of the system  $\left\{ v_k(x) \|v_k\|_{q,2}^{-1} \right\}_{k=1}^{\infty}$  in  $L_p^2(G)$ ,  $1 < p \leq 2$ ,  $p^{-1} + q^{-1} = 1$ , i.e.

$$\sum_{k=1}^{\infty} \left| \left( f, v_k(x) \|v_k\|_{q,2}^{-1} \right) \right|^q \leq M \|f\|_{p,2}^q \quad (4.1)$$

for every vector function  $f(x) \in L_p^2(G)$ .

The inequality (4.1), the condition (1.3) and the  $p$ -closeness of the systems  $\left\{ u_k(x) \|u_k\|_{p,2}^{-1} \right\}_{k=1}^{\infty}$  and  $\{\psi_k(x)\}_{k=1}^{\infty}$  in  $L_p^2(G)$  imply that the series

$\sum_{k=1}^{\infty} \tilde{f}_k \|u_k\|_{p,2} \|v_k\|_{q,2} \left( u_k \|u_k\|_{p,2}^{-1} - \psi_k(x) \right)$  converges in  $L_p^2(G)$  for every  $f(x) \in L_p^2(G)$ ,

where  $\tilde{f}_k = \left( f, v_k \|v_k\|_{q,2}^{-1} \right)$ . Denote the sum of this series by  $Kf$ . As the sequence  $K_n f =$

$\sum_{k=1}^n \tilde{f}_k \|u_k\|_{p,2} \|v_k\|_{q,2} \left( u_k(x) \|u_k\|_{p,2}^{-1} - \psi_k(x) \right)$  is fundamental in  $L_p^2(G)$ , the linear operator  $K$  acts in  $L_p^2(G)$ , i.e.  $Kf \in L_p^2(G)$  for  $f(x) \in L_p^2(G)$ . Obviously,  $\|Kf - K_n f\|_{p,2} =$

$o(1) \|f\|_{p,2}$ , i.e. the sequence of finite dimensional operators  $\{K_n\}$  converges to the operator  $K$ . Consequently, this operator is compact in  $L_p^2(G)$ . Besides,  $K u_k \|u_k\|_{p,2}^{-1} - \psi_k$ , i.e.

$(E - K)u_k \|u_k\|_{p,2}^{-1} = \psi_k$ ,  $k \in N$ , where  $E$  is a unit operator.

Let's show that the operator  $E - K$  is continuously invertible. The compactness of the operator  $K$  and the Fredholm alternative imply that if the operator  $E - K$  is non-invertible, then there exists a non-zero element  $g \in L_q^2(G)$  such that  $(E - K)^*g = 0$ . The element  $g$  satisfies the relation

$$(g, \psi_k) = \left( g, (E - K)u_k \|u_k\|_{p,2}^{-1} \right) = \left( (E - K)^*g, u_k \|u_k\|_{p,2}^{-1} \right) = 0, \quad k \in N.$$

Hence, due to the basicity of the system  $\{\psi_k(x)\}_{k=1}^\infty$  for  $L_p^2(G)$ , it follows that the element  $g$  is equal to 0. The obtained contradiction proves the invertibility of the operator  $E - K$ . Consequently, the system  $\{u_k(x) \|u_k\|_{p,2}^{-1}\}_{k=1}^\infty$  is a basis for  $L_p^2(G)$ , and it is equivalent to the basis  $\{\psi_k(x)\}_{k=1}^\infty$ . If we denote by  $\{z_k(x)\}_{k=1}^\infty$  a biorthogonal adjoint system of  $\{\psi_k(x)\}_{k=1}^\infty$ , then  $v_k(x) \|u_k\|_{p,2} = (E - K)^*z_k(x)$ . This means that the system  $\{v_k(x) \|u_k\|_{p,2}\}_{k=1}^\infty$  is a basis in  $L_q^2(G)$  equivalent to the basis  $\{z_k(x)\}_{k=1}^\infty$ .

Theorem 1.2 is proved.

## References

1. Ilin V.A.: *Unconditional basis property on a closed interval for systems of eigen and associated functions of a second order differential operator*, Dokl. Akad. Nauk SSSR, **273**(5), 1048-1053 (1983).
2. Kurbanov V.M.: *On Bessel property and unconditional basicity of the system of root vector-functions of Dirac operator*, Diff. Uravn., **32**(12), 1608-1617 (1996).
3. Kurbanov V.M., Ismailova A.I.: *Componentwise uniform equiconvergence of expansions in root vector functions of the Dirac operator with the trigonometric expansion* Diff. Uravn., **48**(5), 648-662 (2012).
4. Kurbanov V.M., Ismailova A.I.: *Absolute and uniform convergence of expansions in root vector functions of Dirac operator*, Dokl. RAN. , **446**(4), 380-383 (2012) (in Russian)
5. Kurbanov V.M., Ismailova A.I.: *Riesz inequality for the system of root vector functions of Dirac operator*, Diff. Uravn., **48**(3), 334-340 (2012).
6. Kurbanov V.M., Ismailova A.I.: *Two-sided estimates for root vector functions of Dirac operator*. Diff. Uravn., 48(4), 487-497. (2012).
7. Kurbanov V.M., Ibadov E.J., Hajiyeva G.R.: *On Bessel property and unconditional basicity of the systems of root vector-functions of a Dirac-type operator*, Azerbaijan Journal of Mathematics 7(2), 21-32 (2017).
8. Trooshin I. and Yamamoto M.: *Riesz basis of root vectors of a nonsymmetric system of first-order ordinary differential operators and application to inverse eigenvalue problems*, Appl. Anal., **80** (12), 1951. (2001).
9. Djakov P. and Mityagin B.: *Unconditional convergence of spectral decompositions of 1D Dirac operators with regular boundary conditions*, Indiana Univ. Math. J., **61**(1), 359398.(2012).
10. Savchuk A.M. and Shkalikov A.A.: *The Dirac operator with complex-valued summable potential*, Math. Notes, **96**(5), 336 (2014).
11. Savchuk A. M., Sadovnichaya I.V.: *Riesz basis property with parentheses for Dirac system with integrable potential*, Sovrem. Math. Fundam. Napravl. **58**, 128152 (2015).
12. Lunyov A.A., Malamud M.M.: *On the Riesz basis property of root vectors system for  $2 \times 2$  Dirac-type operators*, Journal of Mathematical Analysis and Applications **441**(1), 57-103 (2016).

13. Kornev V.V., Khromov A.P.: *Dirac system with nondifferentiable potential and antiperiodic boundary conditions*, *Izv. Saratov Univ. Nov. Ser. Mat. Mekh. Inf.*, **13**(3), 2835 (2013).
14. Mykytnyk Ya.V. and Puyda D.V.: *Bari-Markus property of Dirac operators*, *Mat. Stud.*, **40**(2), 165171 (2013).
15. Abdullaev T.Sh., Nabiev I.M.: *An algorithm for reconstructing the Dirac operator with a spectral parameter in the boundary condition*, *Comput. Math. Math. Phys.*, **56**(2), 256267 (2016).
16. Mamedov Kh.R. and Akcay O.: *Inverse eigenvalue problem for a class of Dirac operators with discontinuous coefficient*, *Boundary Value Probl.*, **110** (2014). doi: 10.1186 / 1687-2770-2014-110.
17. Zygmund A.: *Trigonometricheskie ryady (Trigonometric Series)*, *Moscow: Mir*, **2** (1965).