

## Corrigendum to : ”Commutator of anisotropic maximal function with $BMO$ functions on total anisotropic Morrey spaces”

Mehriban N. Omarova, Gulnara A. Abasova\*

Received: 16.07.2024 / Revised: 09.08.2024 / Accepted: 14.08.2024

**Abstract.** *The purpose of this note is to correct an error in an earlier paper by the authors : Commutator of anisotropic maximal function with  $BMO$  functions on total anisotropic Morrey spaces, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci., Issue Mathematics.*

**Keywords.** total anisotropic Morrey spaces, anisotropic maximal operator, commutator,  $BMO$  spaces

**Mathematics Subject Classification (2010):** Primary 42B20, 42B25, 42B35

In [1] we study anisotropic maximal commutators  $M_b^d$  and commutators of the anisotropic maximal operator  $[b, M^d]$  in total anisotropic Morrey spaces  $L_{p,\lambda,\mu}^d(\mathbb{R}^n)$  when  $b$  belongs to  $BMO(\mathbb{R}^n)$ . The main goal of the paper [1] is to give necessary and sufficient conditions for the boundedness of the anisotropic maximal commutator  $M_b^d$  and the commutators of the anisotropic maximal operator  $[b, M^d]$  on  $L_{p,\lambda,\mu}^d(\mathbb{R}^n)$  when  $b$  belongs to  $BMO(\mathbb{R}^n)$ . New characterizations of some subclasses of  $BMO(\mathbb{R}^n)$  are obtained.

We assume that the reader is familiar with the contents and notation in the aforementioned paper [1].

In the paper [1] the author omitted the condition  $\mu \leq \lambda$  in inequalities (2.1), (2.2) and in Lemmas 2.1, 2.2. Otherwise, inequalities (2.1), (2.2) and the statements of Lemmas 2.1, 2.2 are false.

Therefore, the correct form of inequalities (2.1), (2.2) and Lemmas 2.1, 2.2 must be the following form.

$$L_{p,\lambda,\mu}^d(\mathbb{R}^n) \subset\supset L_{p,\lambda}^d(\mathbb{R}^n), \mu \leq \lambda \text{ and } \|f\|_{L_{p,\lambda}^d} \leq \|f\|_{L_{p,\lambda,\mu}^d},$$
$$L_{p,\lambda,\mu}^d(\mathbb{R}^n) \subset\supset L_{p,\mu}^d(\mathbb{R}^n), \mu \leq \lambda \text{ and } \|f\|_{L_{p,\mu}^d} \leq \|f\|_{L_{p,\lambda,\mu}^d}$$

**Lemma 0.1** *If  $0 < p < \infty$ ,  $0 \leq \mu \leq \lambda \leq |d|$ , then*

$$L_{p,\lambda,\mu}^d(\mathbb{R}^n) = L_{p,\lambda}^d(\mathbb{R}^n) \cap L_{p,\mu}^d(\mathbb{R}^n)$$

and

$$\|f\|_{L_{p,\lambda,\mu}^d(\mathbb{R}^n)} = \max \left\{ \|f\|_{L_{p,\lambda}^d}, \|f\|_{L_{p,\mu}^d} \right\}.$$

\* Corresponding author

M.N. Omarova  
Baku State University, Baku, Azerbaijan  
Institute of Mathematics and Mechanics, Baku, Azerbaijan  
E-mail: mehriban\_omarova@yahoo.com

G.A. Abasova  
Institute of Mathematics and Mechanics, Baku, Azerbaijan  
Azerbaijan State University of Economics, Baku, Azerbaijan  
E-mail: abasovag@yahoo.com

**Lemma 0.2** *If  $0 < p < \infty$ ,  $0 \leq \mu \leq \lambda \leq |d|$ , then*

$$WL_{p,\lambda,\mu}^d(\mathbb{R}^n) = WL_{p,\lambda}(\mathbb{R}^n) \cap WL_{p,\mu}(\mathbb{R}^n)$$

*and*

$$\|f\|_{WL_{p,\lambda,\mu}^d(\mathbb{R}^n)} = \max \left\{ \|f\|_{WL_{p,\lambda}^d}, \|f\|_{WL_{p,\mu}^d} \right\}.$$

## References

1. Omarova, M.N., Abasova, G.A.: *Commutator of anisotropic maximal function with BMO functions on total anisotropic Morrey spaces*, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci. 43 (1), Mathematics, 3-15 (2023).