

Commutator of Marcinkiewicz integral on total mixed Morrey spaces

Sahib A. Aliyev · Mubariz G. Hajibayov · Fatai A.
Isayev* · Ruhiyya O. Jafarova

Received: 12.10.2024 / Revised: 14.07.2025 / Accepted: 11.08.2025

Abstract. *In this paper, we study the boundedness of the Marcinkiewicz operator μ_Ω and its commutator $\mu_{b,\Omega}$ on total mixed Morrey spaces $L^{p,\lambda,\mu}(\mathbb{R}^n)$.*

Keywords. Total mixed Morrey spaces, Marcinkiewicz operator, commutators, BMO .

Mathematics Subject Classification (2010): 42B20, 42B25, 35J10

1 Introduction

In 1961, Benedek and Panzone [7] introduced Lebesgue spaces L^p with mixed norm over Euclidean spaces, which extend Lebesgue spaces and their related properties. In 1975, Bagby [6] investigated the boundedness of the Hardy-Littlewood maximal operator for functions taking values in spaces $l^p(\mathbb{R}^n)$. Since then, many papers focus various mixed norm spaces and the bounded properties of integral operators on spaces with mixed norm. In 2019, Nogayama [22, 23] considered a new Morrey space, with the L^p norm replaced by the mixed Lebesgue norm $L^{p,\lambda}(\mathbb{R}^n)$, which is called mixed Morrey spaces.

Classical Morrey spaces $L^{p,\lambda}$ were originally introduced by Morrey in [21] to study the local behavior of solutions of second-order elliptic partial differential equations. In 2022, Guliyev [12] introduced a variant of Morrey spaces called total Morrey spaces $L^{p,\lambda,\mu}(\mathbb{R}^n)$, $0 < p < \infty$, $\lambda \in \mathbb{R}$ and $\mu \in \mathbb{R}$. Total Morrey spaces generalize the classical Morrey spaces

* Corresponding author

Sa.A. Aliyev
Nakhchivan Teacher Institute, Department of Mathematics and Informatics, Nakhchivan, Azerbaijan
Nakhchivan State University, Department of General Mathematics, Nakhchivan, Azerbaijan
E-mail: sahib1960elm@gmail.com

M.G. Hajibayov
National Aviation Academy, Baku, Azerbaijan
Institute of Mathematics and Mechanics, Ministry of Science
E-mail: hajibayovm@yahoo.com

F.A. Isayev
Institute of Mathematics and Mechanics, Ministry of Science
and Educations of the Republic of Azerbaijan, Baku, Azerbaijan
E-mail: isayevfatai@yahoo.com

R.O. Jafarova
Nakhchivan State University, Department of General Mathematics, Nakhchivan, Azerbaijan
E-mail: cebr2012@mail.ru

$L^{p,\lambda}(\mathbb{R}^n)$ so that $L^{p,\lambda,\lambda}(\mathbb{R}^n) \equiv L^{p,\lambda}(\mathbb{R}^n)$ and the modified Morrey spaces $\tilde{L}^{p,\lambda}(\mathbb{R}^n)$ so that $L^{p,\lambda,0}(\mathbb{R}^n) = \tilde{L}^{p,\lambda}(\mathbb{R}^n)$. Necessary and sufficient conditions for the boundedness of the maximal commutator operator M_b and the commutator of the maximal operator $[b, M]$ on $L^{p,\lambda,\mu}(\mathbb{R}^n)$ when b belongs to the spaces $BMO(\mathbb{R}^n)$, are given in [12, Theorems 3 and 4], see also [9, 14–16, 24, 25].

In [16], the authors consider the total mixed Morrey spaces $L^{p,\lambda,\mu}(\mathbb{R}^n)$ introduced by Guliyev in [12] in the case $\mathbf{p} = (p, \dots, p)$. These spaces generalize mixed Morrey spaces so that $L^{p,\lambda,\lambda}(\mathbb{R}^n) \equiv L^{p,\lambda}(\mathbb{R}^n)$ and the modified mixed Morrey spaces so that $L^{p,\lambda,0}(\mathbb{R}^n) = \tilde{L}_{\mathbf{p},\lambda}(\mathbb{R}^n)$. The main properties of the spaces $L^{p,\lambda,\lambda}(\mathbb{R}^n)$ were presented and some embeddings into the Morrey space $L^{p,\lambda,\mu}(\mathbb{R}^n)$ are studied. Necessary and sufficient conditions for the boundedness of the maximal commutator operator M_b and the commutator of the maximal operator $[b, M]$ on $L^{p,\lambda,\mu}(\mathbb{R}^n)$ are also given. New characteristics for some subclasses of $BMO(\mathbb{R}^n)$ are obtained.

For any $r > 0$ and $x \in \mathbb{R}^n$, let $B(x, r) = \{y : |y - x| < r\}$ be the ball centered at x with radius r . Let $\mathcal{B} = \{B(x, r) : x \in \mathbb{R}^n, r > 0\}$ be the set of all such balls. We also use χ_E and $|E|$ to denote the characteristic function and the Lebesgue measure of a measurable set E . Let $S^{n-1} = \{x \in \mathbb{R}^n : |x| = 1\}$ be the unit sphere of \mathbb{R}^n ($n \geq 2$) equipped with the normalized Lebesgue measure. Suppose that Ω satisfies the following conditions.

(i) Ω is a homogeneous function of degree zero on \mathbb{R}^n . That is,

$$\Omega(tx) = \Omega(x) \quad (1.1)$$

for all $t > 0$ and $x \in \mathbb{R}^n$.

(ii) Ω has mean zero on S^{n-1} . That is,

$$\int_{S^{n-1}} \Omega(x') dx' = 0, \quad (1.2)$$

where $x' = x/|x|$ for any $x \neq 0$.

The Marcinkiewicz integral operator of higher dimension μ_Ω is defined by

$$\mu_\Omega(f)(x) = \left(\int_0^\infty |F_{\Omega,t}(f)(x)|^2 \frac{dt}{t^3} \right)^{1/2},$$

where

$$F_{\Omega,t}(f)(x) = \int_{B(x,t)} \frac{\Omega(x-y)}{|x-y|^{n-1}} f(y) dy.$$

It is well known that the Littlewood-Paley g -function is a very important tool in harmonic analysis and the Marcinkiewicz integral is essentially a Littlewood-Paley g -function. In this paper, we will consider the commutator $\mu_{\Omega,b}$ which is given by the following expression

$$\mu_{\Omega,b}f(x) = \left(\int_0^\infty |F_{\Omega,t}^b(x)|^2 \frac{dt}{t^3} \right)^{1/2},$$

where

$$F_{\Omega,t}^b(x) = \int_{B(x,t)} \frac{\Omega(x-y)}{|x-y|^{n-1}} [b(x) - b(y)] f(y) dy.$$

The study of Schrödinger operator $L = -\Delta + V$ recently attracted much attention. In particular, Shen [26] considered L_p estimates for Schrödinger operators L with certain potentials which include Schrödinger Riesz transforms $R_j^L = \frac{\partial}{\partial x_j} L^{-\frac{1}{2}}$, $j = 1, \dots, n$. Then, Dziubanński and Zienkiewicz [10] introduced the Hardy type space $H_L^1(\mathbb{R}^n)$ associated

with the Schrödinger operator L , which is larger than the classical Hardy space $H^1(\mathbb{R}^n)$, see also [2–5, 8, 13, 17, 18].

Similar to the classical Marcinkiewicz function, we define the Marcinkiewicz functions $\mu_{j,\Omega}$ associated with the Schrödinger operator L by

$$\mu_{j,\Omega}^L f(x) = \left(\int_0^\infty \left| \int_{B(x,t)} |\Omega(x-y)| K_j^L(x,y) f(y) dy \right|^2 \frac{dt}{t^3} \right)^{1/2},$$

where $K_j^L(x,y) = \widetilde{K_j^L}(x,y)|x-y|$ and $\widetilde{K_j^L}(x,y)$ is the kernel of $R_j = \frac{\partial}{\partial x_j} L^{-\frac{1}{2}}$, $j = 1, \dots, n$. In particular, when $V = 0$, $K_j^\Delta(x,y) = \widetilde{K_j^\Delta}(x,y)|x-y| = \frac{(x-y)_j/|x-y|}{|x-y|^{n-1}}$ and $\widetilde{K_j^\Delta}(x,y)$ is the kernel of $R_j = \frac{\partial}{\partial x_j} \Delta^{-\frac{1}{2}}$, $j = 1, \dots, n$. From now on, we will write $K_j(x,y) = K_j^\Delta(x,y)$ and

$$\mu_{j,\Omega} f(x) = \left(\int_0^\infty \left| \int_{B(x,t)} |\Omega(x-y)| K_j(x,y) f(y) dy \right|^2 \frac{dt}{t^3} \right)^{1/2}.$$

Obviously, $\mu_{j,\Omega} f$ are classical Marcinkiewicz functions with rough kernel. Therefore, it will be an interesting to study the properties of the operator $\mu_{j,\Omega}^L$.

The commutator of the classical Marcinkiewicz function with rough kernel is defined by

$$\mu_{j,\Omega,b} f(x) = \left(\int_0^\infty \left| \int_{B(x,t)} |\Omega(x-y)| K_j(x,y) [b(x) - b(y)] f(y) dy \right|^2 \frac{dt}{t^3} \right)^{1/2}.$$

The commutator $\mu_{j,\Omega,b}^L$ formed by $b \in BMO(\mathbb{R}^n)$ and the Marcinkiewicz function with rough kernel $\mu_{j,\Omega}^L$ is defined by

$$\mu_{j,\Omega,b}^L f(x) = \left(\int_0^\infty \left| \int_{B(x,t)} |\Omega(x-y)| K_j^L(x,y) [b(x) - b(y)] f(y) dy \right|^2 \frac{dt}{t^3} \right)^{1/2}.$$

The main goal of this paper is to show that Marcinkiewicz operators with rough kernel associated with the Schrödinger operators $\mu_{j,\Omega}^L$, $j = 1, \dots, n$, are bounded on the total mixed Morrey space $L^{\mathbf{p},\lambda,\mu}(\mathbb{R}^n)$, $1 < \mathbf{p} < \infty$, $0 \leq \lambda \leq n$, $0 \leq \mu \leq n$.

The well-known classical Hardy-Littlewood maximal operator M is defined by

$$Mf(x) = \sup_{r>0} \frac{1}{|B(x,r)|} \int_{B(x,r)} |f(y)| dy,$$

where $f \in L_{loc}^1(\mathbb{R}^n)$ and $|B(x,r)|$ is the Lebesgue measure of the ball $B(x,r)$. As we know, the Hardy-Littlewood maximal operator M is bounded on $L^{\mathbf{p}}(\mathbb{R}^n)$, $1 < \mathbf{p} < \infty$ (see [22, 23]), but there is no complete boundedness results for some other operators on the mixed Lebesgue spaces.

We find the conditions with $b \in BMO(\mathbb{R}^n)$ which ensures the boundedness of the operators $\mu_{j,\Omega,b}^L$, $j = 1, \dots, n$ on total mixed Morrey space $L^{\mathbf{p},\lambda,\mu}(\mathbb{R}^n)$, $1 < \mathbf{p} < \infty$, $0 \leq \lambda \leq n$, $0 \leq \mu \leq n$.

By $A \lesssim B$, we mean that $A \leq CB$ for some constant $C > 0$, and $A \approx B$ means that $A \lesssim B$ and $B \lesssim A$.

2 Definitions and preliminaries

We first recall the definition of mixed Lebesgue space defined in [7].

Let $\mathbf{p} = (p_1, \dots, p_n) \in (0, \infty]^n$. Then the mixed Lebesgue norm $\|\cdot\|_{L^{\mathbf{p}}}$ or $\|\cdot\|_{L^{(p_1, \dots, p_n)}}$ is defined by

$$\begin{aligned} \|f\|_{L^{\mathbf{p}}} &= \|f\|_{L^{(p_1, \dots, p_n)}} \\ &= \left(\int_{\mathbb{R}} \cdots \left(\int_{\mathbb{R}} \left(\int_{\mathbb{R}} |f(x_1, x_2, \dots, x_n)|^{p_1} dx_1 \right)^{\frac{p_2}{p_1}} dx_2 \right)^{\frac{p_3}{p_2}} \dots dx_n \right)^{\frac{1}{p_n}} \end{aligned}$$

where $f : \mathbb{R}^n \rightarrow \mathbb{C}$ is a measurable function. If $p_j = \infty$ for some $j = 1, n$, then we have to make appropriate modifications. We define the mixed Lebesgue space $L^{\mathbf{p}}(\mathbb{R}^n) = L^{(p_1, \dots, p_n)}(\mathbb{R}^n)$ to be the set of all locally integrable functions f with $\|f\|_{L^{\mathbf{p}}} < \infty$.

Definition 2.1 Let $0 < \mathbf{p} < \infty$, $\lambda \in \mathbb{R}$, $\mu \in \mathbb{R}$, $[t]_1 = \min\{1, t\}$, $t > 0$. We denote by $L^{\mathbf{p}, \lambda}(\mathbb{R}^n)$ the mixed Morrey space [23], by $\tilde{L}_{\mathbf{p}, \lambda}(\mathbb{R}^n)$ the modified mixed Morrey space [11], and by $L^{\mathbf{p}, \lambda, \mu}(\mathbb{R}^n)$ the total mixed Morrey space the set of all classes of locally integrable functions f with the finite norms

$$\begin{aligned} \|f\|_{L^{\mathbf{p}, \lambda}} &= \sup_{x \in \mathbb{R}^n, t > 0} t^{-\frac{\lambda}{n} \left(\sum_{i=1}^n \frac{1}{p_i} \right)} \|f\|_{L^{\mathbf{p}}(B(x, t))}, \\ \|f\|_{\tilde{L}_{\mathbf{p}, \lambda}} &= \sup_{x \in \mathbb{R}^n, t > 0} [t]_1^{-\frac{\lambda}{n} \left(\sum_{i=1}^n \frac{1}{p_i} \right)} \|f\|_{L^{\mathbf{p}}(B(x, t))}, \\ \|f\|_{L^{\mathbf{p}, \lambda, \mu}} &= \sup_{x \in \mathbb{R}^n, t > 0} [t]_1^{-\frac{\lambda}{n} \left(\sum_{i=1}^n \frac{1}{p_i} \right)} [1/t]_1^{\frac{\mu}{n} \left(\sum_{i=1}^n \frac{1}{p_i} \right)} \|f\|_{L^{\mathbf{p}}(B(x, t))}, \end{aligned}$$

respectively.

Definition 2.2 Let $0 < \mathbf{p} < \infty$, $\lambda \in \mathbb{R}$ and $\mu \in \mathbb{R}$. We define the weak mixed Morrey space $WL^{\mathbf{p}, \lambda}(\mathbb{R}^n)$ [23], the weak modified mixed Morrey space $W\tilde{L}_{\mathbf{p}, \lambda}(\mathbb{R}^n)$ [11] and the weak total mixed Morrey space $WL^{\mathbf{p}, \lambda, \mu}(\mathbb{R}^n)$ as the set of all locally integrable functions f with finite norms

$$\begin{aligned} \|f\|_{WL^{\mathbf{p}, \lambda}} &= \sup_{x \in \mathbb{R}^n, t > 0} t^{-\frac{\lambda}{n} \left(\sum_{i=1}^n \frac{1}{p_i} \right)} \|f\|_{WL^{\mathbf{p}}(B(x, t))}, \\ \|f\|_{W\tilde{L}_{\mathbf{p}, \lambda}} &= \sup_{x \in \mathbb{R}^n, t > 0} [t]_1^{-\frac{\lambda}{n} \left(\sum_{i=1}^n \frac{1}{p_i} \right)} \|f\|_{WL^{\mathbf{p}}(B(x, t))}, \\ \|f\|_{WL^{\mathbf{p}, \lambda, \mu}} &= \sup_{x \in \mathbb{R}^n, t > 0} [t]_1^{-\frac{\lambda}{n} \left(\sum_{i=1}^n \frac{1}{p_i} \right)} [1/t]_1^{\frac{\mu}{n} \left(\sum_{i=1}^n \frac{1}{p_i} \right)} \|f\|_{WL^{\mathbf{p}}(B(x, t))}, \end{aligned}$$

respectively.

Note that

$$\begin{aligned} L^{\mathbf{p}, 0, 0}(\mathbb{R}^n) &= \tilde{L}_{\mathbf{p}, 0}(\mathbb{R}^n) = L^{\mathbf{p}, 0}(\mathbb{R}^n) = L^{\mathbf{p}}(\mathbb{R}^n), \\ WL^{\mathbf{p}, 0, 0}(\mathbb{R}^n) &= W\tilde{L}_{\mathbf{p}, 0}(\mathbb{R}^n) = WL^{\mathbf{p}, 0}(\mathbb{R}^n) = WL^{\mathbf{p}}(\mathbb{R}^n), \\ L^{\mathbf{p}, \lambda, \lambda}(\mathbb{R}^n) &= L^{\mathbf{p}, \lambda}(\mathbb{R}^n), \quad L^{\mathbf{p}, \lambda, 0}(\mathbb{R}^n) = \tilde{L}_{\mathbf{p}, \lambda}(\mathbb{R}^n), \\ \|f\|_{WL^{\mathbf{p}, \lambda, \mu}} &\leq \|f\|_{L^{\mathbf{p}, \lambda, \mu}} \quad \text{and therefore} \quad L^{\mathbf{p}, \lambda, \mu}(\mathbb{R}^n) \subset WL^{\mathbf{p}, \lambda, \mu}(\mathbb{R}^n) \end{aligned}$$

and

$$L^{\mathbf{p},\lambda,\mu}(\mathbb{R}^n) \subset_{\succ} L^{\mathbf{p},\lambda}(\mathbb{R}^n), \mu \leq \lambda \text{ and } \|f\|_{L^{\mathbf{p},\lambda}} \leq \|f\|_{L^{\mathbf{p},\lambda,\mu}},$$

$$L^{\mathbf{p},\lambda,\mu}(\mathbb{R}^n) \subset_{\succ} L^{\mathbf{p},\mu}(\mathbb{R}^n), \mu \leq \lambda \text{ and } \|f\|_{L^{\mathbf{p},\mu}} \leq \|f\|_{L^{\mathbf{p},\lambda,\mu}}$$

$$\tilde{L}_{\mathbf{p},\lambda}(\mathbb{R}^n) \subset_{\succ} L^{\mathbf{p}}(\mathbb{R}^n) \text{ and } \|f\|_{L^{\mathbf{p}}} \leq \|f\|_{\tilde{L}_{\mathbf{p},\lambda}}$$

and if $\lambda < 0$ or $\lambda > n$, then $L^{\mathbf{p},\lambda}(\mathbb{R}^n) = \tilde{L}_{\mathbf{p},\lambda}(\mathbb{R}^n) = WL^{\mathbf{p},\lambda}(\mathbb{R}^n) = W\tilde{L}_{\mathbf{p},\lambda}(\mathbb{R}^n) = \Theta$, where $\Theta \equiv \Theta(\mathbb{R}^n)$ is the set of all functions equivalent to 0 on \mathbb{R}^n .

Lemma 2.1 [16] *If $0 < \mathbf{p} < \infty$, $0 \leq \mu \leq \lambda \leq n$, then*

$$L^{\mathbf{p},\lambda,\mu}(\mathbb{R}^n) = L^{\mathbf{p},\lambda}(\mathbb{R}^n) \cap L^{\mathbf{p},\mu}(\mathbb{R}^n)$$

and

$$\|f\|_{L^{\mathbf{p},\lambda,\mu}(\mathbb{R}^n)} = \max \left\{ \|f\|_{L^{\mathbf{p},\lambda}(\mathbb{R}^n)}, \|f\|_{L^{\mathbf{p},\mu}(\mathbb{R}^n)} \right\}.$$

Lemma 2.2 [16] *If $0 < \mathbf{p} < \infty$, $0 \leq \mu \leq \lambda \leq n$, then*

$$WL^{\mathbf{p},\lambda,\mu}(\mathbb{R}^n) = WL^{\mathbf{p},\lambda}(\mathbb{R}^n) \cap WL^{\mathbf{p},\mu}(\mathbb{R}^n)$$

and

$$\|f\|_{WL^{\mathbf{p},\lambda,\mu}(\mathbb{R}^n)} = \max \left\{ \|f\|_{WL^{\mathbf{p},\lambda}(\mathbb{R}^n)}, \|f\|_{WL^{\mathbf{p},\mu}(\mathbb{R}^n)} \right\}.$$

Remark 2.1 If $0 < \mathbf{p} < \infty$, and $\mu < 0$ or $\lambda > n$, then

$$L^{\mathbf{p},\lambda,\mu}(\mathbb{R}^n) = WL^{\mathbf{p},\lambda,\mu}(\mathbb{R}^n) = \Theta(\mathbb{R}^n).$$

3 Marcinkiewicz operator μ_{Ω} in total mixed Morrey spaces

In this section, we investigate the boundedness of Marcinkiewicz operator μ_{Ω} satisfies the conditions (1.1), (1.2) and $\Omega \in L^{\infty}(S^{n-1})$ on the total mixed Morrey space $L^{\mathbf{p},\lambda,\mu}$.

The following lemma gives us explicit estimates for the $L^{\mathbf{p}}(\mathbb{R}^n)$ norm of μ_{Ω} on a given ball $B(x_0, r)$.

Lemma 3.1 [1, Lemma 3.1] *Let Ω be satisfies the conditions (1.1), (1.2) and $\Omega \in L^{\infty}(S^{n-1})$.*

Then for $1 < \mathbf{p} < \infty$, the inequality

$$\|\mu_{\Omega}f\|_{L^{\mathbf{p}}(B(x_0,r))} \lesssim r^{\sum_{i=1}^n \frac{1}{p_i}} \int_{2r}^{\infty} t^{-1-\sum_{i=1}^n \frac{1}{p_i}} \|f\|_{L^{\mathbf{p}}(B(x_0,t))} dt \quad (3.1)$$

holds for any ball $B(x_0, r)$ and all $f \in L_{loc}^{\mathbf{p}}(\mathbb{R}^n)$.

Now we can present the first main result in this section.

Theorem 3.1 *Let Ω be satisfies the conditions (1.1), (1.2) and $\Omega \in L^{\infty}(S^{n-1})$. Let also $1 < \mathbf{p} < \infty$, $0 \leq \lambda \leq n$ and $0 \leq \mu \leq n$. Then the operator μ_{Ω} is bounded on $L^{\mathbf{p},\lambda,\mu}$. Moreover,*

$$\|\mu_{\Omega}f\|_{L^{\mathbf{p},\lambda,\mu}} \leq \|f\|_{L^{\mathbf{p},\lambda,\mu}}.$$

Proof. From the inequality (3.1) we get

$$\begin{aligned}
\|\mu_\Omega f\|_{L^{\mathbf{p},\lambda,\mu}} &= \sup_{x \in \mathbb{R}^n, r > 0} [r]_1^{-\frac{\lambda}{n}(\sum_{i=1}^n \frac{1}{p_i})} [1/r]_1^{\frac{\mu}{n}(\sum_{i=1}^n \frac{1}{p_i})} \|\mu_\Omega f\|_{L^{\mathbf{p}}(B(x,r))} \\
&\lesssim \sup_{x \in \mathbb{R}^n, r > 0} [r]_1^{-\frac{\lambda}{n}(\sum_{i=1}^n \frac{1}{p_i})} [1/r]_1^{\frac{\mu}{n}(\sum_{i=1}^n \frac{1}{p_i})} r^{\sum_{i=1}^n \frac{1}{p_i}} \int_{2r}^\infty t^{-1-\sum_{i=1}^n \frac{1}{p_i}} \|f\|_{L^{\mathbf{p}}(B(x_0,t))} dt \\
&\lesssim \|f\|_{L^{\mathbf{p},\lambda,\mu}} \sup_{x \in \mathbb{R}^n, r > 0} [r]_1^{-\frac{\lambda}{n}(\sum_{i=1}^n \frac{1}{p_i})} [1/r]_1^{\frac{\mu}{n}(\sum_{i=1}^n \frac{1}{p_i})} r^{\sum_{i=1}^n \frac{1}{p_i}} \\
&\quad \times \int_r^\infty t^{-\sum_{i=1}^n \frac{1}{p_i}} [t]_1^{\frac{\lambda}{n}(\sum_{i=1}^n \frac{1}{p_i})} [1/t]_1^{-\frac{\mu}{n}(\sum_{i=1}^n \frac{1}{p_i})} \frac{dt}{t} \\
&\lesssim \|f\|_{L^{\mathbf{p},\lambda,\mu}} \sup_{x \in \mathbb{R}^n, r > 0} [r]_1^{\frac{n-\lambda}{n}(\sum_{i=1}^n \frac{1}{p_i})} [1/r]_1^{-\frac{n-\mu}{n}(\sum_{i=1}^n \frac{1}{p_i})} \\
&\quad \times \int_r^\infty [t]_1^{-\frac{n-\lambda}{n}(\sum_{i=1}^n \frac{1}{p_i})} [1/t]_1^{\frac{n-\mu}{n}(\sum_{i=1}^n \frac{1}{p_i})} \frac{dt}{t} \\
&= \|f\|_{L^{\mathbf{p},\lambda,\mu}} \sup_{x \in \mathbb{R}^n, r > 0} \int_1^\infty [t]_1^{-\frac{n-\lambda}{n}(\sum_{i=1}^n \frac{1}{p_i})} [1/t]_1^{\frac{n-\mu}{n}(\sum_{i=1}^n \frac{1}{p_i})} \frac{dt}{t} \\
&\lesssim \|f\|_{L^{\mathbf{p},\lambda,\mu}}.
\end{aligned}$$

Thus the proof of the theorem is completed.

Note that if we take $\mathbf{p} = (p, \dots, p)$ in Theorem 3.1, we obtain the boundedness of μ_Ω on the total Morrey spaces.

4 Commutator of Marcinkiewicz operator $\mu_{\Omega,b}$ in total mixed Morrey spaces

In this section, we investigate the boundedness of commutator of Marcinkiewicz operator $\mu_{\Omega,b}$ satisfies the conditions (1.1), (1.2) and $\Omega \in L^\infty(S^{n-1})$ on the total mixed Morrey space $L^{\mathbf{p},\lambda,\mu}$. First, we review the definition of $BMO(\mathbb{R}^n)$, the bounded mean oscillation space. A function $f \in L^1_{loc}(\mathbb{R}^n)$ belongs to the bounded mean oscillation space $BMO(\mathbb{R}^n)$ if

$$\|f\|_{BMO} = \sup_{x \in \mathbb{R}^n, r > 0} \frac{1}{|B(x,r)|} \int_{B(x,r)} |f(y) - f_{B(x,r)}| dy < \infty. \quad (4.1)$$

If one regards two functions whose difference is a constant as one, then the space $BMO(\mathbb{R}^n)$ is a Banach space with respect to norm $\|\cdot\|_{BMO}$. The John-Nirenberg inequality for BMO yields that for any $1 < q < \infty$ and $f \in BMO(\mathbb{R}^n)$, the BMO norm of f is equivalent to

$$\|f\|_{BMO^q} = \sup_{x \in \mathbb{R}^n, r > 0} \left(\frac{1}{|B(x,r)|} \int_{B(x,r)} |f(y) - f_{B(x,r)}|^q dy \right)^{\frac{1}{q}}$$

Recall that for any $\mathbf{p} = (p_1, \dots, p_n) \in (1, \infty)^n$, the John-Nirenberg inequality for mixed norm space [19, 20] shows that the BMO norm of all $f \in BMO(\mathbb{R}^n)$ is also equivalent to

$$\|f\|_{BMO^{\mathbf{p}}} = \sup_{x \in \mathbb{R}^n, r > 0} \frac{\|(f - f_{B(x,r)})\chi_{B(x,r)}\|_{L^{\mathbf{p}}}}{\|\chi_{B(x,r)}\|_{L^{\mathbf{p}}}}. \quad (4.2)$$

The following property for BMO functions is valid.

Lemma 4.1 *Let $f \in BMO(\mathbb{R}^n)$. Then for all $0 < 2r < t$, we have*

$$|f_{B(x,r)} - f_{B(x,t)}| \lesssim \|f\|_{BMO} \ln \frac{t}{r}. \quad (4.3)$$

The following lemma gives us explicit estimates for the $L^{\mathbf{p}}(\mathbb{R}^n)$ norm of $\mu_{\Omega,b}$ on a given ball $B(x_0, r)$.

Lemma 4.2 *[1, Lemma 4.2] Let Ω be satisfy the conditions (1.1), (1.2) and $\Omega \in L^\infty(S^{n-1})$. Let also $1 < \mathbf{p} < \infty$ and $b \in BMO(\mathbb{R}^n)$. Then the inequality*

$$\begin{aligned} & \|\mu_{\Omega,b} f\|_{L^{\mathbf{p}}(B(x_0,r))} \\ & \lesssim \|b\|_{BMO} r^{\sum_{i=1}^n \frac{1}{p_i}} \int_{2r}^{\infty} \left(1 + \ln \frac{t}{r}\right) t^{-1 - \sum_{i=1}^n \frac{1}{p_i}} \|f\|_{L^{\mathbf{p}}(B(x_0,t))} dt \end{aligned} \quad (4.4)$$

holds for any ball $B(x_0, r)$ and all $f \in L_{loc}^{\mathbf{p}}(\mathbb{R}^n)$.

Now we give the boundedness of $\mu_{\Omega,b}$ on the total mixed Morrey space.

Theorem 4.1 *Let Ω be satisfy the conditions (1.1), (1.2) and $\Omega \in L^\infty(S^{n-1})$. Let also $1 < \mathbf{p} < \infty$, $b \in BMO(\mathbb{R}^n)$, $0 \leq \lambda \leq n$ and $0 \leq \mu \leq n$. Then the operator $\mu_{\Omega,b}$ is bounded on $L^{\mathbf{p},\lambda,\mu}$. Moreover,*

$$\|\mu_{\Omega,b} f\|_{L^{\mathbf{p},\lambda,\mu}} \leq \|b\|_{BMO} \|f\|_{L^{\mathbf{p},\lambda,\mu}}.$$

Proof. From the inequality (4.4) we get

$$\begin{aligned} \|\mu_{\Omega,b} f\|_{L^{\mathbf{p},\lambda,\mu}} &= \sup_{x \in \mathbb{R}^n, r > 0} [r]_1^{-\frac{\lambda}{n} \left(\sum_{i=1}^n \frac{1}{p_i}\right)} [1/r]_1^{\frac{\mu}{n} \left(\sum_{i=1}^n \frac{1}{p_i}\right)} \|\mu_{\Omega,b} f\|_{L^{\mathbf{p}}(B(x,r))} \\ &\lesssim \|b\|_{BMO} \sup_{x \in \mathbb{R}^n, r > 0} [r]_1^{-\frac{\lambda}{n} \left(\sum_{i=1}^n \frac{1}{p_i}\right)} [1/r]_1^{\frac{\mu}{n} \left(\sum_{i=1}^n \frac{1}{p_i}\right)} r^{\sum_{i=1}^n \frac{1}{p_i}} \\ &\quad \times \int_{2r}^{\infty} \left(1 + \ln \frac{t}{r}\right) t^{-1 - \sum_{i=1}^n \frac{1}{p_i}} \|f\|_{L^{\mathbf{p}}(B(x_0,t))} dt \\ &\lesssim \|b\|_{BMO} \|f\|_{L^{\mathbf{p},\lambda,\mu}} \sup_{x \in \mathbb{R}^n, r > 0} [r]_1^{-\frac{\lambda}{n} \left(\sum_{i=1}^n \frac{1}{p_i}\right)} [1/r]_1^{\frac{\mu}{n} \left(\sum_{i=1}^n \frac{1}{p_i}\right)} r^{\sum_{i=1}^n \frac{1}{p_i}} \\ &\quad \times \int_r^{\infty} \left(1 + \ln \frac{t}{r}\right) t^{-\sum_{i=1}^n \frac{1}{p_i}} [t]_1^{\frac{\lambda}{n} \left(\sum_{i=1}^n \frac{1}{p_i}\right)} [1/t]_1^{-\frac{\mu}{n} \left(\sum_{i=1}^n \frac{1}{p_i}\right)} \frac{dt}{t} \\ &\lesssim \|b\|_{BMO} \|f\|_{L^{\mathbf{p},\lambda,\mu}} \sup_{x \in \mathbb{R}^n, r > 0} [r]_1^{\frac{n-\lambda}{n} \left(\sum_{i=1}^n \frac{1}{p_i}\right)} [1/r]_1^{-\frac{n-\mu}{n} \left(\sum_{i=1}^n \frac{1}{p_i}\right)} \\ &\quad \times \int_r^{\infty} \left(1 + \ln \frac{t}{r}\right) [t]_1^{-\frac{n-\lambda}{n} \left(\sum_{i=1}^n \frac{1}{p_i}\right)} [1/t]_1^{\frac{n-\mu}{n} \left(\sum_{i=1}^n \frac{1}{p_i}\right)} \frac{dt}{t} \\ &= \|b\|_{BMO} \|f\|_{L^{\mathbf{p},\lambda,\mu}} \sup_{x \in \mathbb{R}^n, r > 0} \int_1^{\infty} (1 + \ln t) [t]_1^{-\frac{n-\lambda}{n} \left(\sum_{i=1}^n \frac{1}{p_i}\right)} [1/t]_1^{\frac{n-\mu}{n} \left(\sum_{i=1}^n \frac{1}{p_i}\right)} \frac{dt}{t} \\ &\lesssim \|b\|_{BMO} \|f\|_{L^{\mathbf{p},\lambda,\mu}}. \end{aligned}$$

Thus the proof of the theorem is completed.

By taking $\mathbf{p} = (p, \dots, p)$ in Theorem 4.1, we obtain the boundedness of $\mu_{\Omega,b}$ on the total Morrey spaces.

5 Marcinkiewicz operators with rough kernel associated with the Schrödinger operators $\mu_{j,\Omega}^L$ and its commutator $\mu_{j,\Omega,b}^L$ in total mixed Morrey spaces

Let us consider the Schrödinger operator

$$L = -\Delta + V \text{ on } \mathbb{R}^n, \quad n \geq 3,$$

where V is a non-negative, $V \neq 0$, and belongs to the reverse Hölder class B_q for some $q \geq n/2$, i.e., there exists a constant $C > 0$ such that the reverse Hölder inequality

$$\left(\frac{1}{|B(x,r)|} \int_{B(x,r)} V^q(y) dy \right)^{1/q} \leq \frac{C}{|B(x,r)|} \int_{B(x,r)} V(y) dy$$

holds for every $x \in \mathbb{R}^n$ and $0 < r < 1$. In particular, if V is a nonnegative polynomial, then $V \in B_1$.

Obviously, $B_{q_2} \subset B_{q_1}$, if $q_2 > q_1$. The most important property of the class B_q is its self-improvement, that is, if $V \in B_q$, then $V \in B_{q+\epsilon}$ for some $\epsilon > 0$.

In this section, we prove the boundedness of the Marcinkiewicz operators with rough kernel associated with the Schrödinger operators $\mu_{j,\Omega}^L$ and its commutator $\mu_{j,\Omega,b}^L$ on total mixed Morrey space $L^{\mathbf{p},\lambda,\mu}$.

For $x \in \mathbb{R}^n$, the function $\rho(x)$ is defined by

$$\rho(x) = \sup_{r>0} \left\{ r : \frac{1}{r^{n-2}} \int_{B(x,r)} V(y) dy \leq 1 \right\}.$$

Lemma 5.1 [26] *Let $V \in B_q$ with $q \geq n/2$. Then there exists $l_0 > 0$ such that*

$$\frac{l}{C} \left(1 + \frac{|x-y|}{\rho(x)} \right)^{-l_0} \leq \frac{\rho(y)}{\rho(x)} \leq C \left(1 + \frac{|x-y|}{\rho(x)} \right)^{l_0/(l_0+1)}.$$

In particular, $\rho(x) \sim \rho(y)$ if $|x-y| < C\rho(x)$.

Lemma 5.2 [26] *Let $V \in B_q$ with $q \geq n/2$. For any $l > 0$, there exists $C_l > 0$ such that*

$$\left| K_j^L(x, y) \right| \leq \frac{C_l}{\left(1 + \frac{|x-y|}{\rho(x)} \right)^l} \frac{1}{|x-y|^{n-1}},$$

and

$$\left| K_j^L(x, y) - K_j(x-y) \right| \leq C \frac{\rho(x)}{|x-y|^{n-2}}.$$

Analogously proof of Lemma 3.1 and Theorem 3.1 the following results is valid.

Lemma 5.3 *Let Ω be satisfy the conditions (1.1), (1.2), $\Omega \in L^\infty(S^{n-1})$ and $V \in B_n$. Then for $1 < \mathbf{p} < \infty$, the inequality*

$$\|\mu_{j,\Omega}^L f\|_{L^{\mathbf{p}}(B(x_0,r))} \lesssim r^{\sum_{i=1}^n \frac{1}{p_i}} \int_{2r}^\infty t^{-1-\sum_{i=1}^n \frac{1}{p_i}} \|f\|_{L^{\mathbf{p}}(B(x_0,t))} dt$$

holds for any ball $B(x_0, r)$ and all $f \in L_{loc}^{\mathbf{p}}(\mathbb{R}^n)$.

Theorem 5.1 Let Ω be satisfy the conditions (1.1), (1.2), $\Omega \in L^\infty(S^{n-1})$ and $V \in B_n$. Let also $1 < \mathbf{p} < \infty$, $0 \leq \lambda \leq n$ and $0 \leq \mu \leq n$. Then the operator $\mu_{j,\Omega}^L$ is bounded on $L^{\mathbf{p},\lambda,\mu}$. Moreover,

$$\|\mu_{j,\Omega}^L f\|_{L^{\mathbf{p},\lambda,\mu}} \leq \|f\|_{L^{\mathbf{p},\lambda,\mu}}.$$

Analogously proof of Lemma 4.2 and Theorem 4.1 the following results is valid.

Lemma 5.4 Let Ω be satisfy the conditions (1.1), (1.2), $\Omega \in L^\infty(S^{n-1})$ and $V \in B_n$. Then for $1 < \mathbf{p} < \infty$ and $b \in BMO(\mathbb{R}^n)$, the inequality

$$\|\mu_{j,\Omega,b}^L f\|_{L^{\mathbf{p}}(B(x_0,r))} \lesssim \|b\|_{BMO} r^{\sum_{i=1}^n \frac{1}{p_i}} \int_{2r}^{\infty} \left(1 + \ln \frac{t}{r}\right) t^{-1 - \sum_{i=1}^n \frac{1}{p_i}} \|f\|_{L^{\mathbf{p}}(B(x_0,t))} dt$$

holds for any ball $B(x_0, r)$ and all $f \in L_{loc}^{\mathbf{p}}(\mathbb{R}^n)$.

Theorem 5.2 Let Ω be satisfy the conditions (1.1), (1.2), $\Omega \in L^\infty(S^{n-1})$ and $V \in B_n$. Let also $1 < \mathbf{p} < \infty$, $b \in BMO(\mathbb{R}^n)$, $0 \leq \lambda \leq n$ and $0 \leq \mu \leq n$. Then the operator $\mu_{j,\Omega,b}^L$ is bounded on $L^{\mathbf{p},\lambda,\mu}$. Moreover,

$$\|\mu_{j,\Omega,b}^L f\|_{L^{\mathbf{p},\lambda,\mu}} \leq \|b\|_{BMO} \|f\|_{L^{\mathbf{p},\lambda,\mu}}.$$

Acknowledgements. The authors would like to express their gratitude to the referees for his very valuable comments and suggestions.

References

1. Akbarov, A.A., Isayev, F.A., Ismayilov, M.I.: *Marcinkiewicz integral and its commutator on mixed Morrey spaces*, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci **45**(1) Mathematics, 3-16 (2025).
2. Akbulut, A., Celik, S., Omarova, M.N.: *Fractional maximal operator associated with Schrödinger operator and its commutators on vanishing generalized Morrey spaces*, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci **44**(1) Mathematics, 3-19 (2024).
3. Akbulut, A., Guliyev, R., Ekincioglu, I.: *Calderon-Zygmund operators associated with Schrödinger operator and their commutators on vanishing generalized Morrey spaces*, TWMS J. Pure Appl. Math. **13**(2), 144-157 (2022).
4. Akbulut, A., Kuzu, O.: *Marcinkiewicz integrals associated with Schrödinger operator on generalized Morrey spaces*, J. Math. Inequal. **8**(4), 791-801 (2014).
5. Akbulut, A., Omarova, M.N., Serbetci, A.: *Generalized local mixed Morrey estimates for linear elliptic systems with discontinuous coefficients*, Socar Proceedings No. 1, 136-142 (2025).
6. Bagby, R.L.: *An extended inequality for the maximal function*, Proc. Amer. Math. Soc. **48**(2), 419-422 (1975).
7. Benedek, A., Panzone, R.: *The spaces L^p with mixed norm*, Duke Math. J. **28**(3), 301-324 (1961).
8. Celik, S., Guliyev, V.S., Akbulut, A.: *Commutator of fractional integral with Lipschitz functions associated with Schrödinger operator on local generalized mixed Morrey spaces*, Open Math. **22**, 20240082 (2024).
9. Celik, S., Akbulut, A., Omarova, M.N.: *Characterizations of anisotropic Lipschitz functions via the commutators of anisotropic maximal function in total anisotropic Morrey spaces*, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci **45**(1) Mathematics, 25-37 (2025).

10. Dziubański, J., Zienkiewicz, J.: *Hardy space H^1 associated to Schrödinger operator with potential satisfying reverse Hölder inequality*, Rev. Mat. Iber. **15**, 279-296 (1999).
11. V.S. Guliyev, J.J. Hasanov, Y. Zeren, *Necessary and sufficient conditions for the boundedness of the Riesz potential in modified Morrey spaces*, J. Math. Inequal. **5** (2011), no. 4, 491-506.
12. Guliyev, V.S. : *Maximal commutator and commutator of maximal function on total Morrey spaces*, J. Math. Inequal. **16**(4), 15091524 (2022).
13. Guliyev, V.S., Akbulut, A., Celik, S.: *Fractional integral related to Schrödinger operator on vanishing generalized mixed Morrey spaces*, Bound. Value Probl. (2024), Article number: 137 (2024).
14. Guliyev, V.S., Isayev, F.A., Serbetci, A.: *Boundedness of the anisotropic fractional maximal operator in total anisotropic Morrey spaces*, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci. Math. **44**(1) Mathematics, 41-50 (2024).
15. Guliyev, V.S. : *Characterizations of commutators of the maximal function in total Morrey spaces on stratified Lie groups*, Anal. Math. Phys. **15**:42 (2025).
16. Guliyev, V.S., Akbulut, A., Isayev, F.A., Serbetci, A.: *Commutators of maximal function with BMO functions on total mixed Morrey spaces*, Journal of Contemporary Applied Mathematics **16**(1), 1-15 (2026).
17. Hasanov, A., Hasanov, S.G., Nazkipinar, A.: *Marcinkiewicz integral with rough kernel in local Morrey-type spaces*, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci. **43**(4) Mathematics, 96-104 (2023).
18. Hamzayev, V.H., Mammadov, Y.Y.: *Commutators of Marcinkiewicz integral with rough kernels on generalized weighted Morrey spaces*, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci. **43**(1) Mathematics, 55-65 (2023).
19. Ho, K.P.: *Strong maximal operator on mixed-norm spaces*, Ann. Univ. Ferrara, **62**(2), 275-291 (2016).
20. Ho, K.P.: *Mixed norm lebesgue spaces with variable exponents and applications*, Riv. Mat. Univ. Parma **9**(1), 21-44 (2018).
21. Morrey, C.B.: *On the solutions of quasi-linear elliptic partial differential equations*, Trans. Amer. Math. Soc. **43**(1), 126-166 (1938).
22. Nogayama, T.: *Boundedness of commutators of fractional integral operators on mixed Morrey spaces*, Integral Transforms Spec. Funct. **30**(10), 790-816 (2019).
23. Nogayama, T.: *Mixed Morrey spaces*, Positivity **23**(4), 961-1000 (2019).
24. Omarova, M.N. : *Commutators of parabolic fractional maximal operators on parabolic total Morrey spaces*, Math. Meth. Appl. Sci. **48**(11), 11037-11044 (2025).
25. Omarova, M.N. : *Commutators of anisotropic maximal operators with BMO functions on anisotropic total Morrey spaces*, Azerb. J. Math. **15**(2), 150-162 (2025).
26. Shen, Z.: *L^p estimates for Schrödinger operators with certain potentials*, Ann. Inst. Fourier (Grenoble) **45**, 513-546 (1995).