

Riesz property criterion for the system of eigen and associated functions of second order ordinary differential operator

Vali M. Kurbanov*, Kh.R. Gojayeve

Received: 22.02.2025 / Revised: 15.07.2025 / Accepted: 06.08.2025

Abstract. *On the interval $G = (0, 1)$ an ordinary differential operator of second order with summable coefficients is considered. Riesz property for the system of eigen and associated functions of the given operator is studied. The criterion of Riesz property in $L_p(G)$, $1 < p \leq 2$, of eigen and associated functions system is established.*

Keywords. differential operator · eigen and associated functions · Riesz property criterion.

Mathematics Subject Classification (2010): 34L10, 42A20

1 Introduction and statement of main result

On the interval $G = (0, 1)$ consider the ordinary differential operator

$$Lu = u'' + q_1(x)u' + q_2(x)u,$$

with summable coefficients $q_i(x) \in L_1(G)$, $i = 1, 2$.

The root functions (i.e. eigenfunctions and associated function) of the operator L are understood in the generalized sense (irrespective of the boundary conditions) [3].

Consider an arbitrary system $\{u_k(x)\}_{k=1}^{\infty}$ consisting of eigen and associated functions (root functions) of the operator L . Let $\{\lambda_k\}_{k=1}^{\infty}$ be the corresponding system of eigenvalues of this operator. We require that, together with each associated function of order s , $s \geq 1$, the system $\{u_k(x)\}_{k=1}^{\infty}$ also contains the corresponding eigenfunction and all associated functions of order less than s . This means that each element of the system $\{u_k(x)\}_{k=1}^{\infty}$ is identically non-zero, is absolutely continuous together with its first order derivative on $\overline{G}$, and almost everywhere in G satisfies the equation $Lu_k + \lambda_k u_k = \theta_k u_{k-1}$, where θ_k equals 0 (in this case $u_k(x)$ is an eigenfunction), or 1 (in this case $u_k(x)$ is an associated functions

* Corresponding author

V.M.Kurbanov
Ministry of Science and Education, Institute of Mathematics and Mechanics;
Azerbaijan State Pedagogical University, Baku, Azerbaijan ;
E-mail: q.vali@yahoo.com

Kh.R.Gojayeve
Azerbaijan State Pedagogical University, Baku, Azerbaijan
E-mail: mehdizade.xedice@gmail.com

of order $r \geq 1$, and $u_{k-1}(x)$ is an associated functions of order $r - 1$, $\lambda_k = \lambda_{k-1} = \dots = \lambda_{k-r}$, $\theta_k = \theta_{k-1} = \theta_{k-r+1} = 1$ and $\theta_{k-r} = 0$).

The highest order of root functions corresponding to the given eigenfunction will be called the rank of this eigenfunction.

In such a generalized understanding of eigen and associated functions (root functions) Ilin V.A. (see [3]) first established the Bessel property unconditional basic property criterion in $L_2(G)$ of the system of root functions of the operator L for $q_1(x) \equiv 0$, $q_2(x) \in L_1(G)$.

In the papers (see [2],[4],[6-9], [16]) these problems were studied for a higher order ordinary differential operator. This time, for Bessel property it was always assumed that $q_1(x) \in L_2(G)$, $q_2(x) \in L_1(G)$, $f(x) \in L_2(G)$, for Riesz property it was always assumed that $q_1(x) \in L_p(G)$, $1 < p < 2$, $f(x) \in L_p(G)$.

It should be note that the Riess inequality (in particular, the Bessel inequality) is widely used to study the convergence of spectral expansions of functions (see [1-4], [9-16]).

In the present paper we study the Riesz property (in case $p = 2$ Bessel property) for the system of root functions of the given operator with the coefficients $q_i(x) \in L_1(G)$, $i = 1, 2$, and the criterion of Riesz property of root functions system is established.

Definition 1.1 A system $\{\varphi_k(x)\}_{k=1}^{\infty} \subset L_q(G)$ is called Riesz system if there exists a constant $M = M(p)$ such that for each function $f(x) \in L_p(G)$, $1 < p \leq 2$, $q = p/(p-1)$, one has the inequality

$$\left(\sum_{k=1}^{\infty} |(f, \varphi_k)|^q \right)^{1/q} \leq M \|f\|_p.$$

Denote $\mu_k = \sqrt{\lambda_k}$, $Re\mu_k \geq 0$.

The main result of this work is the following theorem.

Theorem (Riesz property criterion). Let $q_i(x) \in L_1(G)$, $i = 1, 2$, the ranks of eigenfunctions be uniformly bounded, and assume that there exists a constant C_0 such that

$$|Im\mu_k| \leq C_0, \quad k = 1, 2, \dots \quad (1.1)$$

Then a necessary and sufficient condition for the Riesz property of the system $\left\{ u_k(x) \|u_k\|_q^{-1} \right\}_{k=1}^{\infty}$ in $L_p(G)$, $1 < p \leq 2$, $q = p/(p-1)$ is that there a constant M_1 such that

$$\sum_{\tau \leq Re\mu_k \leq \tau+1} 1 \leq M_1, \quad \forall \tau \geq 0. \quad (1.2)$$

Remark 1.1 In the sufficient part of the Theorem, the uniform boundedness of the ranks of eigenfunctions holds automatically as a consequence of inequality (1.2).

2 Proof of the Theorem

Necessity: We fix an arbitrary number $t \geq 0$. Introduce following set

$$\Omega_{\tau} = \{k \in N : \tau \leq Re\mu_k \leq \tau + 1, |Im\mu_k| \leq C_0, |\mu_k| \geq 1\},$$

and number

$$R_0 = (n_0(1 + C_0)^{1+1/q})^{-1},$$

where $n_0 \geq 1$ is chosen so that $R_0 \leq 1/4$ and for any set $E \subset \overline{G}$, $mes E \leq 2R_0$, the following is fulfilled:

$$\omega(R_0) = \sup_{E \subset \overline{G}} \left\{ \|q_i\|_{1,E}, \quad i = 1, 2 \right\} \leq N_0^{-1}, \quad (2.1)$$

where $\|q_i\|_{1,E} = \int_E |q_i(x)| dx$, $(\|\cdot\|_1 = \|\cdot\|_{1,G})$; N_0 is a positive number, the choice of the value of which will be determined later.

Let $x \in [0, \frac{1}{2}]$, $k \in \Omega_\tau$. We write the mean value formula (see [3], [8], [9]) for the points $x, x+t, x+2t$ at $t \leq R_0$:

$$u_k(x) = 2u_k(x+t) \cos \mu_k t - u_k(x+2t) + \mu_k^{-1} \int_x^{x+2t} \left\{ q_1(\xi) u'_k(\xi) + \right. \\ \left. + q_2(\xi) u_k(\xi) - \theta_k u_{k-1}(\xi) \right\} \sin \mu_k (|x+t-\xi| - t) d\xi. \quad (2.2)$$

Since, according to the condition of the theorem, the rank of eigenfunctions is uniformly bounded, then it suffices to consider only the eigen-functions $u_k(x)$, i.e. the case $\theta_k = 0$. Then adding and subtracting on the right hand of formula (2.2) the expression $2u_k(x+t) \cos \tau t$ and by using the identity

$$\cos a - \cos \beta = 2 \sin \frac{a+\beta}{2} \sin \frac{\beta-a}{2},$$

then applying the operation $R_0^{-1} \int_0^{R_0} dt$, in the result we can obtain

$$u_k(x) = R_0^{-1} (u_k, v) + 4R_0^{-1} \int_0^{R_0} u_k(x+t) \sin \frac{\mu_k + \tau}{2} t \sin \frac{\tau - \mu_k}{2} t dt + \\ + (R_0 \mu_k)^{-1} \int_0^{R_0} \int_x^{x+2t} \left\{ q_1(\xi) u'_k(\xi) + q_2(\xi) u_k(\xi) \right\} \sin \mu_k (|x+t-\xi| - t) d\xi dt, \quad (2.3)$$

where $v(t) = 2 \cos \tau (x-t) - \frac{1}{2}$ for $t \in [x, x+R_0]$, $v(t) = -1/2$ for $t \in [x+R_0, x+2R_0]$ and $v(t) = 0$ for $t \notin [x, x+2R]$.

In the second addend on the right hand side of the formula (2.3) applying to the expression $u_k(x+t)$ following shift formula (see [3], [9], [10])

$$u_k(x \pm t) = u_k(x) \cos \mu_k t \pm \mu_k^{-1} \sin \mu_k t u'_k(x) \pm \\ \pm \mu_k^{-1} \int_x^{x \pm t} \left\{ q_1(\xi) u'_k(\xi) + q_2(\xi) u_k(\xi) - \theta_k u_{k-1}(\xi) \right\} \sin \mu_k (|x-\xi| - t) d\xi \quad (2.4)$$

for $\theta_k = 0$, we get

$$u_k(x) = R_0^{-1} (u_k, v) + 4R_0^{-1} u_k(x) \int_0^{R_0} \cos \mu_k t \sin \frac{\mu_k + \tau}{2} t \sin \frac{\tau - \mu_k}{2} t dt + \\ + 4(\mu_k R_0)^{-1} u'_k(x) \int_0^{R_0} \sin \mu_k t \sin \frac{\mu_k + \tau}{2} t \sin \frac{\tau - \mu_k}{2} t dt + \\ + 4(\mu_k R_0)^{-1} \int_0^{R_0} \sin \frac{\mu_k + \tau}{2} t \sin \frac{\tau - \mu_k}{2} t dt \int_x^{x+t} \left\{ q_1(\xi) u'_k(\xi) + \right. \\ \left. + q_2(\xi) u_k(\xi) \right\} \sin \mu_k (|x-\xi| - t) d\xi dt +$$

$$\begin{aligned}
& + (\mu_k R_0)^{-1} \int_0^{R_0} \int_x^{x+2t} \{q_1(\xi) u'_k(\xi) + q_2(\xi) u_k(\xi)\} \sin \mu_k (|x+t-\xi|-t) d\xi dt = \\
& = R_0^{-1} (u_k, v) + \sum_{j=1}^4 T_j.
\end{aligned} \tag{2.5}$$

Let estimate the integrals T_j , $j = \overline{1, 4}$. Taking into account $k \in \Omega_\tau$ and using the inequality

$$|\sin z| \leq 2, \quad |\cos z| \leq 2, \quad |\sin z| \leq 2|z| \quad \text{for } \operatorname{Im} z \leq 1$$

we get

$$\begin{aligned}
|T_1| & \leq 8R_0 |\tau - \mu_k| |u_k(x)| \leq 8R_0 (1 + C_0) |u_k(x)| \leq 8n_0^{-1} |u_k(x)|, \\
|T_2| & \leq 8R_0 |\tau - \mu_k| |\mu_k^{-1}| |u'_k(x)| \leq 8R_0 (1 + C_0) |\mu_k^{-1}| |u'_k(x)| \leq 8n_0^{-1} |\mu_k|^{-1} |u'_k(x)|
\end{aligned}$$

We estimate T_3 for $k \in \Omega_\tau$. Using the estimations (see [8],[9])

$$\begin{aligned}
& \|u_k^{(s)}\|_\infty \leq C_1 (1 + |\mu_k|)^s (1 + |\operatorname{Im} \mu_k|)^{1/p} \times \\
& \times \left\{ \|u_k\|_p + \theta_k |\mu_k|^{-1} (1 + |\operatorname{Im} \mu_k|)^{-1} \|\mu_{k-1}\|_p \right\}, \quad 1 \leq p < \infty, s = 0, 1;
\end{aligned} \tag{2.6}$$

$$\|u'_k\|_p \leq C_2 (1 + |\mu_k|) \left\{ \|u_k\|_p + \theta_k |\mu_k|^{-1} (1 + |\operatorname{Im} \mu_k|)^{-1} \|\mu_{k-1}\|_p \right\}, \quad p \geq 1, \tag{2.7}$$

for $\theta_k = 0$ and applying the above elementary inequalities, we have

$$\begin{aligned}
|T_3| & \leq 32R_0 |\mu_k^{-1}| \int_0^{R_0} \int_x^{x+t} |q_1(\xi)| |u'_k(\xi)| d\xi dt + \\
& + 32R_0 \int_0^{R_0} \int_x^{x+t} |q_2(\xi)| d\xi dt \|u_k\|_\infty \leq 32\omega(R_0) |\mu_k|^{-1} \|u'_k\|_\infty + 16R_0\omega(R_0) \|u_k\|_\infty \leq \\
& \leq 32\omega(R_0) C_1 \left(1 + |\mu_k|^{-1}\right) (1 + |\operatorname{Im} \mu_k|)^{1/q} \|u_k\|_q + \\
& + 16R_0\omega(R_0) C_1 (1 + |\operatorname{Im} \mu_k|)^{1/q} \|u_k\|_q \leq \\
& \leq 64\omega(R_0) (1 + C_0)^{1/2} C_1 \|u_k\|_q + 16C_1 (1 + C_0)^{1/2} R_0\omega(R_0) \|u_k\|_q \leq \\
& \leq C_3\omega(R_0) \|u_k\|_q \leq C_4 N_0^{-1} \|u_k\|_q.
\end{aligned}$$

The same estimation is fulfilled also for the integral T_4 for $k \in \Omega_t$.

Consequently, from (2.5) we get

$$\begin{aligned}
& |u_k(x)| \|u_k\|_q^{-1} \leq R_0^{-1} \left| (u_k \|u_k\|_q^{-1}, v) \right| + \\
& + 8n_0^{-1} |u_k(x)| \|u_k\|_q^{-1} + 8n_0^{-1} |\mu_k|^{-1} |u'_k(x)| \|u_k\|_q^{-1} + C_4 N_0^{-1}
\end{aligned} \tag{2.8}$$

By virtue of symmetry (see formulas (2.2), (2.4)) this inequality is valid in the case $x \in [0, \frac{1}{2}]$, as well. This time the function $v(t)$ is determined by the formula: $v(t) = \frac{1}{2}$ for $t \in [x - 2R_0, x - R_0]$, $v(t) = 2\cos \tau (x - t) - \frac{1}{2}$ for $t \in [x - R_0, x]$, $v(t) = 0$ for $t \notin [x - 2R_0, x]$.

Hence, for $n_0 \geq 16$ it follows that

$$\begin{aligned} |u_k(x)| \|u_k\|_q^{-1} &\leq 2R_0^{-1} \left| \left(u_k \|u_k\|_q^{-1}, v \right) \right| + \\ &+ 16n_0^{-1} |\mu_k|^{-1} \left| u'_k(x) \right| \|u_k\|_q^{-1} + 2C_4 N_0^{-1}. \end{aligned} \quad (2.9)$$

We raised to the power q every part of inequality (2.9), applying inequality

$$\left(\sum_{i=1}^3 a_i \right)^q \leq 3^{q-1} \sum_{i=1}^3 a_i^q, \quad a_i \geq 0,$$

integrating given result with respect to x from 0 to 1, we get

$$\begin{aligned} 1 &\leq 2^q 3^{q-1} R_0^{-q} \left| \left(u_k \|u_k\|_q^{-1}, v \right) \right|^q + \\ &+ 3^{q-1} 16^q n_0^{-q} |\mu_k|^{-q} \left| u'_k \right|_q^q \|u_k\|_q^{-q} + 3^{q-1} (2C_4 N_0^{-1})^q \end{aligned}$$

applying inequality (2.7) for $p = q$, $\theta_k = 0$, we obtain

$$\begin{aligned} 1 &\leq 2^q 3^{q-1} R_0^{-q} \left| \left(u_k \|u_k\|_q^{-1}, v \right) \right|^q + 3^{q-1} 16^q n_0^{-q} C_2^q \left(1 + |\mu_k|^{-1} \right)^q + \\ &+ 3^{q-1} (2C_4 N_0^{-1})^q \end{aligned}$$

Summing above inequality over $k \in \Omega_\tau$, we get

$$\begin{aligned} \sum_{k \in \Omega_\tau} 1 &\leq 2^q 3^{q-1} R_0^{-q} \sum_{k \in \Omega_\tau} \left| \left(u_k \|u_k\|_q^{-1}, v \right) \right|^q + \\ &+ 3^{q-1} 16^q n_0^{-q} C_2^q \sum_{k \in \Omega_\tau} \left(1 + |\mu_k|^{-1} \right)^q + 3^{q-1} (2C_4 N_0^{-1})^q \sum_{k \in \Omega_\tau} 1. \end{aligned}$$

since, $\left\{ u_k(x) \|u_k\|_q^{-1} \right\}$ is Riesz system, and $1 + |\mu_k|^{-1} \leq 2$ for $k \in \Omega_\tau$, then

$$\sum_{k \in \Omega_\tau} 1 \leq 2^q 3^{q-1} R_0^{-q} \|v\|_p^q + 3^{q-1} \left(\left(\frac{32C_2}{n_0} \right)^q + (2C_4 N_0^{-1})^q \right) \sum_{k \in \Omega_\tau} 1.$$

Taking into account estimate $\|v\|_p^q \leq (6R_0)^{q/p}$, we have

$$\sum_{k \in \Omega_\tau} 1 \leq 6^{q/p} R_0^{-1} + \left\{ C_5 n_0^{-q} + C_6 N_0^{-q} \right\} \sum_{k \in \Omega_\tau} 1,$$

where $C_5 = 3^{q-1} (32C_2)^q$, $C_6 = 3^{q-1} (2C_4)^q$.

Choosing the numbers n_0 ($n_0 \geq 16$) and N_0 so that $C_5 n_0^{-q} + C_6 N_0^{-q} \leq \frac{1}{2}$, we have at the inequality

$$\sum_{k \in \Omega_\tau} 1 \leq \text{const.}$$

Consequently, for $|\mu_k| \geq 1$ necessity of condition (1.2) is established.

For $|\mu_k| < 1$ the validity of condition (1.2) is proved in the following way. We consider the equation $Lu_k - 2u_k + \lambda'_k u_k = 0$, where $\lambda'_k = \lambda_k + 2$, $|\lambda_k| < 1$. Then $\left| \text{Re} \lambda'_k \right| \geq 1$

and the system $\{u_k(x)\}$ does not change. Therefore inequality (1.2) is fulfilled in the case $|\lambda_k| < 1$ as well. The necessity of condition (1.2) is established.

Sufficiency. Let conditions (1.1) and (1.2) be fulfilled. Prove that $\left\{u_k(x) \|u_k\|_q^{-1}\right\}_{k=1}^{\infty}$ is Riesz system. Firstly we prove that $\left\{u_k(x) \|u_k\|_2^{-1}\right\}_{k=1}^{\infty}$ is Bessel system in $L_2(G)$. By formula (2.4), conditions (1.1) and (1.2) for the convergence of the series $\sum_{k=1}^{\infty} \left| \left(f, u_k \|u_k\|_2^{-1} \right) \right|^2$ for any $f \in L_2(G)$, it suffices to prove the validity of the following inequalities:

$$\sum_{|\mu_k| \geq 1} \left| \int_0^1 \overline{f(t)} \cos \mu_k t dt \right|^2 \|u_k\|_2^{-2} |u_k(0)|^2 \leq C \|f\|_2^2; \quad (2.10)$$

$$\sum_{|\mu_k| \geq 1} |\mu_k|^{-2} \left| \int_0^1 \overline{f(t)} \sin \mu_k t dt \right|^2 |u_k(0)|^2 \leq C \|f\|_2^2; \quad (2.11)$$

$$\sum_{|\mu_k| \geq 1} |\mu_k|^{-2} \left| \int_0^1 \overline{f(t)} \int_0^t q_1(\xi) u'_k(\xi) \sin \mu_k(\xi - t) d\xi dt \right|^2 \|u_k\|_2^{-2} \leq C \|f\|_2^2; \quad (2.12)$$

$$\sum_{|\mu_k| \geq 1} |\mu_k|^{-2} \left| \int_0^1 \overline{f(t)} \int_0^t q_2(\xi) u_k(\xi) \sin \mu_k(\xi - t) d\xi dt \right|^2 \|u_k\|_2^{-2} \leq C \|f\|_2^2; \quad (2.13)$$

$$\sum_{|\mu_k| \geq 1} \theta_k |\mu_k|^{-2} \left| \int_0^1 \overline{f(t)} \int_0^t u_{k-1}(\xi) \sin \mu_k(\xi - t) d\xi dt \right|^2 \|u_k\|_2^{-2} \leq C \|f\|_2^2, \quad (2.14)$$

where $f \in L_2(G)$.

Under conditions (1.1), (1.2), the validity of inequalities (2.10), (2.11), (2.13) follows from [3].

Prove the validity of the inequality (2.12). Denote

$$g(t, \xi) = \begin{cases} f(t + \xi), & 0 \leq t \leq 1 - \xi, \\ 0, & 1 - \xi < t \leq 1, \end{cases} \quad \text{where } \xi \in [0, 1].$$

Then

$$\begin{aligned} S_k(f) &= |\mu_k|^{-2} \left| \int_0^1 \overline{f(t)} \int_0^t q_1(\xi) u'_k(\xi) \sin \mu_k(\xi - t) d\xi dt \right|^2 \|u_k\|_2^{-2} = \\ &= \int_0^t q_1(\xi) u'_k(\xi) \mu_k^{-1} \|u_k\|_2^{-1} \int_0^1 \overline{g(t, \xi)} \sin \mu_k t dt d\xi \times \\ &\times \int_0^1 \overline{q_1(z) u'_k(z) \mu_k^{-1}} \|u_k\|_2^{-1} \int_0^1 g(r, z) \overline{\sin \mu_k r} dr dz = \\ &= \int_0^1 \int_0^1 q_1(\xi) \overline{q_1(z)} u'_k(\xi) \mu_k^{-1} \|u_k\|_2^{-1} \overline{u'_k(z) \mu_k^{-1}} \|u_k\|_2^{-1} \times \\ &\times \int_0^1 \overline{g(t, \xi)} \sin \mu_k t dt \int_0^1 g(r, z) \overline{\sin \mu_k r} dr d\xi dz \end{aligned}$$

Applying here the estimation (see [5])

$$\left\| u_k' \right\|_{\infty} \leq C_7 (1 + |\mu_k|) \|u_k\|_2$$

we get that at $|\mu_k| \geq 1$ for $S_k(f)$ the following inequality is fulfilled:

$$S_k(f) \leq C_8 \int_0^1 \int_0^1 |q_1(\xi)| |q_1(z)| \left| \int_0^1 \overline{g(t, \xi)} \sin \mu_k t dt \right| \left| \int_0^1 \overline{g(r, z)} \sin \mu_k r dr \right| d\xi dz.$$

For an arbitrary finite subset J' of the set of indices $J = \{k : |\mu_k| \geq 1\}$, we obtain

$$\begin{aligned} \sum_{k \in J'} S_k(f) &\leq C_8 \int_0^1 \int_0^1 |q_1(\xi)| |q_1(z)| \left(\sum_{k \in J'} \left| \int_0^1 \overline{g(t, \xi)} \sin \mu_k t dt \right|^2 \right)^{1/2} \times \\ &\quad \times \left(\sum_{k \in J'} \left| \int_0^1 \overline{g(r, z)} \sin \mu_k r dr \right|^2 \right)^{1/2} d\xi dz \leq \\ &\leq C_9 \int_0^1 \int_0^1 |q_1(\xi)| |q_1(z)| \|g(\cdot, \xi)\|_2 \|g(\cdot, z)\| d\xi dz. \end{aligned}$$

Taking into account that for any $\xi \in [0, 1]$ we have the inequality $\|g(\cdot, \xi)\|_2 \leq \|f\|_2$, we obtain

$$\sum_{k \in J'} S_k(f) \leq C_{10} \|q_1\|_1^2 \|f\|_2^2.$$

Hence, from arbitrariness of $J' \subset J$, it follows the inequality

Therefore, for an arbitrary function $f(x) \in L_2(G)$ the following Bessel inequality is satisfied:

$$\left(\sum_{k=1}^{\infty} \left| \left(f, u_k \|u_k\|_2^{-1} \right) \right|^2 \right)^{1/2} \leq M \|f\|_2. \quad (2.15)$$

On the other hand, for any $f \in L_1(G)$

$$\sup_k \left(f, u_k \|u_k\|_2^{-1} \right) \leq \sup_k \left(\|u_k\|_{\infty} \|u_k\|_2^{-1} \right) \|f\|_1$$

Hence, by virtue of the inequality (see [5])

$$\|u_k\|_r \leq c_{11} \|u_k\|_{\gamma}, \quad \gamma \geq 1, \quad r \geq 1 \quad (2.16)$$

we get that for any function $f \in L_1(G)$ the following inequality is valid:

$$\sup_k \left(f, u_k \|u_k\|_2^{-1} \right)^{1/2} \leq M_2 \|f\|_1. \quad (2.17)$$

By virtue of the Riesz-Torin interpolation theorem (see [17], p.144) from inequalities (2.15) and (2.17) it follows that the system $\left\{ u_k(x) \|u_k\|_2^{-1} \right\}_{k=1}^{\infty}$ satisfies the Riesz inequality, i.e.

$$\left(\sum_{k=1}^{\infty} \left| \left(f, u_k(x) \|u_k\|_2^{-1} \right) \right|^q \right)^{1/q} \leq M(p) \|f\|_q$$

for $1 < p \leq 2$, $q = p/(p-1)$.

Since

$$u_k(x) \|u_k\|_2^{-1} = u_k(x) \|u_k\|_q^{-1} \|u_k\|_q \|u_k\|_2^{-1}$$

and by virtue of inequality (2.16), the estimation

$$1 \leq \|u_k\|_q \|u_k\|_2^{-1} \leq C_{12}$$

is fulfilled, then the system $\left\{ u_k(x) \|u_k\|_q^{-1} \right\}_{k=1}^{\infty}$ satisfies the Riesz inequality as well, i.e. the inequality

$$\left(\sum_{k=1}^{\infty} \left| \left(f, u_k(x) \|u_k\|_q^{-1} \right) \right|^q \right)^{1/q} \leq M_2(p) \|f\|_q, \quad q = p/(p-1)$$

is fulfilled for any $f \in L_p(G)$, $1 < p \leq 2$.

Theorem is completely proved.

References

1. Abdullayeva, A.M.: *On local componentwise equiconvergence for one-dimensional Dirac operator*, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci. Mathematics, **39** (1), 3-14 (2019).
2. Budaev, V.D.: *Criteria for the Bessel property and the Riesz basis property of systems of root functions of differential operators. I*, Differ. Uravn. **27**:12 (1991), 2033-2044 ; Differ. Equ. **27** (12), 1421-1432 (1991).
3. Ilin, V.A.: *Unconditional basis property on a closed interval of systems of eigen and associated functions of a second-order differential operator*, Dokl. Akad. Nauk SSSR, **273** (5), 1048-1053 (1983).
4. Kerimov, N.B.: *Unconditional basis property of a system of eigen- and associated functions of a fourth-order differential operator*, Dokl. Akad. Nauk SSSR, **286** (4), 803-808 (1986).
5. Kerimov, N.B.: *Some properties of eigen and associated functions of ordinary differential operators*, Dokl. Akad. Nauk SSSR, **291** (5), 1054-1056 (1986).
6. Kritskov, L.V.: *Bessel property of the system of root functions of a second-order singular operator on an interval*, Diff. Equat. **54**, 1032-1048 (2018).
7. Kurbanov, V.M.: *On the Hausdorff-Young inequality for systems of root vector functions of an n -th order differential operator*, Differ. Uravn. **33**:3 358-367 (1997); Differ. Equ. **33**:3 , 355-365 (1997).
8. Kurbanov, V.M.: *On the eigenvalue distribution of and a Bessel criterion property criterion for root functions of a differential operator. I*, Diff. Equat. **41**, 489-505 (2005).
9. Kurbanov, V.M.: *On the eigenvalue distribution and a Bessel property criterion for root functions of a differential operator: II*, Diff. Equat. **41**, 649-659 (2005).
10. Kurbanov, V.M.: *A theorem on equivalent bases for differential operators*, Dokl. Math. **73**, 11-14 (2006).
11. Kurbanov, V.M.: *Analog of the Riesz theorem and the basis property in L_p of a system of root functions of a differential operator: I*, Diff. Equat. **49**, 7-19 (2013).
12. Kurbanov, V.M.: *On an analog of the Riesz theorem and the basis property of the system of root functions of a differential operator in L_p . II*, Diff Equat. **49**, 437-449 (2013).
13. Kurbanov, V.M., Buksayeva, L.Z., Ismailova, A.I.: *Criteria for componentwise uniform equiconvergence with trigonometric series of spectral expansions responding to discontinuous Dirac operator*, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci. Mathematics, **41** (4), 112-124 (2021).

-
14. Kurbanov, V.M., Gojaeva, Kh. R., Shahbazov, R. I.: *On absolute and uniform convergence of a biorthogonal series in root functions of an odd order differential operator*, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci. Mathematics, **42** (4), 102-111 (2022).
 15. Kurbanov, V.M., Ibadov, E. J.: *On Riesz property and equivalent basis property of the system of root vector functions of Dirac-type operator*, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci. Mathematics, **44** (1), 51-62 (2024).
 16. Lomov, I.S.: *Bessel inequality, Riesz theorem, and unconditional basis property for root vectors of ordinary differential operators*, Vestnik Moskov. Univ. Ser. 1. Mat. Mekh. no. 5, 33-43, (1992).
 17. Zigmund, A. Trigonometric series. Cambridge, (1959), Translated under the title *Trigonometricheskie ryady*, Moscow, **2** (1965).