

## On stability of a basis consisting of an perturbed exponential system in the weighted grand Lebesgue space

Migdad I. Ismailov · Tural X. Hagverdi\*

Received: 02.05.2025 / Revised: 18.01.2026 / Accepted: 12.02.2026

**Abstract.** *In this paper, we investigate the stability of a perturbed exponential system  $\{e^{i\alpha_n x}\}_{n \in \mathbb{Z}}$  in a separable subspace of the weighted grand Lebesgue space  $N_{p,\omega}(-\pi, \pi)$ . Analogs of classical Levinson-type theorems are obtained for the completeness of the system in space  $L_{p,\omega}(-\pi, \pi)$ ,  $1 < p < +\infty$  and using this result, the stability of the perturbed exponential system is established.*

**Keywords.** exponential system, weighted grand Lebesgue space,  $q$ -Hilbert system, separated system, Levinson theorem.

**Mathematics Subject Classification (2010):** 33B10, 46E30, 54D70

### 1 Introduction

The study of exponential systems plays a central role in harmonic analysis and the theory of function spaces. Among them, perturbed exponential systems, which generalize classical exponential systems, have attracted considerable interest due to their connection with eigenfunctions of differential operators subject to integral boundary conditions. The investigation of exponential systems, commonly known as the theory of nonharmonic Fourier series (see [1–5]), traces its roots to the classical works in this direction include those by Paley and Wiener [6], N. Levinson [1], who explored the basis properties of such systems in various functional frameworks. One of the well-known early results in this area states that the trigonometric system  $\{e^{inx}\}_{n=-\infty}^{+\infty}$  possesses a stable basis property in  $L_2(-\pi, \pi)$ . More precisely, the system  $\{e^{i\lambda_n x}\}_{n=-\infty}^{+\infty}$  forms a Riesz basis in  $L_2(-\pi, \pi)$  provided that  $|\lambda_n - n| \leq L < 1/4$ . The optimality of the constant  $\frac{1}{4}$  was established by M. I. Kadec [7], and later by R. M. Redheffer and R. M. Young [8].

In recent years, the investigation of such systems in generalized function spaces, such as grand Lebesgue spaces (see [9–16]) and weighted grand Lebesgue spaces, has gained

\* Corresponding author

This work was supported by the Azerbaijan Science Foundation - Grant No: AEF-MGC-2025-1(54)-20/03/1-M-03

M.I. Ismailov  
Azerbaijan State Oil and Industry University, Baku, Azerbaijan  
E-mail: migdad-ismailov@rambler.ru

T.X. Hagverdi  
Institute of Mathematics and Mechanics, The Ministry of Science and Education of the Republic of Azerbaijan, Baku, Azerbaijan  
E-mail: turalhagverdi@gmail.com

significant momentum. These spaces provide a flexible framework for analyzing systems with variable integrability properties and are particularly effective in addressing questions of stability and completeness. In this direction, special attention has been paid to the stability of bases generated by perturbed exponential systems, both in classical  $L_p$  (see [17]), Morrey-Lebesgue spaces (see [18]) and in grand Lebesgue spaces where the non-uniform integrability structure plays a crucial role in the preservation of basis properties (see also [19–21]).

Now, let us consider a certain sequence of real numbers  $\{\alpha_n\}_{-\infty}^{\infty}$  of real numbers. We examine the following exponential system

$$\{e^{i\alpha_n x}\}_{n \in \mathbb{Z}}. \quad (1.1)$$

In this paper, we focus on the stability of a basis formed by a perturbed exponential system within a separable subspace  $N_{p,\omega}(-\pi, \pi)$  of the weighted grand Lebesgue space  $L_{p,\omega}(-\pi, \pi)$ ,  $1 < p < +\infty$ . We establish analogues of classical Levinson-type theorems, providing criteria for the completeness and minimality of the system in this setting. The stability of system (1.1) in space  $N_{p,\omega}(-\pi, \pi)$ ,  $1 < p < +\infty$ , is studied.

## 2 Preliminaries

Let  $\omega(x)$  be a weight function defined on  $[-\pi, \pi]$ . We denote by  $L_{p,\omega}(-\pi, \pi)$ ,  $1 < p < +\infty$ , the weighted grand Lebesgue space consisting of measurable functions defined on  $[-\pi, \pi]$  with finite norm

$$\|f\|_{p,\omega} = \sup_{0 < \varepsilon < p-1} \left( \frac{\varepsilon}{2\pi} \int_{-\pi}^{\pi} |f(t)\omega(t)|^{p-\varepsilon} dt \right)^{\frac{1}{p-\varepsilon}} < +\infty.$$

We denote by  $A_p(-\pi, \pi)$ ,  $1 < p < +\infty$ , the Muckenhoupt weight class consisting of periodic functions  $\omega(x)$  with period  $2\pi$ , satisfying the following condition

$$\sup_{I \subset [-\pi, \pi]} \left( \frac{1}{|I|} \int_I \omega^p(t) dt \right) \left( \frac{1}{|I|} \int_I \omega^{-\frac{p}{p-1}}(t) dt \right)^{p-1} < +\infty,$$

where the supremum is taken over all finite intervals  $I$  in  $[-\pi, \pi]$ .

Since the space  $L_{p,\omega}(-\pi, \pi)$ ,  $1 < p < +\infty$ , is non-separable, we consider its separable subspace  $N_{p,\omega}(-\pi, \pi)$  of functions  $f \in L_{p,\omega}(-\pi, \pi)$ :

$$\|f(\cdot + \delta) - f(\cdot)\|_{L_{p,\omega}(-\pi, \pi)} \rightarrow 0, \quad \delta \rightarrow 0.$$

**Theorem 2.1 ([22])** *Let there is  $\varepsilon \in (0, p-1)$  such that  $\omega^{-1} \in L_{\frac{1}{p-\varepsilon-1}}$ . Then set of infinitely differentiable functions  $C^\infty[-\pi, \pi]$  is dense in  $N_{p,\omega}(-\pi, \pi)$ .*

Let  $X$  be a b.f.s. over  $(M; \mu)$  with norm  $\|\cdot\|_X$ . We will also use certain concepts and results from the theory of Banach function spaces (see [23]). The necessary facts from this theory are stated below.

**Definition 2.1 ([23])** *A function  $f$  in a b.f.s. (Banach function space)  $X$  is said to have continuous norm in  $X$  if  $\|f\chi_{E_n}\|_X \rightarrow 0$ , for every sequence  $\{E_n\}_{n=1}^{\infty}$  satisfying  $E_n \rightarrow \emptyset$ ,  $\mu - \text{a.e.}$  The set of all functions in  $X$  of absolutely continuous norm is denoted by  $X_a$ .*

**Theorem 2.2 ([23])** *The dual  $X^*$  of a b.f.s.  $X$  is canonically isometrically isomorphic to the associate space  $X'$  if and only if  $X$  have absolutely continuous norm, where*

$$X' = \left\{ g \in L_1 : \|g\|_{X'} = \sup_{\|f\|_X \leq 1} \int_M |f(t)g(t)| dt < +\infty \right\}.$$

We also need the following theorems.

**Theorem 2.3 (Hausdorff-Young)** *Let  $1 \leq p \leq 2$ ,  $\frac{1}{p} + \frac{1}{q} = 1$ ,  $f \in L^p(-\pi, \pi)$ , and define its Fourier coefficients by  $\hat{f}(n) = c_n := \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x)e^{-inx} dx$ ,  $n \in \mathbb{Z}$ . Then Hausdorff-Young inequality states that*

$$\| \{c_n\}_{n \in \mathbb{Z}} \|_{\ell_q} \leq C_p \|f\|_{L_p},$$

where

$$\| \{c_n\} \|_{\ell_q} = \left( \sum_{n \in \mathbb{Z}} |c_n|^q \right)^{1/q}$$

and  $C_p$  is a constant depending only on  $p$ .

We will also use the following analogue of the Paley-Wiener theorem in Banach spaces.

**Theorem 2.4** *Let  $X$  be a Banach space and let  $\{x_n\}_{n=1}^{\infty} \in X$ , be a (Schauder) basis of  $X$ . Let  $\{y_n\}_{n=1}^{\infty} \in X$ , be another sequence such that there exists a constant  $\theta \in [0, 1)$  satisfying*

$$\left\| \sum_{n=1}^m a_n (y_n - x_n) \right\| \leq \theta \left\| \sum_{n=1}^m a_n x_n \right\|$$

for all finite scalar sequences  $(a_1, \dots, a_m)$ . Then  $\{y_n\}$  is also a Schauder basis of  $X$ . Moreover,  $\{x_n\}$  and  $\{y_n\}$  are equivalent bases.

### 3 A Necessary Condition for the Sequence $\{\alpha_n\}$ to be Separated

**Definition 3.1 ([24])** *A system  $\{f_n\}_{n \in \mathbb{N}} \subset L_{p,\omega}(-\pi, \pi)$ ,  $1 < p < +\infty$ , is called a  $q$ -Hilbert system, if there exists  $C > 0$ , such that for any finite system  $\{c_n\}_{n \in \mathbb{Z}}$  of complex numbers the following inequality holds*

$$\left( \sum_n |c_n|^q \right)^{\frac{1}{q}} \leq C \left\| \sum_n c_n f_n \right\|_{L_{p,\omega}},$$

where  $\frac{1}{q} + \frac{1}{p} = 1$ .

**Definition 3.2 ([25])** *A sequence  $\{\lambda_n\}_{n \in \mathbb{Z}} \subset \mathbb{R}$  is called separated if*

$$\inf_{n \neq m} |\lambda_n - \lambda_m| > 0,$$

holds.

**Lemma 3.1** *Let  $\{\alpha_n\}_{n \in \mathbb{Z}}$  be a sequence of real numbers. If for  $\omega \neq 0$  and  $\omega \in L_p(-\pi, \pi)$  the system  $\{e^{i\alpha_n x}\}_{n \in \mathbb{Z}}$  is a  $q$ -Hilbert system in the space  $L_{p,\omega}(-\pi, \pi)$ ,  $1 < p < +\infty$ , then the sequence  $\{\alpha_n\}_{n \in \mathbb{Z}}$  is separated.*

**Proof.** By the definition of q-Hilbert system, we have:

$$\left( \sum_{k=-\infty}^{\infty} |c_k|^q \right)^{\frac{1}{q}} \leq C \sup_{0 < \varepsilon < p-1} \left( \frac{\varepsilon}{2\pi} \int_{-\pi}^{\pi} \left| \sum_{k=-\infty}^{\infty} c_k f_k \omega \right|^{p-\varepsilon} dt \right)^{\frac{1}{p-\varepsilon}}.$$

If we choose the coefficients as  $c_n = 1$ ,  $c_m = -1$ ,  $c_k = 0$ ,  $k \neq n \neq m$  and  $f_k = e^{i\alpha_k t}$  then we obtain

$$\begin{aligned} 2^{\frac{1}{q}} &\leq C \sup_{0 < \varepsilon < p-1} \left( \frac{\varepsilon}{2\pi} \int_{-\pi}^{\pi} |(e^{i\alpha_n t} - e^{i\alpha_m t}) \omega|^{p-\varepsilon} dt \right)^{\frac{1}{p-\varepsilon}} \Rightarrow \\ &2^{\frac{1}{q}} \leq C \|e^{i\alpha_n t} - e^{i\alpha_m t}\|_{p, \omega}. \end{aligned} \quad (3.1)$$

On the other hand

$$|e^{i\alpha_n t} - e^{i\alpha_m t}| = 2 \left| \sin \left( \frac{\alpha_n - \alpha_m}{2} t \right) \right| \leq |\alpha_n - \alpha_m| |t| \leq \pi |\alpha_n - \alpha_m|.$$

It follows from this that

$$\|e^{i\alpha_n t} - e^{i\alpha_m t}\|_{L_p, \omega(-\pi, \pi)} \leq \pi |\alpha_n - \alpha_m| \|\omega\|_{L_p(-\pi, \pi)}.$$

Then, from relation (4.1), we obtain

$$\begin{aligned} 2^{\frac{1}{q}} &\leq \|e^{i\alpha_n t} - e^{i\alpha_m t}\|_{L_p, \omega(-\pi, \pi)} \leq \pi |\alpha_n - \alpha_m| \|\omega\|_{L_p(-\pi, \pi)} \Rightarrow \\ |\alpha_n - \alpha_m| &\geq \frac{2^{\frac{1}{q}}}{\pi \|\omega\|_{L_p(-\pi, \pi)}} \Rightarrow \inf_{n \neq m} |\alpha_n - \alpha_m| > 0. \end{aligned}$$

The theorem is thus proved.

#### 4 Analogues of Levinson-Type Theorems in the Space $N_{p, \omega}(-\pi; \pi)$

We establish the following analogues of Levinson's theorems [26].

**Theorem 4.1** *Let  $\{\alpha_k\}_{k=1}^{\infty}$  be a sequence of complex numbers and  $\omega \in A_p$ ,  $1 < p < \infty$ . A necessary and sufficient condition for the exponential system  $\{e^{i\alpha_k t}\}_{k \in \mathbb{N}}$  to be incomplete in the space  $N_{p, \omega}(-\pi, \pi)$  is the existence of a non-zero entire function  $F(z)$  of exponential type such that*

$$F(\alpha_n) = 0 \quad \text{for all } n.$$

Here

$$F(z) = \int_{-\pi}^{\pi} f(x) e^{izx} dx, \quad (4.1)$$

where  $f(x) \in N'_{p, \omega}(-\pi, \pi)$  and  $f \neq 0$ .

**Proof.** Suppose that the system  $\{e^{i\alpha_k t}\}_{k \in \mathbb{N}}$  is not complete in the space. Then there exists a non-zero functional  $L \in N'_{p,\omega}(-\pi, \pi)$  such that

$$L(e^{i\alpha_k x}) = 0 \text{ for all } k.$$

Let us show that the spaces  $N'_{p,\omega}(-\pi, \pi)$  and  $N^*_{p,\omega}(-\pi, \pi)$  are isometrically isomorphic. According to Theorem 2.2 it is sufficient to prove that the space  $N_{p,\omega}(-\pi, \pi)$ ,  $1 < p < +\infty$ , admits an absolutely continuous norm. By Theorem 2.1, we have

$$\overline{C^\infty([-\pi, \pi])} = N_{p,\omega}(-\pi, \pi).$$

Let  $f \in L_{p,\omega}$  and let  $\varepsilon > 0$  be arbitrary. Then there exists  $f_0 \in C[-\pi, \pi]$  such that

$$\|f - f_0\|_{L_{p,\omega}} < \varepsilon.$$

Suppose that  $\{E_n\}_{n \in \mathbb{N}} \subset (-\pi, \pi)$  is a sequence of Lebesgue measurable sets such that  $E_n \rightarrow \emptyset$ ,  $\mu - a.e.$ , i.e.  $\chi_{E_n} \rightarrow 0$ ,  $\mu - a.e.$  We need to show

$$\|f\chi_{E_n}\|_{L_{p,\omega}} \rightarrow 0.$$

We have

$$\begin{aligned} \|f\chi_{E_n}\|_{L_{p,\omega}} &= \|(f - f_0)\chi_{E_n} + f_0\chi_{E_n}\|_{L_{p,\omega}} \\ &\leq \|(f - f_0)\chi_{E_n}\|_{L_{p,\omega}} + \|f_0\chi_{E_n}\|_{L_{p,\omega}} \\ &< \varepsilon + \|f_0\chi_{E_n}\|_{L_{p,\omega}}. \end{aligned} \quad (4.2)$$

Let  $M = \|f_0\|_{L^\infty(-\pi, \pi)}$ . Then we obtain

$$\|f_0\chi_{E_n}\|_{L_{p,\omega}} \leq M\|\chi_{E_n}\|_{L_{p,\omega}} \leq M_1\|\chi_{E_n}\|_{L_{p,\omega}} < \infty.$$

Obviously,  $\|\chi_{E_n}\|_{L_{p,\omega}} \rightarrow 0$ .

Consequently, from (4.2) it follows that

$$\|f\chi_{E_n}\|_{L_{p,\omega}} \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Thus, we conclude that the space  $N_{p,\omega}(-\pi, \pi)$  possesses an absolutely continuous norm. From Theorem 2.2, it follows that

$$N'_{p,\omega}(-\pi, \pi) \cong N^*_{p,\omega}(-\pi, \pi),$$

which means that there exists a function  $g \in N'_{p,\omega}(-\pi, \pi)$  such that for every  $f \in N_{p,\omega}(-\pi, \pi)$ , we have

$$L(f) = \int_{-\pi}^{\pi} f(x)g(x)dx. \quad (4.3)$$

Substituting  $f(x) = e^{i\alpha_n x}$  into (4.3), we obtain:

$$L(e^{i\alpha_n x}) = \int_{-\pi}^{\pi} e^{i\alpha_n x} g(x)dx = 0.$$

It thus becomes clear that  $F(z) = \int_{-\pi}^{\pi} f(x)e^{izx}dx$  is an entire function and hence

$$F(\alpha_n) = 0 \text{ for all } n.$$

Conversely, suppose that for the exponential system defined by (4.1), the entire function  $F(z)$  satisfies  $F(\alpha_n) = 0$  for all  $n$ . Then one may consider the functional  $L(f)$  defined by (4.3). By Hlder's inequality, the functional  $L$  is a linear bounded functional on  $N_{p,\omega}(-\pi, \pi)$ . On the other hand, since  $L \neq 0$  and

$$L(e^{i\alpha_n x}) = 0 \quad \text{for all } n,$$

the system  $\{e^{i\alpha_n x}\}_{n \in Z}$  is not complete.

The theorem is thus proved.

**Theorem 4.2** *Suppose that the function  $F(z)$  is represented in the form (4.1) and  $g \in N'_{p,\omega}(-\pi, \pi)$ ,  $1 < p < +\infty$ ,  $F(\alpha_0) = 0$  and  $\beta$  any complex number. Then*

$$F_1(\alpha) = \frac{\alpha - \beta}{\alpha - \alpha_0} F(\alpha),$$

is also represented in the form (4.1).

**Proof.** As in the classical Levinson theorem, let us define the new  $h(x)$  function in the following way

$$h(x) = g(x) + i(\beta - \alpha_0) e^{-i\alpha_0 x} \int_{-\pi}^x e^{i\alpha_0 y} g(y) dy. \quad (4.4)$$

If we multiply both sides of equation (4.4) by  $e^{i\alpha x}$  and integrate, we obtain

$$\begin{aligned} \int_{-\pi}^{\pi} e^{i\alpha x} h(x) dx &= \int_{-\pi}^{\pi} e^{i\alpha x} g(x) dx + i(\beta - \alpha_0) \int_{-\pi}^{\pi} e^{i(\alpha - \alpha_0)x} \left( \int_{-\pi}^x e^{i\alpha_0 y} g(y) dy \right) dx = \\ &= F(\alpha) + i(\beta - \alpha_0) \int_{-\pi}^{\pi} \int_y^{\pi} e^{i(\alpha - \alpha_0)x} e^{i\alpha_0 y} g(y) dx dy. \end{aligned}$$

By Fubini's theorem, changing the order of integration yields

$$\begin{aligned} \int_{-\pi}^{\pi} e^{i\alpha x} h(x) dx &= F(\alpha) + \frac{\beta - \alpha_0}{\alpha - \alpha_0} \int_{-\pi}^{\pi} \left( e^{i(\alpha - \alpha_0)\pi} - e^{i(\alpha - \alpha_0)y} \right) e^{i\alpha_0 y} g(y) dy \\ &= F(\alpha) + \frac{\beta - \alpha_0}{\alpha - \alpha_0} \int_{-\pi}^{\pi} e^{i\alpha y} g(y) dy \\ &= F(\alpha) - \frac{\beta - \alpha_0}{\alpha - \alpha_0} F(\alpha) = \frac{\alpha - \beta}{\alpha - \alpha_0} F(\alpha). \end{aligned}$$

Thus

$$F_1(\alpha) = \int_{-\pi}^{\pi} e^{i\alpha x} h(x) dx.$$

The theorem is proved.

**Corollary 4.1** *Suppose  $\omega \in A_p$  and the system  $\{e^{i\alpha_k x}\}_{k \in Z}$  is complete in the space  $N_{p,\omega}(-\pi, \pi)$ . Then, if we remove any  $n$  functions from this system and replace them with any other  $n$  exponential functions  $\{e^{i\beta_k x}\}_{k=1}^n$  (here  $\beta_k \neq \alpha_k, \forall k = \overline{1, n}$ ) then the newly obtained system will be complete in space  $N_{p,\omega}(-\pi, \pi)$ .*

### 5 Stability of the System $\{e^{i\alpha_n x}\}_{n \in \mathbb{Z}}$ in the Space $N_{p,\omega}(-\pi, \pi)$

Let us consider the following theorem concerning the stability of an exponential system with respect to the sequence  $\{\alpha_n\}_{n \in \mathbb{Z}}$  in the space  $N_{p,\omega}(-\pi, \pi)$ .

**Theorem 5.1** *Let  $1 < p < +\infty$ ,  $\omega \in A_p$ ,  $r \in (1, \min(p, q))$ ,  $r < p_1 < p$ ,  $\omega^{-1} \in L_{\frac{p_1 r}{p_1 - r}}$ ,  $\alpha_n, \beta_n \in \mathbb{R}$ ,  $n \in \mathbb{Z}$ , such that,  $\alpha_i \neq \alpha_j$ ,  $\beta_i \neq \beta_j$ , for  $i \neq j$ , and*

$$\sum_{n=-\infty}^{+\infty} |\alpha_n - \beta_n|^r < +\infty. \quad (5.1)$$

*If the system  $\{e^{i\alpha_n x}\}_{n \in \mathbb{Z}}$  forms a basis isomorphic to the system  $\{e^{inx}\}_{n \in \mathbb{Z}}$  in space  $N_{p,\omega}(-\pi, \pi)$ , then  $\{e^{i\beta_n x}\}_{n \in \mathbb{Z}}$  forms a basis isomorphic to the system  $\{e^{inx}\}_{n \in \mathbb{Z}}$  in  $N_{p,\omega}(-\pi, \pi)$ .*

**Proof.** Let's denote

$$e_n(x) = e^{inx}, \varphi_n(x) = e^{i\alpha_n x}$$

and

$$\psi_n(x) = e^{i\beta_n x}, n \in \mathbb{Z}, x \in [-\pi, \pi].$$

According to the Lemma 3.1,

$$\|\varphi_n - \psi_n\|_{p,\omega} \leq M |\alpha_n - \beta_n|, n \in \mathbb{Z}, \quad (5.2)$$

here  $M = \pi \|\omega\|_p$ . Let fix number  $m \in \mathbb{Z}_+$  and consider the system of functions as following

$$f_n = \begin{cases} \varphi_n, & |n| < m \\ \psi_n, & |n| \geq m \end{cases}, n \in \mathbb{Z}.$$

Applying inequality (5.2) together with Hölder's inequality, we obtain the following for any finite complex sequence  $\{c_n\}$

$$\begin{aligned} \left\| \sum_n c_n (f_n - \varphi_n) \right\|_{L_{p,\omega}} &\leq \sum_n |c_n| \|f_n - \varphi_n\|_{L_{p,\omega}} \\ &\leq M \sum_n |c_n| |\alpha_n - \beta_n| \\ &\leq M \left( \sum_n |c_n|^{r'} \right)^{\frac{1}{r'}} \left( \sum_n (\alpha_n - \beta_n)^r \right)^{\frac{1}{r}}. \end{aligned} \quad (5.3)$$

Since  $1 < r < 2$  and  $\omega \in A_p$  the system  $\{e^{inx}\}_{n \in \mathbb{Z}}$  becomes basis in  $N_{p,\omega}(-\pi, \pi)$  (see result [22]) so we can apply here by the Theorem 2.4 about Hausdroff-Young inequality

$$\left( \sum_n |c_n|^{r'} \right)^{\frac{1}{r'}} \leq \left\| \sum_n c_n e_n \right\|_r. \quad (5.4)$$

Since  $\omega^{-1} \in L_{\frac{p_1 r}{p_1 - r}}$  for  $\forall f \in L_{p,\omega}$  by using Hölder's inequality we obtain

$$\|f\|_r = \left( \int_{-\pi}^{\pi} |f\omega|^r \cdot \omega^{-r} dt \right)^{\frac{1}{r}} \leq$$

$$\leq \left( \int_{-\pi}^{\pi} |f\omega|^{p_1} dt \right)^{\frac{1}{p_1}} \left( \int_{-\pi}^{\pi} \omega^{-\frac{p_1 r}{p_1 - r}} dt \right)^{\frac{p_1 - r}{p_1 r}} \leq K \|f\|_{p, \omega}. \quad (5.5)$$

Taking inequality (5.5) into account in inequality (5.4), we obtain

$$\left( \sum_n |c_n|^{r'} \right)^{\frac{1}{r'}} \leq K \left\| \sum_n c_n e_n \right\|_{p, \omega}. \quad (5.6)$$

Since the systems  $\{\varphi_n\}$  and  $\{\psi_n\}$  are isomorphic bases, there exists a constant  $L > 0$  such that

$$\left\| \sum_n c_n e_n \right\|_{p, \omega} \leq L \left\| \sum_n c_n \varphi_n \right\|_{p, \omega} \quad (5.7)$$

Thus, taking inequality (5.7) into account in inequality (5.6), and using inequality (5.5), we have

$$\left\| \sum_n c_n (f_n - \varphi_n) \right\|_{p, \omega} \leq M_1 \left( \sum_{|n| \geq m} (\alpha_n - \beta_n)^r \right)^{\frac{1}{r}} \left\| \sum_n c_n \varphi_n \right\|_{p, \omega}.$$

Since the series  $\sum_{n=-\infty}^{+\infty} |\alpha_n - \beta_n|^r$  converges, for a sufficiently large  $m \in \mathbb{Z}_+$  we can choose it so that  $\left( \sum_{|n| \geq m} |\alpha_n - \beta_n|^r \right)^{\frac{1}{r}} < \frac{1}{M_1}$ , holds. Thus, for an arbitrarily chosen  $m \in \mathbb{Z}_+$ :

$$\theta = M_1 \left( \sum_{|n| \geq m} |\alpha_n - \beta_n|^r \right)^{\frac{1}{r}} < 1,$$

and

$$\left\| \sum_n c_n (f_n - \varphi_n) \right\|_{p, \omega} \leq \theta \left\| \sum_n c_n \varphi_n \right\|_{p, \omega}$$

holds.

Hence, by the Paley–Wiener theorem, the system  $\{f_n\}$  is an isomorphic basis to the system  $\{\varphi_n\}$  in  $N_{p, \omega}$ . Since the systems  $\{f_n\}$  and  $\{\psi_n\}$  differ from each other countable number functions, more precisely system  $\{\psi_n\}$  consist of functions

$$e^{i\mu_{-m+1}x}, e^{i\mu_{-m+2}x}, \dots, e^{i\mu_{m-2}x}, e^{i\mu_{m-1}x},$$

which are different functions in system  $\{f_n\}$ , it follows that according to the theorem, the fact that the system  $\{\varphi_n\}$  is an isomorphic basis to the system  $\{f_n\}$  is equivalent to its completeness. On the other hand, according to Theorem 2.2, which is an analogue of Levinson's theorem in  $N_{p, \omega}$  space, if in the system  $\{f_n\}$  the elements with indices  $|n| < m$  are replaced, respectively, by the elements  $\{\psi_n\}$ , then the resulting system,  $\{\psi_n\}$  will be complete in  $N_{p, \omega}$ . Therefore, the system  $\{\psi_n\}$  is an isomorphic basis to the system  $\{\varphi_n\}$  in  $N_{p, \omega}$ . The theorem is proved.

**Corollary 5.1** *Let  $1 < p < +\infty$ ,  $1 < p_0 < p$ ,  $\omega \in A_p \cap A_{p_0}$ , and*

$$r \in \left( 1, \min \left( \frac{p}{p-1}, \frac{pp_0}{p_0 + p(p_0 - 1)} \right) \right),$$

*$\{\alpha_n\}, \{\beta_n\} \subset \mathbb{R}$  is a sequence of distinct numbers and  $\sum_{n \in \mathbb{Z}} |\alpha_n - \beta_n|^r < +\infty$ . Then if the system  $\{e^{i\alpha_n x}\}_{n \in \mathbb{Z}}$  is an isomorphic basis to the system  $\{e^{i\beta_n x}\}_{n \in \mathbb{Z}}$  in space  $N_{p, \omega}$  then the system  $\{e^{i\mu_n x}\}_{n \in \mathbb{Z}}$  is an isomorphic basis to the  $\{e^{i\alpha_n x}\}_{n \in \mathbb{Z}}$ .*

**Proof.** From the condition

$$r < \frac{pp_0}{p_0 + p(p_0 - 1)},$$

we obtain

$$\frac{rp_0}{p_0 + r - rp_0} < p.$$

Hence, there exists a number  $p_1$  such that

$$\frac{rp_0}{p_0 + r - rp_0} < p_1 < p.$$

Choosing such a  $p_1$ , we have

$$\frac{rp_1}{p_1 - r} < \frac{p_0}{p_0 - 1}.$$

Thus, since  $\omega^{-1} \in L_{\frac{p_0}{p_0-1}}$ , it follows that  $\omega^{-1} \in L_{\frac{rp_1}{p_1-r}}$ . By Theorem 5.1, if the system  $\{e^{i\alpha_n x}\}$  is an isomorphic basis to  $\{e^{inx}\}$  in  $N_{p,\omega}$ , then the system  $\{e^{i\mu_n x}\}$  is also an isomorphic basis to  $\{e^{inx}\}$  in  $N_{p,\omega}$ .

The result is proved.

## References

1. Levinson, N.: *Gap and Density Theorems*, AMS. Col. Public. **26**, 1940.
2. Paley, R.E., Wiener, N.: *Fourier transforms in the complex domain*, AMS. Col. Public. **19**, 1934.
3. Sedletskii, A.M.: *Fourier transforms and approximations*, Amsterdam. Gordon and Breach Science Publ. 2000.
4. Sedletskii, A.M.: *Classes of analytic Fourier transforms and exponential approximations*, Moscow. Physmatlit 2005.
5. Young, R.M.: *An Introduction to Nonharmonic Fourier Series*, Academic Press 1980.
6. Paley, R.E., Wiener, N.: *Fourier transforms in the complex domain*, AMS. Col. Public. **19**, 1934.
7. Kadets, M.I.: *On the exact value of Paley-Wiener's constant*, Dokl. Akad. Nauk SSSR, **155**(6) (1964), 12531254. (in Russian).
8. Redheffer, R.M., Young, R.M.: *Completeness and basis properties of complex exponentials*, Trans. Amer. Math. Soc. **277**, 93–111 (1983).
9. Bilalov, B.T., Seyidova, F.Sh.: *Basicity of a system of exponents with a piecewise linear phase in Morrey-type spaces*, Turkish J. Math. **43**, 1850–1866 (2019).
10. Bilalov, B.T.: *The Basis Property of a Perturbed System of Exponentials in Morrey-Type Spaces*, Siberian Math. J. **60**(2), 249–271 (2019).
11. Bilalov, B.T., Gasymov, T.B., Guliyeva, A.A.: *On solvability of Riemann boundary value problem in Morrey-Hardy classes*, Turkish J. Math. **40**(5), 1085–1101 (2016).
12. Bilalov, B.T., Guliyeva, A.A.: *On basicity of exponential systems in Morrey-type spaces*, International Journal of Mathematics **25**, No. 6, 1450054 (2014).
13. Bilalov, B.T.: *The weighted grand Lebesgue class of harmonic functions and the Dirichlet problem*, Lobachevskii J. Math. **46**, 377–387 (2025).
14. Bilalov, B., Sadigova, S., Kasumov, Z.: *Some solvability problems of differential equations in non-standard Sobolev spaces*, In: Differential Equations (Working Title). In-techOpen, London (2022).
15. Bilalov, B., Zeren, Y., Sadigova, S., Cetin, S.: *On solvability in the small and Schauder-type estimates for higher order elliptic equations in grand Sobolev spaces (nonseparable case)*, Appl. Anal. **102**(11), 3064–3077 (2023).

16. Bilalov, B.T., Huseynli, A.A., El-Shabrawy, S.R.: *Basis properties of trigonometric systems in weighted Morrey spaces*, Azerb. J. Math. **9**(2), 166-192, (2019).
17. Huseynli, A.A.: *On the stability of basisness in  $L_p$  ( $1 < p < +\infty$ ) of cosines and sines*, Turkish J. Math. **35**, 47–54 (2011).
18. Huseynli, A.A., Mirzoyev, V.S., Quliyeva, A.A.: *On basicity of the perturbed system of exponents in Morrey-Lebesgue space*, Azerb. J. Math. **7**(2), 191-209, (2017).
19. Haqverdi, T.: *On stability of bases consisting of perturbed exponential systems in grand Lebesgue spaces*, Journal of Contemporary Applied Mathematics **11**(2), 81-92, (2021).
20. Gasymov, T.B., Ahmadli, B.Q., Behbudova, J.H.: *On strong solvability of one nonlocal boundary value problem for Poisson's equation in grand Sobolev space in rectangle*, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci. **45**(4), Mathematics, 68-75 (2025).
21. Ismailov, M.I., Hasanova, C.M., Ashirova, H.Z.: *On some properties of grand sequence spaces*, Trans. Natl. Acad. Sci. Azerb. Ser. Phys.-Tech. Math. Sci. **45**(4), Mathematics, 88-96 (2025).
22. Ismailov, M.I., Aliyarova, I.F.: *On the basis property of the system of exponentials and trigonometric systems of sine and cosine functions in weighted grand Lebesgue spaces*, Moscow Univ. Math. Bull. **79**, 85–97 (2024).
23. Bennett, C., Sharpley, R.: *Interpolation of Operators*, Academic Press, 1988, 469 p.
24. Bilalov, B.T.: *Some Problems of Approximation*, Publisher: Elm Publishing House, 2016.
25. Young, R.M.: *An Introduction to Nonharmonic Fourier Series*, Rev. 1st ed., Academic Press, San Diego, 2001.
26. Levin, B.Y.: *Distribution of Roots of Entire Functions*, Moscow, GITL, 1956 (in Russian).